



Interactive
Comment

Interactive comment on “The double layers in the plasma sheet boundary layer during magnetic reconnection” by J. Guo and B. Yu

R. Wang

rongsheng.wang@mail.iggcas.ac.cn

Received and published: 12 November 2014

The paper focused on the study of double layers (DL) during reconnection. By PIC simulations, the authors claimed that the double layers can be created during reconnection due to the beam instability in the plasma sheet boundary layer. The isolated electrostatic waves (double layers, electron holes (EH)) have been observed in previous simulations and are believed to play a key role during reconnection. Electron holes are frequently observed by the spacecraft measurement during reconnection. Double layers are frequently observed in the ionosphere. Only recently, Ergun et al. reported the double layers in the plasma sheet in the magnetotail. But, they could not confirm the relation between the double layers and reconnection in the work. This year, Wang et al. presented the first evidence of double layers during reconnection

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



[Interactive
Comment](#)

[Wang et al., GRL 2014], and found that the double layers are moving away from X-line along the separatrix region. The double layer can accelerate charged particle effectively on its way. In this paper, the simulation domain is only about $25.6 * 12.8 d_i$ which is too small to simulate the plasma boundary layer, since the plasma sheet boundary layer denotes a huge region between the lobe region and plasma sheet in the magnetotail. In my view, the so-called 'plasma sheet boundary layer' in the text corresponds to the separatrix region during reconnection. Then, the simulation results are basically consistent with the observations [Wang et al., GRL 2013; 2014]. Based on the observations, DL and EHs are observed together. EHs are observed in the high potential side of the DL. Moreover, the DL propagates in the local thermal velocity. However, the authors found that the DL almost does not move in the simulation. The authors need to provide a reasonable explanation. According to the observations, the spatial size of the DL is about Debye length. But, it reaches the ion inertial scale in the simulation. In addition, the authors claimed the DL is produced by the beam instability. Then, please show the evidence also. The language should be improved as well. The quality of the figures should be improved. In some figures, no physical quantity can be found, like Figures 3 and 4.

Please also note the supplement to this comment:

<http://www.nonlin-processes-geophys-discuss.net/1/C600/2014/npgd-1-C600-2014-supplement.pdf>

Interactive comment on Nonlin. Processes Geophys. Discuss., 1, 1657, 2014.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)