

Local Finite Time Lyapunov Exponent, Local Sampling and Probabilistic Source and Destination Regions

Amir E. BozorgMagham¹, Shane D. Ross², and David G. Schmale III³

¹Dept. of Atmospheric and Oceanic Science, University of Maryland, College Park, MD, 20742, USA

²Dept. of Biomedical Engineering and Mechanics, Virginia Tech, Blacksburg, VA 24061, USA

³Dept. of Plant Pathology, Physiology, and Weed Science, Virginia Tech, Blacksburg, VA 24061, USA

Correspondence to: Amir E. BozorgMagham (bozorgmagham@gmail.com)

Abstract. The time-varying finite time Lyapunov exponent (FTLE) is a powerful Lagrangian concept widely used for describing large-scale flow patterns and transport phenomena. However, field experiments usually have modest scales. Therefore, it is necessary to bridge between the concept of FTLE and field experiments. In this paper two observations related to approximation of the local FTLE and the distance between the destination (respectively, source) points of consecutively released (resp., collected) particles at a fixed location are presented. These two observations can practically assist in field experiments where samples are collected at a fixed location and it is desirable to attribute source locations to the collected particles. Also, results from this study may aid in planning of optimal local sampling of passive particles for maximal diversity monitoring of atmospheric assemblages of microorganisms in aerobiology. The presented observations can be used to (i) estimate the local FTLE time-series at fixed location as a function of known distances between the destination (resp., source) points and local velocity, and (ii) estimate the distances between the destination (resp., source) points of the released (resp., collected) particles when consecutive release (resp., sampling) events are performed at a fixed location. In addition to deterministic flows, the more realistic case of unresolved turbulence and low resolution flow data that yield probabilistic source (resp., destination) regions are studied. It is shown that, similar to deterministic flows, Lagrangian coherent structures (LCS) separate probabilistic source (resp., destination) regions corresponding to consecutively collected (resp., released) particles.

1 Introduction

20 The classical interpretation of finite time Lyapunov exponent (FTLE) fields and the associated hyperbolic Lagrangian coherent structures (LCSs) provides useful information about large-scale flow patterns and transport and mixing phenomena in flow domains (Haller and Poje, 1998; Haller and Yuan, 2000; Mancho et al., 2004; Shadden et al., 2005; Haller, 2011). There are an increasing number of studies that apply various concepts of LCSs, based on the classic right Cauchy-Green tensor, 25 to describe and predict the time evolution of Lagrangian features in geophysical systems (Haller, 2015). In some of these studies, geophysical information (e.g., wind or oceanic velocity fields) have been used as the input data and Lagrangian results (e.g., the distribution of an oil spill in the ocean or volcanic ash in the atmosphere) over a large area are compared with the behavior of the geophysical system via satellite data or simulations (Dellnitz et al., 2009; Peng and Peterson, 2012; Olascoaga and Haller, 2012; Mendoza and Mancho, 2012; Olascoaga et al., 2012). A large scale distribution of particles is a common characteristic among these studies. In contrast, this study is motivated by a series of field experiments and studies regarding the long distance transport of airborne microorganisms where only a limited number of localized and temporally consecutive measurements of the atmospheric structure of microbial assemblages are available (Schmale III et al., 2008; Tallapragada et al., 2011; Schmale et al., 2012; Lin et al., 2013; Schmale and Ross, 2015). Therefore, there is a need to bridge the powerful concept of FTLE and local field experiments.

In this paper, we present two observations related to the estimation of the local FTLE and the distance between destination (or source) points of released (or collected) particles, respectively. These observations are associated with the concept of local Lyapunov exponents and the direction of maximum expansion in ordinary differential equation systems (Abarbanel et al., 1992; Branicki and Wiggins, 2009). These observations may help investigate long distance transport phenomena as a possible cause of variation in successively collected airborne samples such as the presence or absence of a unique strain or species of microorganism. In addition, this analysis is useful for planning atmospheric collections at a fixed location with respect to forecast FTLE fields (BozorgMagham et al., 2013). Because this study is motivated by aerial measurements in realistic conditions, i.e., hundreds of collections of microorganisms from the atmosphere with drones, it is necessary to consider the spatiotemporal limitations of the available velocity field data. These limitations are manifested in unresolved turbulence and impose uncertainties on the location of the source and destination points. For this reason, we use a Lagrangian particle dispersion model to determine the probabilistic source 50 (or destination) regions and show how the concept of local FTLE can explain the degree of separation between probabilistic source (or destination) regions, and may contribute to understanding the degree of geographic and genetic diversity observed in aerial samples (Fay et al., 1995; Draxler and Hess, 1998; BozorgMagham and Ross, 2015).

Results from this study can be applied to environmental applications such as early warning systems for airborne pathogens, integrated pest management in crops, and the collection of samplers 55

from geophysical flows (Tallapragada et al., 2011; BozorgMagham et al., 2013; BozorgMagham and Ross, 2015).

This paper is outlined as follows. In §2 we present two observations associated with the estimation of the local FTLE and the dispersion of destination (or source) points in flow fields. In §3 we show some numerical examples and applications of presented observations in periodic and aperiodic systems. In §4 we consider the unresolved turbulence and investigate the uncertainty of the backward and forward trajectories and the resulting probabilistic source and destination regions.

2 Local Finite Time Lyapunov Exponent

In this section we present two observations related to the estimation of the local FTLE and the distance between destination (or source) points of successively released (or collected) particles in a time-varying n -dimensional vector field,

$$\frac{d\mathbf{x}}{dt} = \mathbf{v}(\mathbf{x}, t), \quad (1)$$

where $n = 2$ for two-dimensional flows and $n = 3$ for three-dimensional flows.

By local FTLE we mean the time-varying value of the FTLE field at an arbitrary location \mathbf{x} . Classically, the time-varying FTLE measures the maximum separation rate between nearby particles when they are released in the flow field at the *same* time (isochronic particles). Fig. 1(a) refers to this classical description. This figure shows two isochrone particles which are close to each other at an initial time t_0 . Under the effect of the flow field the small displacement vector between the two particles, $\delta\mathbf{x}$, changes. After an elapsed time T , the new vector between the two particles is

$$\begin{aligned} \delta\mathbf{x}(t_0 + T) &= \phi_{t_0}^{t_0+T}(\mathbf{x} + \delta\mathbf{x}) - \phi_{t_0}^{t_0+T}(\mathbf{x}) \\ &= D\phi_{t_0}^{t_0+T}(\mathbf{x}) + \mathcal{O}(\|\delta\mathbf{x}(t_0)\|^2), \end{aligned} \quad (2)$$

where $\phi_{t_0}^{t_0+T}$ is the flow map for the vector field (1) from time t_0 to t_0+T and $D\phi_{t_0}^{t_0+T} = d\phi_{t_0}^{t_0+T}(\mathbf{x})/d\mathbf{x}$ is the Jacobian of the flow map, and $\|\cdot\|$ is the usual Euclidean norm.

Consider the right Cauchy-Green strain tensor, $C(\mathbf{x}, t_0, T) = D\phi_{t_0}^{t_0+T}(\mathbf{x})^\top D\phi_{t_0}^{t_0+T}(\mathbf{x})$. For sake of the following discussion, consider the situation of incompressible two-dimensional flows, $n = 2$. The eigenvalues λ_i and normalized eigenvectors ξ_i of C satisfy (Haller, 2015),

$$C\xi_i = \lambda_i\xi_i, \quad \|\xi_i\| = 1, \quad i = 1, 2, \quad 0 < \lambda_1 < 1 < \lambda_2, \quad \xi_1 \perp \xi_2, \quad (3)$$

where the (\mathbf{x}, t_0, T) dependence of C , λ_i , and ξ_i is understood. As illustrated in Fig. 1(b), the two eigenvectors, ξ_1 and ξ_2 , are carried along by the flow $\phi_{t_0}^{t_0+T}$ to the two vectors r_1 and r_2 , respectively, where

$$r_i = D\phi_{t_0}^{t_0+T}(\mathbf{x})\xi_i, \quad (4)$$




Figure 1. (a) Separation of nearby particles during time interval T due to the flow map ϕ . The two particles are released in the flow field at the same time t_0 ; isochronic particles. (b) ξ_2 is the direction of maximum growth at the initial point \mathbf{x} , which evolves into the direction $r_2 = D\phi_{t_0}^{t_0+T}(\mathbf{x})\xi_2$ at the evolved point $\phi_{t_0}^{t_0+T}(\mathbf{x})$. The closer the initial displacement vector δx is to the ξ_2 direction, the more it will be stretched to that maximum perturbation .

85 whose lengths are scaled by a factor $\sqrt{\lambda_i}$ compared with the normalized eigenvectors. The maximum possible separation between the released particles after a time interval T , assuming a sufficiently small initial distance $\|\delta\mathbf{x}(t_0)\|$, is

$$\max \|\delta\mathbf{x}(t_0 + T)\| = \sqrt{\lambda_{\max}(C(\mathbf{x}, t_0, T))} \|\delta\mathbf{x}(t_0)\| \quad (5)$$

where $\lambda_{\max} = \lambda_n$.

The finite-time Lyapunov exponent (FTLE), with t_0 and T fixed, is considered a scalar field of the
90 Lyapunov exponent as a function of initial position, \mathbf{x} ,

$$\sigma_{t_0}^T(\mathbf{x}) = \frac{1}{|T|} \ln \sqrt{\lambda_{\max}(C)}. \quad (6)$$

Similar to the calculation of maximum separation between two initially neighboring points in a system of ordinary differential equations (ODEs) and the corresponding maximum Lyapunov exponents, $\sigma_{t_0}^T$ can be used, via (5) and (6), to describe $\max \|\delta\mathbf{x}(t_0 + T)\|$ as

$$\max \|\delta\mathbf{x}(t_0 + T)\| = \exp(\sigma_{t_0}^T(\mathbf{x}, t_0) |T|) \|\delta\mathbf{x}(t_0)\|. \quad (7)$$

In this study we are interested in particles that are released (or collected) *sequentially in time at*
95 *a fixed location*. Thus, the standard concept of the FTLE, i.e., separation rate of nearby *isochronic* points, might not be applicable. Therefore, we present two observations and show that we can (i) approximate the local FTLE by using the of the successive destination (or source) points, and (ii) estimate the distance of the destination (or source) points by having the local FTLE and velocity. These two observations require the assumption of a time-dependent vector field, so that that the
100 initial displacement vector is not along a common trajectory for sequential particles.




Figure 2. Two sequentially released (left panel)/collected (right panel) particles at a fixed location shown by a bold cross. Particles 1 and 2 are released/collected at t_1 and $t_2 = t_1 + \delta t$ respectively (time interval between the two sequential samplings is δt). The integration time between the destination/sources and the release/sampling location is approximately T for both particles ($|T| \gg \delta t$). The displacement of the first particle during δt is shown by δ^* .

Referring to Fig. 2, the left panel, assume that two particles are sequentially released at t_1 and $t_2 = t_1 + \delta t$ at the release location shown by the \times . The right panel corresponds to the analogous situation of sequentially collected particles.

Observation I: The local FTLE value over the time interval $[t_1, t_2]$, provided an appropriate $0 < \delta t \ll |T|$, can be approximated by,

$$\sigma_{[t_1, t_2]}^T(\mathbf{x}) = \frac{1}{|T|} \ln \frac{\delta(\mathbf{x}, T, t_1, \delta t)}{\|\bar{\mathbf{v}}(\mathbf{x}, t_1, t_2) \delta t\|}. \quad (8)$$

where $\delta(\mathbf{x}, T, t_1, \delta t)$ is the distance between successive destination/source points corresponding to the elapsed time T , and $\bar{\mathbf{v}}(\mathbf{x}, t_1, t_2)$ is the [non-zero] average velocity at the release/sampling location during $[t_1, t_2]$.

Observation II: The distance between the destination (resp., source) points of consecutively released (resp., collected) particles can be estimated, provided an appropriate δt , by the local velocity and the true local FTLE at the release (resp., sampling) location as,

$$\delta(\mathbf{x}, T, t_1, \delta t) = \exp(|T| \sigma_{[t_1, t_2]}^T(\mathbf{x})) \|\bar{\mathbf{v}}(\mathbf{x}, t_1, t_2) \delta t\|. \quad (9)$$

We suggest that observation I provides with a recovered FTLE field, based on mild assumptions which tend to hold in geophysical flows. Observation II is important for sampling purposes, because it enables us to estimate the distance between the source positions of consecutively collected particles, if we have the local velocity and local FTLE data by separate means. We note that observations I and II are independent and also the information on the right-hand sides of (8) and (9) are assumed known. For example, the local velocity could be obtained from an anemometer or high-frequency radar in ocean, (Shadden et al., 2009)), or the local FTLE could be obtained from a nowcast or forecast velocity field.

Two remarks are in order regarding these observations. (i) A proper choice of δt , which depends on the spatiotemporal variability of the velocity field, is critical for a good approximation of the local

FTLE or the distance between the destination (resp., source) points. If δt is chosen to be too small, then variation of the velocity field would not be observed, the two particles would essentially be following one another along a nearly identical trajectory, and consequently the separation between the two particles would lead to a null LE value, not maximal growth. However, if δt is too large, then the initial particle separation at time t_2 is too large to justify the linearization assumption underlying FTLE calculations; see, e.g., (2). Thus, a good selection of δt depends on the spatiotemporal variability of the velocity field. (ii) A larger true local FTLE of the real flow field yields a smaller error of estimations for the recovered local FTLE and the distance between the destination (resp., source) points. This comes from numerical evidence in the following sections.

The fundamental idea behind these observations is related to more general methods of analysis of chaotic dynamical systems, often used in experimental settings, namely that the direction of maximum expansion dominates the dynamics of typical displacement vector growth (see Fig. 1(b)) (Oseledec, 1968; Abarbanel et al., 1992; Rosenstein et al., 1993; Tanaka and Ross, 2009). This notion is generally accepted in settings assumed to be modeled by underlying autonomous ODEs (whether known or unknown), but to our knowledge, there is no similar theorem for non-autonomous ODEs. Our observations show that in a time-dependent velocity field, with a proper choice of δt and sufficiently large $|T|$, $\delta(\mathbf{x}, T, t_1, \delta t)$ is often close to the maximum possible distance between the two particles. These observations may help stimulate rigorous results related to separation of nearby particles in a non-autonomous ODE setting.

In following section we demonstrate some numerical verification and applications of these observations.

3 Numerical Examples and Applications

3.1 Numerical Examples of Observations I & II, for Periodic and Aperiodic Velocity Fields

First, we study the well-known example of a periodic double-gyre. We consider the same model and parameters introduced in Section 6, Example 1, of (Shadden et al., 2005). For observation I, we need to know $\delta(\mathbf{x}, T, t_1, \delta t)$ and the local velocity. Therefore, we use the double-gyre model to generate the velocity field and then exploit that data to calculate the trajectories and the corresponding distance between successively released particles after integration time T . We repeat this procedure for all the grid points of the gyre domain $[0, 2] \times [0, 1]$.

Fig. 3(a) shows the benchmark (true) forward FTLE field corresponding to $t_0 = 1$, calculated by (6), and panel (b) shows the approximated forward FTLE field calculated by (8). A 4-th order Runge-Kutta integration scheme with constant integration time-step 0.01 and total integration time $T = 15$ time units is implemented for both panels. For the recovered (approximated) FTLE field (panel (b)), we consider $\delta t = 0.2$ time unit in (8). One can adjust parameters, e.g., T or δt , to investigate their




Figure 3. Forward FTLE field of a periodic double-gyre velocity field. (a) The benchmark FTLE field at $t_0 = 1$ calculated by (6). (b) The approximated FTLE field calculated by (8), and $\delta t = 0.2$ time unit. A 4-th order Runge-Kutta integration scheme with constant integration time-step 0.01 and total integration time $T = 15$ time units is implemented for both panels.

impact on the FTLE field. Our results (not presented here) show stable outcomes when a periodic double-gyre is considered.

To investigate observation I for an aperiodic system we use a two-dimensional model of time-dependent Rayleigh-Bénard convection model developed by Solomon and Gollub (1988) and implemented by Lekien and Haller (2008) to study unsteady flow separation on slip boundaries. The streamfunction of this model is a function of position and a stochastic time-dependent forcing term.

$$\psi(x, y, t) = \frac{A}{k} \sin \{k[x - g(t)]\} \sin(2y). \quad (10)$$

Following Lekien and Haller (2008), we generate the stochastic forcing, $g(t)$, based on a random Fourier spectrum with zero mean and unit impulse covariance (see Fig. 5 in (Lekien and Haller, 2008)). Fig. 4(a) shows the benchmark forward FTLE field calculated by (6), and panel (b) shows the recovered forward FTLE field calculated by (8) in the domain $[0, 2] \times [0, \pi/2]$. Two panels of this figure correspond to $t_0 = 1$. Similar to the previous example, a 4-th order Runge-Kutta integration scheme with constant integration time-step 0.01 is used. The total integration time is $T = 75$ time units for both panels. For the recovered FTLE field (panel (b)), we consider $\delta t = 0.1$ time unit in (8).

By comparing two panels of Figs. 3 and 4, one sees that the main features of the FTLE field are recovered by (8) and the benchmark and approximated fields are highly correlated. However, in some areas (e.g., near $(1.2, 0.5)$ in Fig 3 and $(1.5, 0.8)$ in Fig 4) we see discontinuities in the recovered FTLE field. The reason might be that the selected δt is not a proper choice at those regions. It is also important to note that we use a common color scale for the two panels of Figs. 3 and 4, respectively. Therefore, minute differences between the true and approximated fields are visually exaggerated because the FTLE values are generally small in magnitude. Numerical comparison of the results (see the next two numerical experiments) shows close approximation of the recovered local FTLE to the true values.




Figure 4. Forward FTLE field of an aperiodic Rayleigh-Bénard convection model. (a) The benchmark FTLE field at $t_0=1$ calculated by (6). (b) The approximated FTLE field calculated by (8), and $\delta t = 0.1$ time unit. A 4-th order Runge-Kutta integration scheme with constant integration time-step 0.01 and total integration time $T = 75$ units is implemented for both panels.

Next, we investigate both observations I and II at an arbitrary point over the time span $[0, 10]$, which is one period of the double-gyre flow. First, we consider the point $(x, y) = (0.3, 0.4)$ in the periodic double-gyre, keeping all the parameters of FTLE computation the same as before (e.g, $\delta t = 0.2$). Fig. 5(a) shows the benchmark and approximated FTLE time-series at that point. The benchmark FTLE is calculated by (6) using the velocity field information and the maximum eigenvalue of the Cauchy-Green strain tensor. The approximated FTLE in this panel is calculated by using (8). Information about δ and the local velocity (i.e., $\bar{\mathbf{v}}$) are assumed to be known (in this numerical experiment we obtain them from the velocity field). Fig. 5(b) shows the benchmark and approximated distance between successively released particles after an elapsed time $T = 15$. To calculate the benchmark time-series we use the velocity field information to generate the trajectories and find the distance between the successive particles. The approximated time-series is generated by (9), with the provided information about the local velocity and the local FTLE value.

Next, we consider the point $(x, y) = (1.3, 1.3)$ in the aperiodic time-dependent Rayleigh-Bénard convection model, keeping all the parameters of FTLE computation the same as before (e.g, $\delta t = 0.1$). Fig. 6(a) shows the benchmark and approximated FTLE time-series at the selected point and Fig. 6(b) shows the benchmark and the approximated distance between successively released particles after an elapsed time $T = 75$.

Figures 5 and 6 show typical time-series of the recovered local FTLE and the distance between successively released (or collected) particles. As one can observe, the two time-series in panels (a) and (b) are highly correlated and the error of approximation is generally small.




Figure 5. Observation I & II at $(x, y) = (0.3, 0.4)$ for a periodic double-gyre velocity field. (a) Local benchmark and recovered forward FTLE. (b) Benchmark and approximated final distance between successively released particles.




Figure 6. Observation I & II at $(x, y) = (1.3, 1.3)$ for an aperiodic velocity field corresponding to a time-dependent Rayleigh-Bénard convection model. (a) Local benchmark and recovered forward FTLE. (b) Benchmark and approximated final distance between successively released particles.

The error of approximation in observations I and II depends on many parameters, for example: δt , T , and variation of vector field over the time-scale δt . We leave the analysis of errors of observations I and II for a future study.

3.2 Applications of the Local FTLE Observations I & II

Next, we consider the real-world wind data and focus on the backward FTLE fields and the location of source points. This situation is important for field studies for identifying potential source regions of plant pathogens and their relative risk of transport to previously unexposed regions (Lin et al.,

205 2014; Prussin et al., 2014b, a, 2015). For this purpose, we use observations I & II to compare the
benchmark and the recovered local backward FTLE time-series and also the true and the estimated
distance of the source locations corresponding to the particles that were collected at Virginia Tech’s
Kentland Farm, located at $37^{\circ}11'$ N and $80^{\circ}35'$ W. A large variety of microbial samples have been
collected at this location over the past 7 years (2006 to 2013) (Schmale et al., 2012). We refer to this
210 point as $(0, 0)$ in our plots.

The flow maps are calculated by using numerical data corresponding to the North America Mesoscale,
NAM-218 provided by the National Oceanic and Atmospheric Administration (NOAA) and National
Centers for Environmental Prediction’s (NCEP) Operational Model Archive and Distribution System
(NOMADS) project¹. Spatial resolution of this data set is about 12.1 km and the temporal resolution
215 is 3 hours. All the trajectories are calculated by a fourth order Runge-Kutta integrator with a constant
integration time step equals to 5 min. We use third order splines for all necessary spatiotemporal in-
terpolations. We consider the time interval 12:00 UTC 29 Sep to 12:00 UTC 30 Sep 2010 for our
numerical experiments and refer to it as the *interrogation window*.

Figs. 7 shows the trajectories and the initial positions of the indexed particles corresponding to
220 the collected particles at the sampling location during the interrogation window. The frequency of
sampling was 1 hour and the backward time integration is 24 hours for all the particles. In addition,
for simplicity and without losing generality of the results we perform the integration on a quasi-
2D 850mb pressure surface (BozorgMagham and Ross, 2015). Indices of this figure indicate the
sampling times of the collected particles, for example index ‘12’ that locates on the North-West of
225 the figure refers to the initial position of a particle that started at 12:00 UTC 28 Sep and was collected
24 hours later, i.e., 12:00 UTC 29 Sep, at the sampling location. In terms of streaklines (Batchelor,
2000), this line (Fig. 7(b)) is composed of contemporaneous points, e.g., 24 hr, from the assembly
of streaklines which pass through the sampling location during the interrogation window. We define
this line as the *isochrone source-line* since the integration time from all points on it to the sampling
230 location is the same, i.e., 24 hr in this example.

Following the assumptions of the local FTLE observations, i.e., a proper δt with respect to the
spatiotemporal variability of the velocity field, we choose sampling periods from 0.1 to 1 hour and
all the integration are done in the same interrogation window. Figure 8(a) shows the benchmark
distance between successive source points, i.e., $\delta(\mathbf{x}, T, t_1, t_2)$, during the interrogation window cal-
235 culated from the available velocity field data. We use the average velocity at the sampling location
to calculate δ^* as $\|\bar{\mathbf{v}}(\mathbf{x}, t_1, t_2) \delta t\|$. Fig. 8(b) shows the recovered local FTLE time-series, assuming
that the true successive distances are available.

Figures 7(b) and 8 demonstrate that we interpret a *local (backward) FTLE time-series as differ-*
ential stretching of line elements, i.e., strain, along an isochrone source-line. To verify this result
240 and to study the effect of different δt ’s on the recovery of local FTLE time-series we calculate the

¹<http://nomads.ncdc.noaa.gov/data.php>




Figure 7. (a) Trajectories of the collected particles during 24 hours of integration. (b) Sequential source points and the isochrone source-line. Sampling frequency is one hour between 12:00 UTC 29 Sep to 12:00 UTC 30 Sep 2010 and the sampling location is at (0,0) (Virginia Tech Kentland Farm 37° 11' N and 80° 35' W).




Figure 8. (a) δ as the benchmark (true) distance between successive source points corresponding to $\delta t = 0.1, 0.5$ and 1 hr. Horizontal axis represents the averaged time corresponding to each successive pairs. (b) Approximated local FTLE for different δt 's from 0.1 hr (6 min) to 1 hour. Interrogation window is 12:00 UTC 29 Sep to 12:00 UTC 30 Sep 2010.

benchmark backward FTLE fields for the interrogation window with integration time equal to 24 hr. Figure 9(a) shows an image of the true time-varying FTLE field corresponding to 12:00 UTC 29 Sep 2010. To give a sense about the changes of the FTLE field during the interrogation window, we may describe the motion of the strong ridges of the field in figure 9(a) toward North-West direction, as shown by the arrow. Figure 9(b) shows the benchmark local FTLE value (black line) at the Kentland Farm during the interrogation window. To generate this plot we exploit (6) and calculate the backward FTLE field every 15 minutes, then the time-varying value of FTLE at (0,0) is extracted.




Figure 9. (a) The frozen image corresponding to 12:00 UCT 29 Sep 2010 of the backward FTLE field during the interrogation window. Integration time is 24 hours for FTLE calculations. The bold arrow shows the general wind direction and the motion of the attracting LCS. (b) The true (black) and recovered (red) local FTLE time-series at the reference point (0,0). For the recovered time series (red), δt is equal to 0.1 hr.

Also, for comparing the results, the recovered FTLE time-series corresponding to $\delta t = 0.1$ hr is displayed in the same panel by the red line. Figures 8(b) and 9(b) indicate that: (i) an optimal δt for this example is between 0.1 to 0.5 hr. (ii) The estimation error is smaller for larger values of the true local FTLE. Therefore, we may observe larger errors of estimation when [true] σ is close to zero, e.g., between 00 and 4 UTC in Fig. 9(b). For $\delta t = 0.1$ hr we observe that the true and approximated local FTLE time-series are highly correlated and also their maxima (corresponding to the local maxima of the FTLE field) are at the same times (within $\delta t = \pm 0.1$ hr). Therefore, with a proper choice of δt 's the recovered local FTLE time-series can accurately capture the passage times of moving ridges of a FTLE field. Detecting these ridges is important since they are candidates for hyperbolic LCSs in many geophysical applications (Tallapragada et al., 2011; Haller, 2011; Karrasch, 2012; BozorgMagham et al., 2013).

In addition, we investigate whether we can estimate the distances by using observation II, providing necessary information about local velocity and FTLE. Fig. 10 is a numerical example that shows that the benchmark distance between the successive source points (black line) is well approximated (red line) by observation II, i.e., (9). Note that in this case we have the data of the true local FTLE and the local velocity. In this figure we see that at $\delta t = 0.25$ hr, the estimated differential distance time-series is very close to the true answer and it captures the correct times of the local maxima.

This is an empirically important result, because one can schedule the sampling of geophysical flows (e.g., with drones) based on the available forecast FTLE fields and local velocity such that the successive collected particles originate from the most possible diverse locations (see § 3.2). In Fig. 10 it is evident that there are two optimal time intervals, i.e., before and after 16 UTC, for maximal diversity monitoring. To interpret this, consider Fig. 7(b) and notice that the geographic




Figure 10. Differential distance between the successive source points on the isochrone source-line corresponding to $\delta t = 0.25$ hr. The black line shows the benchmark and the red line shows the approximated time series which is calculated by local FTLE theorem as $\exp(|T|\sigma_{[t_1, t_2]}^T(\mathbf{x}))\|\bar{\mathbf{v}}(\mathbf{x}, t_1, t_2)\delta t\|$. The backward integration time for calculations of the flow maps is $T = 24$ hr and the interrogation window is 12:00 UTC 29 Sep to 12:00 UTC 30 Sep 2010.

270 extent of the line segment from point 15 to point 16 is much larger than the segment from point 13 to point 14.

A direct result of the local FTLE observations is the possibility of planning for maximal geographic (and therefore also genetic) diversity monitoring such that the collected particles come from the most separated source locations. This means incorporating greater potential source areas, which
 275 could drive a greater diversity of sample collection. Suppose that it is desired to maximize the genetic diversity of microorganisms collected in a sample, assuming that all the collected particles have approximately the same flight time. Results of observation II indicate that the optimal time for collecting samples such that they originate from the most possible distant locations is at times corresponding to the maxima of the local FTLE time-series (note the high correlation between the distance and the local FTLE time-series in Fig. 10 and Fig. 9(b)). To ensure that the particles are coming from
 280 significantly separated locations we may use the topology of the FTLE field and collect the samples on either side of a strong attracting LCS feature which corresponds to a local maximum of $\sigma_{[t_1, t_2]}^T$, providing short enough time between sampling periods. In this condition, high value of $\sigma_{[t_1, t_2]}^T$ as the exponent in equation (7) is the reason for having a large δ . Figure 11 schematically shows this
 285 strategy when an attracting LCS feature passes over a fixed sampling location causing a dramatic change in the region of possible source points of collected particles.




Figure 11. An attracting LCS feature (red) passes over the geographically fixed sampling location (indicated by a bold \times). Dashed lines show trajectories of hypothetical particles that are absorbed to a moving attracting LCS. The bold arrow shows the general wind direction and the motion of the attracting LCS at the specified interval. Collected samples on either side of this attracting LCS feature come from two different regions.

As an example in realistic geophysical flow, Fig. 12 shows trajectories of three hypothetical particles that are collected at $(0, -100)$ km with respect to the reference point. Backward integration time for specifying the corresponding source points, i.e., A, B and C, and the trajectories is 40 hr
 290 for those three particles. The sampling times during the interrogation window are 13:40 UTC for the red particle, 14:00 UTC for the blue particle and 14:10 UTC for the green particle. The green and the blue particles are collected on one side of an attracting LCS but the red particle is collected on the other side of the same LCS. As we observe, the source points corresponding to blue and green particles, points B and C, are close. Meanwhile the source point of the red particle, point A, is significantly far from the other two particles. An interesting feature of this figure is that the separation of
 295 the trajectories does not start from the sampling point, but as it is shown, the three trajectories remain close to each other for about 200 km and then begin to diverge. This observation is directly related to the concept of the FTLE, because $\sigma_{t_0}^T$ is a function of the “final” separation between nearby particles and it does not specify the moment of divergence.

300 Referring to this example, observation II can help us to explain the seeming association of sample diversity with high FTLE. There have been some reports of significant characteristic variation of the collected particles, e.g., genetic types or aerial density of the microbial samples, during short intervals when sampling coincides with a high value local FTLE, or similarly, passage of a strong LCS over the sampling location (Schmale et al., 2012; Lin et al., 2013). In addition, a direct result of the
 305 local FTLE observations is that when the local FTLE value is small during the sampling process, it is expected that the collected particles originate from nearby source points, assuming approximately




Figure 12. Three trajectories of hypothetical collected samples. The red and blue trajectories correspond to the samples on either side of a LCS. The blue and green trajectories correspond to the samples on one side of the same LCS. Sampling times are 13:40, 14:00 and 14:10 UTC during the interrogation window (12:00 UTC 29 Sep to 12:00 UTC 30 Sep 2010) for the red, blue and the green particles, respectively. Source points of the collected particles are shown by A, B and C. Integration time for all three particles is $T = 40$ hr.

same flight times for them. This might be the reason that the characteristics of the microbial samples remain quasi-constant in consecutive collections, but differ as the time between sample collections increases (Lin et al., 2013). This situation is similar to sampling from a *coherent set* where the FTLE values are generally small (Froyland et al., 2010; Tallapragada and Ross, 2013) and the particles have similar Lagrangian characteristics. Moreover, in cases where we observe significant changes in collected samples while local FTLE value is small, we speculate that those changes are caused by local sources rather than long range transport phenomena (Lin et al., 2013). Thus, the local FTLE concept helps us to include or exclude rare/unique microbes from specific source regions. This sets the stage for additional work to be performed to test hypotheses concerning the presence/absence of the unique microbes at the potential source locations.

4 Unresolved Turbulence and Probabilistic Regions

In this section we study the uncertainty in calculation of the source (or destination) points due to the unresolved turbulence and also the role of high-value local FTLE in separation of the probabilistic source (or destination) regions.

Precise calculation of the source (or destination) point of any collected (or released) particle and the corresponding flow map require high resolution data of the velocity field. But geophysical data

are always discrete and spatially sparse. For example, spatial and temporal resolution of operational data sets vary from the order of 10 to hundreds of kilometers and 3 hours to longer intervals respectively. Meanwhile, spatiotemporal scales of atmospheric flows can be smaller than the resolution of the available data and we may lose important Lagrangian phenomena such as turbulent diffusion and small size eddies if we just consider available data (Csanady, 1973; Rodean, 1996). Therefore, for realistic calculation of the source (or destination) points it is necessary to consider the uncertainty of the trajectories. For this purpose, we consider a Lagrangian particle dispersion model (LPDM) that provides the stochastic component of the velocity with respect to the available deterministic (background) data (Legg and Raupach, 1982; Fay et al., 1995; Draxler and Hess, 1998; Stohl et al., 2005). In LPDM, the velocity vector at each point, $\mathbf{v}(\mathbf{x}, t)$, is assumed to be the summation of a deterministic term, $\bar{\mathbf{v}}(\mathbf{x}, t)$, and a random variable, $\mathbf{V}(\mathbf{x}, \bar{\mathbf{v}}, t)$, which depends explicitly on the instantaneous position of the particle \mathbf{x} , its deterministic velocity $\bar{\mathbf{v}}$ at that location and the time t ; see equation (11). Later, we see how this dependency dictates two different solutions for the calculations of the probabilistic source and destination regions (BozorgMagham and Ross, 2015).

$$\mathbf{v}(\mathbf{x}, t) = \bar{\mathbf{v}}(\mathbf{x}, t) + \mathbf{V}(\mathbf{x}, \bar{\mathbf{v}}, t). \quad (11)$$

The stochastic term of (11) is a Markov-chain process as a function of the velocity deformation tensor and the Lagrangian time scale of the flow field,

$$V_{(t+\Delta t)} = R_{\Delta t} V_t + (1 - R_{\Delta t}^2)^{0.5} \mathcal{N}(0, 1) \sqrt{\kappa/T_L} \quad (12)$$

where V shows each component of the stochastic velocity term \mathbf{V} , and the correlation coefficient $R_{\Delta t}$ is a measure of the association between stochastic velocities during successive time steps. Also, \mathcal{N} is a normal distribution with mean zero and unit standard deviation. The correlation coefficient

$$R_{\Delta t} = \exp(-\Delta t/T_L), \quad (13)$$

is a function of integration time step, Δt , and the Lagrangian time scale of the flow field, T_L , which is on the order of 10^4 s. The term κ depends on the gradient of the instantaneous deterministic velocity, $\bar{\mathbf{v}} = (\bar{u}, \bar{v})$, the grid size of the meteorological data, χ , and an empirical constant, c ,

$$\kappa = 2^{-0.5} (c\chi)^2 \left[\left(\frac{\partial \bar{v}}{\partial x} + \frac{\partial \bar{u}}{\partial y} \right)^2 + \left(\frac{\partial \bar{u}}{\partial x} - \frac{\partial \bar{v}}{\partial y} \right)^2 \right]^{0.5}. \quad (14)$$

Because κ depends on the gradient of the background velocity one can easily use the set of equations (11)-(14) for *forward* integration. Using this set for simple backward integration requires presumption about the position of a particle at specific times which leads to misleading results. Therefore, we have to consider two distinct cases, (i) calculation of the probabilistic destination region of a released particle, (ii) calculation of the probabilistic source region of a collected particle. In this study we discuss both cases, but like before, emphasize the probabilistic source regions (corresponding to the backward trajectories). We also revisit the problem of local FTLE and successive sampling

in the presence of unresolved turbulence. Our numerical results show that even in the presence of unresolved turbulence, if successive sampling are performed on either side of a strong attracting LCS, represented by the temporal peaks in the local backward FTLE time-series, then the probabilistic source regions are significantly separated similar to the deterministic case.

To focus on the main concerns of this study and to avoid complexity we proceed with a 2D velocity field similar to the previous sections. However, this approach can be extended to 3D fields by adding an appropriate stochastic term in the extended direction (Rodean, 1996).

4.1 Probabilistic Source and Destination Regions

(I) The probabilistic destination region is the probability distribution of the final positions of virtually released particles after integration time T when the initial position is known precisely, e.g., a Dirac delta function. The case of forward integration and related calculations of a probabilistic distribution is equivalent to solving the Fokker-Planck or Kolmogorov forward equations (Rodean, 1996; Risken, 1985) which describes the future of a probability distribution function of a known initial condition that evolves under the dynamics of a system, e.g., a diffusion process.

Because the time-varying vector fields are usually complicated, analytical solutions for probabilistic destination regions are not available and it is necessary to use numerical solutions. For this aim, we discretize the domain of our interest into sufficiently small boxes and then use the Monte Carlo method by releasing sufficient number of independent particles from a box that includes the release point. Fig. 13(a) shows this procedure. By choosing an appropriate integration time step we calculate the trajectories. By completion of the integration process we have a distribution of particles in different boxes. If the total number of released particles is sufficiently large and the boxes' dimensions are sufficiently small, then the ratio of the virtual particles in each box to the total number of released particles show the probability distribution of the destination region. By increasing the number of virtual particles and decreasing the size of the boxes the calculated distribution becomes invariant.

(II) The solution for a probabilistic source region is conceptually the same as solving the Kolmogorov backward problem (Risen, 1985). In mathematical terms, at time $t_0 - T$ ($T > 0$ is the integration time) we investigate for a specific source distribution such that in a future time, i.e., t_0 , the system will be in a given target set. A probabilistic source region cannot be determined by simply performing backward time-integration. Because κ in equation (14) and consequently the stochastic velocity term are determined by the instantaneous background velocity which depends on the location and time. Naively applying the backward time-integration produces a series of "false" displacement vectors. The cumulative effect of these false displacements yields a false probabilistic source region. To solve this problem, we first discretize the domain of the flow field into small boxes. Then, we shift the starting time to $t_0 - T$ and consider the velocity field at this new time frame. By this means, we convert this problem into a forward integration problem from $t_0 - T$ to t_0 . At $t_0 - T$




Figure 13. (a) A solution for the probability distribution of a forward case. Virtual particles are released from a box that includes the release location. Distribution of the final positions after integration time T would specify the probabilistic destination region. Calculation of the probabilistic destination region is equivalent to the solution of a Fokker-Planck equation for finding the future probability distribution of an initially known distribution. Trajectories of the released particles from the initial box are shown in green. (b) A solution for probability distribution of a source region. For a proper [forward-time] integration, the starting time is shifted to $t_0 - T$. Virtual particles are released from *all* the boxes in the domain. Important particles are those who land in the target box which include the sampling location. Trajectories of particles which land in the target box are shown by green, other trajectories are shown by red. A solution for the probabilistic source region is conceptually the same as the solution of backward Kolmogorov equation where an initial probability distribution is the desired solution such that in a future time the system will have a specified probability distribution.

we release a sufficient number of independent particles from *all* boxes of the domain (this step is the major difference between the current and the previous case). By forward integration from $t_0 - T$ to time t_0 we find the landing location of all released particles. The influential particles in this procedure are those which land inside the sampling box, e.g., the green trajectories in Fig. 13(b). In this figure, those boxes that have contribute to the particles ending up in the target box “ j ”, are hatched. As we observe, there may be particles from contributing boxes that do not land in the target box (shown by red trajectories).

In Fig. 13(b) the boxes are labeled by $i = 1, 2, \dots, n_b$ where n_b is the number of boxes and the sampling box is shown by index j . We denote the number of particles which start from box i at time $t_0 - T$ and are in box j at time t_0 by $n_{i \rightarrow j}$. We calculate the relative contribution of each source box as,

$$\gamma_i = \frac{n_{i \rightarrow j}}{\sum_i n_{i \rightarrow j}}, \quad (15)$$

where $\sum_i n_{i \rightarrow j}$ shows the total number of particles that land in the sampling (target) box j and γ_i is the chance of a collected particle to come from a specific box i . Therefore, the distribution of γ over the domain approximates the probability distribution of the source region. This procedure

generates the correct probabilistic source region but its numerical efficiency is not high because many, e.g., 10^6 , independent particles are released from all boxes of the domain but only those particles that land in the sampling box are counted. Thus, there are a huge number of calculated trajectories that are left out. It is not the purpose of this study, but one can increase the efficiency of this procedure by applying some optimization methods. For example, sequential release of particles from large boxes that are inside a circle centered at the sampling box and by identifying the regions with maximum contributions. The radius of that circle can be determined by statistical information about the mean velocity and the integration time. After that, one may focus on those important regions by partitioning them into smaller boxes and increasing the number of released particles to determine fine structures of the probabilistic source region. For more information regarding this problem one can refer to STILT project² (Lin et al., 2003; Nehrkorn et al., 2010; Hegarty et al., 2013).

4.2 Probabilistic Source Region and Local FTLE Observations

To investigate a realistic example of probabilistic source regions and the applicability of the presented observations, we revisit the case study of section §3.2. Fig. 14(a) and(b) show one example of a probabilistic source region where the color intensity shows the relative contribution of each source box. In this case the sampling location is at $(0, -100)$ km with respect to our reference point. Sampling time is 14:10 UTC 29 Sep 2010 and the total elapsed time for trajectory calculations is $T = 40$ hours. This figure is the stochastic equivalent of the source point of the particle that its trajectory is shown by the green line in Fig. 12. For this calculation 10^5 particles are released from each 10×10 km box. After trial and error experiment, the search area for this specific problem is considered to be $900 \text{ km} \times 600 \text{ km}$ rectangular grid. Considering the size of the boxes, the total number of released particles and calculated trajectories is 5.4×10^8 in each integration time step.

An important point for probabilistic source and destination regions, is that although at each time step the stochastic velocity term has a Gaussian distribution (recalling equation (12)), the final distribution of particles is not necessarily Gaussian. The reason for this fact is the cumulative effects of the variability of the variance of normal distribution, $\sqrt{\kappa/T_L}$, that is a function of the gradient of instantaneous velocity. In general, for small integration time the probability distribution of the source (or destination) region is close to a Gaussian distribution but as the integration time increases, the corresponding distribution diverges from a normal one. For example, visual inspection of Fig. 14b indicates that the final distribution of the probable source points is not Gaussian. In Fig. 14c the relative contribution of the source boxes along the specified line PQ is shown. Standard statistical tests such as the Kolmogorov-Smirnov test (Lilliefors, 1967) confirm that the distribution is not Gaussian.

In Fig. 12 we show that the source locations of two collected particles on either side of a attracting LCS are much further apart than the source points of two successive collected particles on one side of

²<http://www.stilt-model.org>




Figure 14. (a) The probabilistic equivalent of the source point of the green particle in Fig. 12. The sampling point S is located at $(0, -100)$ km with respect to our reference point and sampling time is 14:10 UTC 29 Sep 2010. (b) Details of the probabilistic source region which is composed of 5400 boxes, each $10\text{ km} \times 10\text{ km}$. Color intensity shows the relative contribution of each source box. (c) γ , relative contribution of source boxes along the specified line PQ.

the same LCS. We want to investigate whether this result is still valid in the presence of unresolved turbulence. If that result holds then in practical applications such as sampling the microbial structure of the atmosphere we can have reasonable confidence about the separation of the probabilistic source regions based solely on a deterministic analysis, that is, without performing bothersome probabilistic calculations. Therefore, we study a case that we know its deterministic dynamics. Fig. 15 shows the evolution of the probabilistic source regions “A” and “B” (shown in panel (a)) corresponding to red and blue particles of Fig. 12 respectively. The total integration time for this example is 40 hours. In each panel of this figure we also show the attracting hyperbolic LCSs according to Haller (2011) and Karrasch (2012). For calculation of each probabilistic region of this figure, 10^5 particles are released from each small $10\text{ km} \times 10\text{ km}$ box. By comparing Fig. 12 and Fig. 15(a) we observe that the probabilistic source regions contain the deterministic source points and they significantly separated from each other. Also, we see how the two probabilistic regions contract and become closer to the attracting LCS as they get closer to the sampling point. One noticeable feature in this figure is the difference between the shapes of the two source regions, while the two samplings are separated by only 20 minutes.




Figure 15. Sequence of hyperbolic LCSs (blue) and two probabilistic source regions corresponding to two successive samples. Probabilistic regions “A” and “B” (panel (a)) correspond to the virtually red and blue particles in Fig. 12. These six panels correspond to 40, 30, 20, 10, 5 and 0 hours before collecting the corresponding samples at 13:40 and 14:00 UTC during the interrogation window.

Results of this example show that, similar to the consequences of observation II, the probabilistic source regions corresponding to the collected particles on either side of a deterministic attracting LCSs are significantly separated in backward time.

455 5 Conclusions

FTLE fields provide useful information about large-scale transport phenomena and also Lagrangian structures of flow fields, particularly geophysical flows. However, in field experiments the data are on a much more modest scale. Therefore, it is necessary to bridge the gap between the concept of large-scale FTLE fields and local experiments. Our new observations were motivated in part by our
 460 previous work examining the dynamics of assemblages of microorganisms in the lower atmosphere.

We present two observations. (i) The local FTLE can be approximated at a given spatial location, if one knows the local velocity and tracks the distance between sequentially released or collected particles. (ii) One can estimate the distances between the destination (or source) points of sequentially released (or collected) particles from a given spatial location assuming the availability of the
 465 local velocity and local FTLE.

We numerically demonstrate the results of our observations for a periodic velocity field (i.e., a double-gyre), an aperiodic system (i.e., a Rayleigh-Bénard convection model) and the real-world wind data. The suggested notion is useful in practical cases where we have samples of particles (e.g., microbes) collected at a fixed location and we are interested in formulating hypotheses about their origin, structure, and potential transport phenomena driving their atmospheric movement. In addition, we show that the concept of local FTLE and observation II can be applied to scheduling of atmospheric sampling missions to collect samples containing unique or high diversity individuals or species.

We also investigate the unresolved turbulence and the probabilistic description of the source (or destination) points. We use the box discretization method and discuss the important differences between calculation methods of the probabilistic source and destination regions. Furthermore, we show that because the stochastic velocity is a function of instantaneous background velocity, the probabilistic source (or destination) regions are not necessarily Gaussian.

Finally, we study the probabilistic source regions corresponding to successive collected particles on either side of a strong hyperbolic attracting LCS (or equivalently, a local maximum of the local FTLE time-series) and show that the source regions are significantly separated, similar to what was found in the deterministic flow field.

Results of this study can aid in optimizing the sampling schedules of passive particles and also aid in understanding of the outcomes of local observations in geophysical flows, based on large-scale transport features.

Acknowledgements. This material is based upon work supported by the National Science Foundation under Grant Numbers CMMI-1100263 (Dynamical Mechanisms Influencing the Population Structure of Airborne Pathogens: Theory and Observations) and CMMI-1150456 (Integrating Geometric, Probabilistic, and Topological Methods for Phase Space Transport in Dynamical Systems). Part of this research was performed during a visit by SDR to Instituto de Ciencias Matemáticas, Madrid Spain. He thanks ICMAT for its hospitality and support from MINECO: ICMAT Severo Ochoa project SEV-2011-0087.

References

- Abarbanel, H. D., Brown, R., and Kennel, M. B.: Local Lyapunov exponents computed from observed data, *Journal of Nonlinear Science*, 2, 343–365, 1992.
- 495 Batchelor, G. K.: *An Introduction to Fluid Dynamics*, Cambridge University Press, 2000.
- BozorgMagham, A. E. and Ross, S. D.: Atmospheric Lagrangian coherent structures considering unresolved turbulence and forecast uncertainty, *Communications in Nonlinear Science and Numerical Simulation*, 22, 964–979, 2015.
- BozorgMagham, A. E., Ross, S. D., and Schmale, D. G.: Real-time Prediction of Atmospheric Lagrangian
500 Coherent Structures Based on Uncertain Forecast Data: An Application and Error Analysis, *Physica D*, 258, 47–60, 2013.
- Branicki, M. and Wiggins, S.: Finite-time Lagrangian transport analysis: stable and unstable manifolds of hyperbolic trajectories and finite-time Lyapunov exponents, arXiv preprint arXiv:0908.1129, 2009.
- Csanady, G. T.: *Turbulent Diffusion in the Environment*, vol. 3, Springer, 1973.
- 505 Dellnitz, M., Froyland, G., Horenkamp, C., Padberg-Gehle, K., and Sen Gupta, A.: Seasonal variability of the subpolar gyres in the Southern Ocean: a numerical investigation based on transfer operators, *Nonlinear Processes in Geophysics*, 16, 655–663, 2009.
- Draxler, R. and Hess, G.: An overview of the HYSPLIT_4 modelling system for trajectories, dispersion and deposition, *Australian meteorological magazine*, 47, 295–308, 1998.
- 510 Fay, B., Glaab, H., Jacobsen, I., and Schrodin, R.: Evaluation of Eulerian AND Lagrangian atmospheric transport models at the Deutscher-wetterdienst using ANATEX surface tracer data, *Atmospheric environment*, 29, 2485–2497, 1995.
- Froyland, G., Lloyd, S., and Santitissadeekorn, N.: Coherent sets for nonautonomous dynamical systems, *Physica D: Nonlinear Phenomena*, 239, 1527–1541, 2010.
- 515 Haller, G.: A variational theory of hyperbolic Lagrangian Coherent Structures, *Physica D*, 240, 574–598, 2011.
- Haller, G.: Lagrangian coherent structures, *Annual Review of Fluid Mechanics*, 47, 137–162, 2015.
- Haller, G. and Poje, A.: Finite time transport in aperiodic flows, *Physica D*, 119, 352–380, 1998.
- Haller, G. and Yuan, G.: Lagrangian coherent structures and mixing in two-dimensional turbulence, *Physica D*, 147, 352–370, 2000.
- 520 Hegarty, J., Draxler, R. R., Stein, A. F., Brioude, J., Mountain, M., Eluszkiewicz, J., Nehr Korn, T., Ngan, F., and Andrews, A.: Evaluation of Lagrangian particle dispersion models with measurements from controlled tracer releases, *Journal of Applied Meteorology and Climatology*, 52, 2623–2637, 2013.
- Karrasch, D.: Comment on “A variational theory of hyperbolic Lagrangian coherent structures, *Physica D* 240 (2011) 574-598”, *Physica D*, 241, 1470–1473, 2012.
- 525 Legg, B. and Raupach, M.: Markov-chain simulation of particle dispersion in inhomogeneous flows - the mean-drift velocity induced by a gradient in Eulerian velocity variance, *Boundary-Layer Meteorology*, 24, 3–13, 1982.
- Lekien, F. and Haller, G.: Unsteady flow separation on slip boundaries, *Physics of Fluids (1994-present)*, 20, 097101, 2008.
- 530 Lilliefors, H. W.: On the Kolmogorov-Smirnov test for normality with mean and variance unknown, *Journal of the American Statistical Association*, 62, 399–402, 1967.

- Lin, B., BozorgMagham, A. E., Ross, S. D., and Schmale, D. G.: Small fluctuations in the recovery of fusaria across consecutive sampling intervals with unmanned aircraft 100 m above ground level, *Aerobiologia*, 29, 45–54, 2013.
- 535 Lin, B., Ross, S. D., Prussin, A. J., and Schmale, D. G.: Seasonal associations and atmospheric transport distances of fungi in the genus *Fusarium* collected with unmanned aerial vehicles and ground-based sampling devices, *Atmospheric Environment*, 94, 385–391, 2014.
- Lin, J., Gerbig, C., Wofsy, S., Andrews, A., Daube, B., Davis, K., and Grainger, C.: A near-field tool for simulating the upstream influence of atmospheric observations: The Stochastic Time-Inverted Lagrangian Transport (STILT) model, *Journal of Geophysical Research: Atmospheres* (1984–2012), 108, 2003.
- 540 Mancho, A., Small, D., and Wiggins, S.: Computation of hyperbolic trajectories and their stable and unstable manifolds for oceanographic flows represented as data sets, *Nonlinear Processes in Geophysics*, 11, 17–33, 2004.
- Mendoza, C. and Mancho, A. M.: Review Article: "The Lagrangian description of aperiodic flows: a case study of the Kuroshio Current", *Nonlinear Processes in Geophysics*, 19, 449–472, 2012.
- 545 Nehr Korn, T., Eluszkiewicz, J., Wofsy, S. C., Lin, J. C., Gerbig, C., Longo, M., and Freitas, S.: Coupled weather research and forecasting–stochastic time-inverted lagrangian transport (WRF–STILT) model, *Meteorology and Atmospheric Physics*, 107, 51–64, 2010.
- Olascoaga, M., Brown, M., Beron-Vera, F., and Koçak, H.: Brief communication "Stratospheric winds, transport barriers and the 2011 Arctic ozone hole", *Nonlinear Processes in Geophysics*, 19, 687–692, 2012.
- 550 Olascoaga, M. J. and Haller, G.: Forecasting sudden changes in environmental pollution patterns, *PNAS*, 109, 4738–4743, 2012.
- Oseledec, V. I.: A multiplicative ergodic theorem. Lyapunov characteristic numbers for dynamical systems, *Trans. Moscow Math. Soc.*, 19, 197–231, 1968.
- 555 Peng, J. and Peterson, R.: Attracting structures in volcanic ash transport, *Atmospheric Environment*, 48, 230–239, 2012.
- Prussin, A. J., Li, Q., Malla, R., Ross, S. D., and Schmale, D. G.: Monitoring the long distance transport of *Fusarium graminearum* from field-scale sources of inoculum, *Plant Disease*, 98, 504–511, 2014a.
- Prussin, A. J., Szanyi, N. A., Welling, P. I., Ross, S. D., and Schmale, D. G.: Estimating the production and release of ascospores from a field-scale source of *Fusarium graminearum* inoculum, *Plant Disease*, 98, 497–503, 2014b.
- 560 Prussin, A. J., Marr, L. C., Schmale, D. G., Stoll, R., and Ross, S. D.: Experimental validation of a long-distance transport model for plant pathogens: application to *Fusarium graminearum*, *Agricultural and Forest Meteorology*, 203, 118–130, 2015.
- 565 Risken, H.: *The Fokker-Planck equation: methods of solution and applications*, Springer-Verlag, 1985.
- Rodean, H. C.: *Stochastic Lagrangian models of turbulent diffusion*, *Meteorological monographs*, 26, 1–84, 1996.
- Rosenstein, M. T., Collins, J. J., and De Luca, C. J.: A practical method for calculating largest Lyapunov exponents from small data sets, *Physica D: Nonlinear Phenomena*, 65, 117–134, 1993.
- 570 Schmale, D. and Ross, S.: Highways in the Sky: Scales of Atmospheric Transport of Plant Pathogens, *Annual Review of Phytopathology*, 53, 591–611, 2015.

- Schmale, D. G., Ross, S. D., Feters, T., Tallapragada, P., Wood-Jones, A., and Dingus, B.: Isolates of *Fusarium graminearum* collected 40 to 320 meters above ground level cause Fusarium head blight in wheat and produce trichothecene mycotoxins, *Aerobiologia*, 28, 1–11, 2012.
- 575 Schmale III, D. G., Dingus, B. R., and Reinholtz, C.: Development and application of an autonomous unmanned aerial vehicle for precise aerobiological sampling above agricultural fields, *Journal of Field Robotics*, 25, 133–147, 2008.
- Shadden, S. C., Lekien, F., and Marsden, J. E.: Definition and properties of Lagrangian coherent structures from finite-time Lyapunov exponents in two-dimensional aperiodic flows, *Physica D*, 212, 271–304, 2005.
- 580 Shadden, S. C., Lekien, F., Paduan, J. D., Chavez, F. P., and Marsden, J. E.: The correlation between surface drifters and coherent structures based on high-frequency radar in Monterey Bay, *Deep Sea Research Part II: Topical Studies in Oceanography*, 56, 161–172, 2009.
- Solomon, T. and Gollub, J. P.: Chaotic particle transport in time-dependent Rayleigh-Benard convection, *Physical Review A*, 38, 6280, 1988.
- 585 Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note: The Lagrangian particle dispersion model FLEXPART version 6.2, *Atmospheric chemistry and physics*, 5, 2461–2474, 2005.
- Tallapragada, P. and Ross, S. D.: A set oriented definition of finite-time Lyapunov exponents and coherent sets, *Communications in Nonlinear Science and Numerical Simulation*, 18, 1106–1126, 2013.
- Tallapragada, P., Ross, S. D., and Schmale, D. G.: Lagrangian coherent structures are associated with fluctuations in airborne microbial populations, *Chaos*, 21, 033 122, 2011.
- 590 Tanaka, M. L. and Ross, S. D.: Separatrices and basins of stability from time series data: an application to biodynamics, *Nonlinear Dynamics*, 58, 1–21, 2009.