

Limiting amplitudes of fully nonlinear interfacial tides and solitons

Borja Aguiar-González^{1,2} and Theo Gerkema³

¹Departamento de Física, Facultad de Ciencias del Mar, Universidad de Las Palmas de Gran Canaria, E-35017 Las Palmas, Spain.

²NIOZ Royal Netherlands Institute for Sea Research, Department of Ocean Systems Sciences and Utrecht University, P.O. Box 59, 1790 AB Den Burg, Texel, the Netherlands.

³NIOZ Royal Netherlands Institute for Sea Research, Department of Estuarine and Delta Systems, and Utrecht University, P.O. Box 140, 4400 AC Yerseke, the Netherlands.

Correspondence to: B. Aguiar-González
(aguiar@nioz.nl)

Abstract. A new two-fluid layer model consisting of forced rotation-modified Boussinesq equations is derived for studying tidally generated fully nonlinear, weakly nonhydrostatic dispersive interfacial waves. This set is a generalization of the Choi-Camassa equations, extended here with forcing terms and Coriolis effects. The forcing is represented by a horizontally oscillating sill, mimicking a barotropic tidal flow over topography. Solitons are generated by a disintegration of the interfacial tide. Because of strong non-linearity, solitons may attain a limiting table-shaped form, in accordance with soliton theory. More generally, we use the model equations to investigate the role of the initial stages of the internal tide on limiting the amplitudes of solitons under fully nonlinear conditions. Numerical solutions reveal that the internal tide, considered linear but with the inclusion of barotropic advection (the quasi-linear case), reaches a limiting amplitude under increasing barotropic forcing. This extends to the nonlinear case. In particular, internal solitons formed by a disintegration of the internal tide may not reach their table-shaped form with increased forcing because of the limitation in the underlying internal tide itself.

1 Introduction

tidally generated internal solitons are a widespread phenomenon in the oceans and they have been observed and studied for decades (see, e.g., Apel et al. (2006)). They are intrinsically linked to the internal tide, which itself is generated by barotropic tidal flow over topography. As the internal tide steepens, it may split up into groups of internal solitons, which therefore appear at the tidal period.

20 For internal solitons as such, an archetypal model has been the Korteweg-de Vries (KdV) equation,
which is based on the assumption of weak nonlinearity and weak nonhydrostatic effects. The equa-
tion gives prediction for the relation between amplitude, width and phase speed of the solitons, as
well as the shape itself. In the KdV equation there is, mathematically speaking, no limit to the ampli-
25 tude that solitons may reach (although, of course, at some point the underlying assumption of weak
nonlinearity would be violated). This behaviour changes fundamentally if a higher-order (i.e., cubic)
nonlinear term is included, leading to the so-called extended KdV (eKdV) equation, as discussed in,
e.g., Helfrich and Melville (2006). This extended version produces qualitatively different solitons:
their amplitude is limited (for a given configuration of layers) and they broaden as they reach their
maximum amplitude, the so-called ‘table-top’ solitons. This behaviour is confirmed by the fully
30 nonlinear soliton models, as derived by Choi and Camassa (1999) and Miyata (1985, 1988) (denoted
as the MCC equations for brevity).

In this paper, we focus on another limiting factor, which comes into play even before solitons arise,
namely in the internal tide itself. In a purely linear system, the amplitude of the internal tide increases
35 linearly with the strength of the barotropic tidal flow. Here we study how this changes when, at a
next stage, one includes quasi-linear terms, i.e. retaining products of barotropic and baroclinic fields
in the advective terms while still ignoring interactions of the baroclinic field with itself. We demon-
strate that a saturation in the amplitude of the internal tide occurs and increasing the barotropic flow
further does not produce a larger internal tide. As a consequence, when one includes the genuinely
40 nonlinear effects, i. e. products of baroclinic terms, resulting solitons may stay well below their
formal limiting amplitude, no matter how strong the forcing.

To study these effects we derived a set of fully nonlinear, weakly nonhydrostatic model equations,
by extending the MCC equations with a barotropic tidal forcing over topography and with Coriolis
45 effects, which have previously been shown to play a key role in soliton generation from internal
tides (Gerkema and Zimmerman, 1995). To avoid nonlinearities in the barotropic flow itself (which
cannot be formally neglected in a fully nonlinear model), we mimic the generation by barotropic
tidal flow over topography with a horizontally oscillating topography. (There is no complete equiv-
alence, but we demonstrate that for the parameters used here, the difference remains small.) The
50 presence of a topography greatly complicates the subsequent handling of the equation, necessary to
bring them in a form amenable to numerical solving, but we demonstrate that the set of equations
can be obtained.

An extension of the MCC theory with Coriolis effects (MCC- f) was already derived by Helfrich
55 (2007), who investigated on the decay and return of internal solitary waves with rotation. We focus
on the novel aspect of studying the wave evolution and limiting amplitudes of fully nonlinear, weakly

nonhydrostatic internal tides and solitons when a forcing and rotational effects are added. We denote our extension of the MCC theory as forced-MCC- f (forced-MCC in absence of rotation), for brevity.

60 The paper is organized as follows. We derive a new two-fluid layer model consisting of a set of forced rotation-modified Boussinesq equations in Sect. 2. We start with the basic equations and assumptions. Then, we scale equations (Sect. 2.1) and vertically integrated them over the layers (Sect. 2.2). Up to this point, the resulting equations are exact but do not form a closed set. The set is closed by making an expansion in a small parameter measuring the strength of non-hydrostaticity
65 (Sect. 2.3). The resulting model turns out equivalent to Choi-Camassa equations plus additional terms which provide the forcing and rotation effects to the system. Prior to discussion of the numerical experiments, we address in Sect. 3 some preliminaries related to the oscillating topography, the governing nondimensional parameters and the actual environmental parameters used for the runs. In Sect. 4 we investigate the factors limiting the growth of tidally generated solitons by firstly ap-
70 proaching the generation of quasi-linear internal tides within the parameter space of study. Next, in Sect. 5 we solve the full set of forced-MCC- f and explore the conditions by which tide-generated fully nonlinear solitons may actually attain a limiting amplitude. A discussion of main findings and conclusions are presented in Sect. 6.

75 The numerical methods and schemes are described in *Appendix A*. The full set of model equations as solved in the code is presented in *Appendix B* together with its (quasi)-linearization form. In *Appendix C* we evaluate, within the parameter space of study, the oscillating topography against a generation model which works with tidal flow over a topography at rest.

80 2 Derivation of the forced-MCC- f model

We start from the continuity and Euler equations and consider a two-fluid layer system (Fig. 1) with a jump in density across the interface and in which each layer is composed of a homogeneous, inviscid, and incompressible fluid, applying the Boussinesq approximation. We also assume uniformity in one of the horizontal directions, taking $\partial/\partial y = 0$. Hence, the continuity and momentum
85 equations read

$$u_{i,x} + w_{i,z} = 0 \quad (1)$$

$$\bar{\rho} \left(u_{i,t} + u_i u_{i,x} + w_i u_{i,z} - f v_i \right) = -p_{i,x} \quad (2)$$

$$v_{i,t} + u_i v_{i,x} + w_i v_{i,z} + f u_i = 0 \quad (3)$$

$$\bar{\rho} \left(w_{i,t} + u_i w_{i,x} + w_i w_{i,z} \right) = -p_{i,z} - \rho_i g \quad (4)$$

90 where ρ_i is density, (u_i, v_i, w_i) are the velocity components in Cartesian coordinates, p_i is pressure, g the gravitational acceleration, f the Coriolis parameter ($f = 2\Omega\sin\phi$, at latitude ϕ) and $\bar{\rho}$ the mean density. The subscript $i = 1$ ($i = 2$) refers to the upper (lower) layer and a stable stratification, $\rho_1 < \rho_2$, is assumed.




Fig. 1. The two-fluid layer system for which the forced-MCC- f equations are derived. The horizontal dashed grey line indicates the level $z = 0$; the level at which the interface resides at rest.

95 Boundaries are defined at the surface, taken to be a rigid lid, which is located at $z = H_1$, and at the bottom, located at $z = -H_2 + H(x, t)$. The time-dependence of the bottom will later be specified as a horizontal oscillation, mimicking a barotropic tidal flow over topography, the forcing mechanism for internal tides. However, the two are not exactly equivalent, since the transformation from one frame of reference to the other involves an acceleration and is therefore not Galilean. We further
 100 discuss this aspect in *Appendix C*.

The kinematic boundary conditions at the surface and bottom read

$$w_1 = 0 \quad \text{at } z = H_1 \quad (5)$$

$$w_2 = H_t + H_x u_2 \quad \text{at } z = -H_2 + H(x, t). \quad (6)$$

105 At the interface, $z = Z(x, t)$, which if at rest is located at $z = 0$, the boundary conditions are given by the continuity of normal velocity and pressure:

$$w_i = Z_t + u_i Z_x \text{ and } p_1 = p_2 \quad \text{at } z = Z. \quad (7)$$

For later convenience, we write pressure as the sum of hydrostatic and dynamic parts, the latter being denoted by primes:

110 $p_i = \rho_i g H_1 - \rho_i g z + p'_i(t, x, z).$

In the horizontal momentum equation, this amounts to replacing $p_{i,x}$ by $p'_{i,x}$, whereas the vertical momentum equation (4) gives

$$\bar{\rho} (w_{i,t} + u_i w_{i,x} + w_i w_{i,z}) = -p'_{i,z}.$$

Continuity of pressure at the interface, the second equation in (7), now becomes

$$115 \quad (p'_1 - p'_2)|_{z=Z} = (\rho_1 - \rho_2)gZ.$$

2.1 Scaling

The next step is to bring the equations into an appropriate dimensionless form for which we introduce the following scales. The scale for the undisturbed water depth is taken to be D , and the typical wavelength L . Crucially, we will assume waves to be long, i.e. nonhydrostatic effects to be weak.

$$120 \quad \text{This will be expressed by the small parameter}^1, \delta = \left(\frac{D}{L}\right)^2 \ll 1.$$

Since we allow waves to have large amplitudes (i.e. being strongly nonlinear), we may take horizontal current velocities to scale with $c_0 = (g'D)^{1/2}$, where g' is reduced gravity, $g' = g(\rho_2 - \rho_1)/\bar{\rho}$; and c_0 is close to the linear long-wave phase speed for interfacial waves, c_p (which would have

125 $H_1 H_2 / D$ instead of D). Thus, u and v will be scaled with c_0 . For the interfacial displacement being allowed to be large, an appropriate scale of Z is D .

The typical scale of w now follows from the continuity equation as Dc_0/L . Finally, the scale of pressure follows from assuming a primary balance between the acceleration term $\bar{\rho} u_t$ and p_x in the

130 horizontal momentum equation.

In summary, then, we can introduce the following dimensionless variables, indicated by asterisks,

$$\begin{aligned} x &= L x^*, & z &= D z^*, & t &= (L/c_0) t^*, \\ p'_i &= (\bar{\rho} c_0^2) p'^*_i, & u_i &= c_0 u^*_i, & v_i &= c_0 v^*_i, & w_i &= (D/L) c_0 w^*_i. \end{aligned} \quad (8)$$

With these scales, the dimensionless continuity and Euler equations yield (for convenience, we drop

135 the asterisks right away):

$$u_{i,x} + w_{i,z} = 0 \quad (9)$$

$$u_{i,t} + u_i u_{i,x} + w_i u_{i,z} - \mu v_i = -p'_{i,x} \quad (10)$$

$$v_{i,t} + u_i v_{i,x} + w_i v_{i,z} + \mu u_i = 0 \quad (11)$$

$$\delta \left(w_{i,t} + u_i w_{i,x} + w_i w_{i,z} \right) = -p'_{i,z}. \quad (12)$$

140 Here μ is the scaled Coriolis parameter, $\mu = fL/c_0$. Furthermore we introduce the dimensionless quantities ζ , h_i , and h via $(Z, H_1, H_2, H) = D(\zeta, h_1, h_2, h)$, so that the scaled form of the boundary

¹In Choi and Camassa (1999) a small parameter ϵ was used instead, which relates to ours as $\delta = \epsilon^2$.

conditions is

$$w_1 = 0 \quad \text{at } z = h_1 \quad (13)$$

$$w_i = \zeta_t + u_i \zeta_x \quad \text{at } z = \zeta(x, t) \quad (14)$$

$$145 \quad p'_2 - p'_1 = \zeta \quad \text{at } z = \zeta(x, t) \quad (15)$$

$$w_2 = h_t + u_2 h_x \quad \text{at } z = -h_2 + h(x, t) . \quad (16)$$

The goal is now to derive a reduced set of equations from (9)–(12), in which the boundary conditions (13)–(16) are incorporated by vertical integration, exploiting the smallness of the parameter δ . The procedure is identical to that of (Choi and Camassa, 1999), but with the additional complications of
150 the Coriolis force, topography, and tidal forcing.

2.2 Vertically integrated equations

We vertically integrate the equations over the upper and lower layers and expand them to the orders δ^0 and δ^1 to obtain a closed set for the weakly nonhydrostatic equations, following Choi and Camassa (1999). The layer-mean \bar{f}_1 of a function $f_1(x, z, t)$ for the upper layer is being defined as

$$155 \quad \bar{f}_1(x, t) = \frac{1}{\eta_1} \int_{\zeta}^{h_1} dz f_1(x, z, t), \quad \eta_1 = h_1 - \zeta \quad (17)$$

and for the lower layer as

$$\bar{f}_2(x, t) = \frac{1}{\eta_2} \int_{-h_2+h}^{\zeta} dz f_2(x, z, t), \quad \eta_2 = h_2 - h + \zeta . \quad (18)$$

where η_i represents the thickness of the layer (depending on the interfacial displacement ζ). Notice that the boundaries of the integral depend on time and space (x) via the interfacial movement $\zeta(t, x)$,
160 but also, for the lower layer, via the horizontally oscillating topography², $h(t, x)$. Before proceeding, nonlinear terms in horizontal momentum equations (10) and (11) are rewritten as $(u_i^2)_x + (w_i u_i)_z$ and $(u_i v_i)_x + (w_i v_i)_z$, respectively, to facilitate the procedure.

After integration of Eqs. (9)–(11) for $i = 1$ and applying the boundary conditions (13)–(15) we
165 obtain the layer-mean equations for the upper layer

$$\eta_{1,t} + (\eta_1 \bar{u}_1)_x = 0, \quad (19)$$

$$(\eta_1 \bar{u}_1)_t + (\eta_1 \overline{u_1 u_1})_x - \mu \eta_1 \bar{v}_1 = -\overline{\eta_1 p'_{1,x}}, \quad (20)$$

$$170 \quad (\eta_1 \bar{v}_1)_t + (\eta_1 \overline{u_1 v_1})_x + \mu \eta_1 \bar{u}_1 = 0. \quad (21)$$

²For this reason we need to apply the Leibniz integral rule below with respect to x and t .

For the lower layer one proceeds similarly, except that now both boundaries are variable. Applying the boundary conditions (14)–(16), vertical integration of (9)–(11) for $i = 2$ yields

$$\eta_{2,t} + (\eta_2 \bar{u}_2)_x = 0, \quad (22)$$

$$175 \quad (\eta_2 \bar{u}_2)_t + (\eta_2 \overline{u_2 u_2})_x - \mu \eta_2 \bar{v}_2 = -\eta_2 \overline{p'_{2,x}}, \quad (23)$$

$$(\eta_2 \bar{v}_2)_t + (\eta_2 \overline{u_2 v_2})_x + \mu \eta_2 \bar{u}_2 = 0. \quad (24)$$

2.3 Expansion in δ

The six integrated equations (19)–(24) derived so far are exact but do not form a closed set. The variables η_1 , η_2 and ζ count as one unknown, but we have also \bar{u}_i , \bar{v}_i , $\overline{p'_{i,x}}$, $\overline{u_i u_i}$ and $\overline{u_i v_i}$, giving 11 unknowns for 6 equations. To obtain a closed set, the last two expressions will be cast in terms of \bar{u}_i and \bar{v}_i by using the vertical momentum equation, expanded in terms of the small parameter δ . Furthermore, continuity of pressure at the interface is used to connect the pressure in the lower and upper layer (i.e., $\overline{p'_{1,x}}$ and $\overline{p'_{2,x}}$). All in all, the six equations are thus modified to contain only six unknowns. With this aim, we make a formal expansion of the unknowns for the lowest (δ^0) and next (δ) orders, as, for example:

$$\bar{f}_i = \bar{f}_i^{(0)} + \delta \bar{f}_i^{(1)} + \dots$$

At the lowest order (δ^0), $p'^{(0)}$ accounts for hydrostatic effects. At the next order (δ), $p'^{(1)}$ brings weakly nonhydrostatic effects into the system.

190

2.3.1 Lowest order

At lowest order, the vertical momentum equation (12) reduces to $\partial p_i'^{(0)}/\partial z = 0$ as terms of order δ are neglected; therefore, (perturbation) pressure is vertically constant in each layer. For convenience, we introduce $P = p_2'^{(0)}$, being a function of t and x . It then follows from continuity of pressure at the interface, that $p_1'^{(0)} = P - \zeta$. Thus, to this order of approximation,

195

$$\overline{p'_{1,x}} = P_x - \zeta_x + O(\delta), \quad (25)$$

and for the lower layer

$$\overline{p'_{2,x}} = P_x + O(\delta). \quad (26)$$

Given the z -independence of pressure and returning to the original horizontal momentum equations, it is now natural to assume that the horizontal velocities, too, are independent of z within each layer:

200

$$\overline{u_i u_i} = \bar{u}_i^2 + O(\delta), \quad \overline{u_i v_i} = \bar{u}_i \bar{v}_i + O(\delta).$$

At lowest order, then, the set of integrated equations is closed; together with the (exact) integrated continuity equations (19) and (22), we have the momentum equations in terms of the six variables $\bar{u}_i, \bar{v}_i, \zeta$ and P :

$$205 \quad (\eta_1 \bar{u}_1)_t + (\eta_1 \bar{u}_1^2)_x - \mu \eta_1 \bar{v}_1 = -\eta_1 (P_x - \zeta_x) + O(\delta), \quad (27)$$

$$(\eta_2 \bar{u}_2)_t + (\eta_2 \bar{u}_2^2)_x - \mu \eta_2 \bar{v}_2 = -\eta_2 P_x + O(\delta), \quad (28)$$

$$(\eta_1 \bar{v}_1)_t + (\eta_1 \bar{u}_1 \bar{v}_1)_x + \mu \eta_1 \bar{u}_1 = O(\delta), \quad (29)$$

$$(\eta_2 \bar{v}_2)_t + (\eta_2 \bar{u}_2 \bar{v}_2)_x + \mu \eta_2 \bar{u}_2 = O(\delta). \quad (30)$$

Recall that $\eta_{1,2}$ can be expressed in terms of ζ and thus involve just one unknown.

210

2.3.2 Next order

To include terms of order δ , the key problem is, again, to close the set of six vertically integrated equations by deriving closed expressions for the horizontal pressure gradients $\overline{p'_{i,x}}$ as well as for the contributions of $\overline{u_i u_i}$ and $\overline{u_i v_i}$ in the nonlinear terms. The latter problem is particularly simple. At

215 order δ , the products contain one lowest-order field, which is independent of z (e.g., $u_i^{(0)} = \bar{u}_i^{(0)}$), hence

$$\begin{aligned} \overline{u_i u_i} &= \frac{1}{\eta_i} \int dz u_i^2 = \frac{1}{\eta_i} \int dz (u_i^{(0)2} + 2\delta u_i^{(0)} u_i^{(1)} + \dots) \\ &= \bar{u}_i^{(0)2} + 2\delta \bar{u}_i^{(0)} \bar{u}_i^{(1)} + \dots \\ &= \bar{u}_i^2 + O(\delta^2) \end{aligned}$$

220 so that

$$\overline{u_i u_i} = \bar{u}_i^2 + O(\delta^2), \quad \overline{u_i v_i} = \bar{u}_i \bar{v}_i + O(\delta^2).$$

This allows us to write the horizontal momentum equations as

$$(\eta_1 \bar{u}_1)_t + (\eta_1 \bar{u}_1^2)_x - \mu \eta_1 \bar{v}_1 = -\eta_1 \overline{(p_1^{(0)} + \delta p_1^{(1)})_x} + O(\delta^2) \quad (31)$$

$$(\eta_2 \bar{u}_2)_t + (\eta_2 \bar{u}_2^2)_x - \mu \eta_2 \bar{v}_2 = -\eta_2 \overline{(p_2^{(0)} + \delta p_2^{(1)})_x} + O(\delta^2) \quad (32)$$

$$225 \quad (\eta_1 \bar{v}_1)_t + (\eta_1 \bar{u}_1 \bar{v}_1)_x + \mu \eta_1 \bar{u}_1 = O(\delta^2) \quad (33)$$

$$(\eta_2 \bar{v}_2)_t + (\eta_2 \bar{u}_2 \bar{v}_2)_x + \mu \eta_2 \bar{u}_2 = O(\delta^2) \quad (34)$$

The remaining problem is to find an expression for $p_i^{(1)}$. At order δ , Eq. (12) reads in terms of the lowest order vertical velocities,

$$w_i^{(0)}{}_t + u_i^{(0)} w_i^{(0)}{}_x + w_i^{(0)} w_i^{(0)}{}_z = -p_i^{(1)}{}_z \quad (35)$$

230 From vertically integrating the continuity equation (9), we obtain an expression for $w_i^{(0)}$:

$$w_i^{(0)} = -z \bar{u}_{i,x}^{(0)} + c_i(t, x)$$

where c_i are ‘constants’ of integration which are determined by using the boundary conditions at the surface (13) and bottom (16). Thus, $w_i^{(0)}$ for the upper- and lower layers become, respectively,

$$w_1^{(0)} = (h_1 - z) \bar{u}_{1,x}^{(0)}, \quad (36)$$

$$235 \quad w_2^{(0)} = (h - h_2 - z) \bar{u}_{2,x}^{(0)} + D_2 h, \quad (37)$$

where the operator D_i is defined as $\partial/\partial t + \bar{u}_i^{(0)}\partial/\partial x$. Substituting $w_1^{(0)}$ from Eq. (36) and $w_2^{(0)}$ from Eq. (37) into Eq. (35), and vertically integrating the result, we get an expression for $p_1^{(1)}$ and $p_2^{(1)}$. Taking their derivative with respect to x and their mean over each layer, we finally obtain an expression for $\overline{p_i^{(1)x}}$ at the upper and lower layer at order δ . Including the lowest order terms (25)

240 and (26), this allow us to write the horizontal pressure gradient for the upper layer

$$\overline{p'_{1,x}} = \overline{p'_{1,x}{}^{(0)}} + \delta \overline{p'_{1,x}{}^{(1)}} + O(\delta^2) = P_x - \zeta_x - \delta \left[\frac{1}{3\eta_1} (\eta_1^3 G_1)_x \right] + O(\delta^2), \quad (38)$$

and, for the lower layer,

$$\overline{p'_{2,x}} = \overline{p'_{2,x}{}^{(0)}} + \delta \overline{p'_{2,x}{}^{(1)}} + O(\delta^2) = P_x - \delta \left[\frac{1}{3\eta_2} (\eta_2^3 G_2)_x + \frac{1}{2} \eta_2 G_2 h_x - \frac{\eta_2}{2} (D_2^2 h)_x - \zeta_x D_2^2 h \right] + O(\delta^2), \quad (39)$$

245 where we introduced for simplicity the term G_i (as in Choi and Camassa (1999)),

$$G_i = \bar{u}_{i,xt}^{(0)} + \bar{u}_i^{(0)} \bar{u}_{i,xx}^{(0)} - (\bar{u}_{i,x}^{(0)})^2. \quad (40)$$

With this, the horizontal momentum equations (31) and (32) become

$$(\eta_1 \bar{u}_1)_t + (\eta_1 \bar{u}_1^2)_x - \mu \eta_1 \bar{v}_1 = -\eta_1 \left\{ P_x - \zeta_x - \delta \left[\frac{1}{3\eta_1} (\eta_1^3 G_1)_x \right] \right\} + O(\delta^2) \quad (41)$$

$$(\eta_2 \bar{u}_2)_t + (\eta_2 \bar{u}_2^2)_x - \mu \eta_2 \bar{v}_2 = -\eta_2 \left\{ P_x - \delta \left[\frac{1}{3\eta_2} (\eta_2^3 G_2)_x + \frac{1}{2} \eta_2 G_2 h_x - \frac{\eta_2}{2} (D_2^2 h)_x - \zeta_x D_2^2 h \right] \right\} + O(\delta^2) \quad (42)$$

250

We have thus obtained a closed set of six dimensionless equations, namely the exact continuity equations (19) and (22), the horizontal momentum equations (41) and (42), as well as (33) and (34); the last four equations involve the the weakly non-hydrostatic assumption. The six unknowns are \bar{u}_1 , \bar{u}_2 , \bar{v}_1 , \bar{v}_2 , P , and (via $\eta_{1,2}$) ζ . In the absence of an interfacial wave forcing and neglecting Earth's rotation effects, our set of equations reduces to that of Choi and Camassa (1999).

255

Before proceeding to numerical solving, we further specify the model by prescribing the oscillating topography, i.e., the forcing to the system, with

$$260 \quad h = h(X) \quad \text{with} \quad X(x, t) = x - U_0 \cos t \quad (U_0 \text{ being an arbitrary positive constant}). \quad (43)$$

We combine the continuity equations (19) and (22) into

$$(\eta_1 + \eta_2)_t + (\eta_1 \bar{u}_1 + \eta_2 \bar{u}_2)_x = 0, \quad (44)$$

Given that $\eta_1 + \eta_2 = h_1 + h_2 - h$, with the two-fluid system depth $h_1 + h_2 = 1$, this leads to

$$-h_t + (\eta_1 \bar{u}_1 + \eta_2 \bar{u}_2)_x = 0. \quad (45)$$

265 If we now substitute the time derivative of the oscillating topography (43) above, it yields

$$(\eta_1 \bar{u}_1 + \eta_2 \bar{u}_2)_x = U \frac{\partial h}{\partial x}, \quad (46)$$

with

$$U = U_0 \sin t, \quad (47)$$

which mimicks a barotropic tidal flow over the oscillating topography (i. e. the velocity of the
270 moving topography), where U_0 represents its dimensionless velocity amplitude. Then, Eq. (46) can
be integrated in x ,

$$\eta_1 \bar{u}_1 + \eta_2 \bar{u}_2 = Uh + C(t). \quad (48)$$

Far from the sill (i.e., $h \rightarrow 0$ for $x \rightarrow \pm\infty$), we impose the flow to be purely baroclinic, so that the
left-hand side must be zero and hence it follows that $C(t) = 0$. Notice that the right-hand side is
275 prescribed via the forcing and is thus a known quantity. It allows us to express \bar{u}_2 in terms of \bar{u}_1 .

We can thus combine the horizontal momentum equations (41) and (42), eliminating P ,

$$\begin{aligned} \bar{u}_{1,t} + \bar{u}_1 \bar{u}_{1,x} + \mu \bar{v}_1 = \zeta_x + \frac{1}{(1-h)} \left((Uh)_t + (\eta_1 \bar{u}_1^2 + \eta_2 \bar{u}_2^2)_x - \mu(\eta_1 \bar{v}_1 + \eta_2 \bar{v}_2) - \eta_1 \zeta_x \right) + \\ \delta \left(1 - \frac{\eta_1}{(1-h)} \right) \left[\eta_1 G_1 \eta_{1,x} + \frac{\eta_1^2}{3} G_{1,x} \right] \\ 280 + \frac{\delta \eta_2}{(1-h)} \left[-\eta_2 G_2 \zeta_x - \frac{\eta_2^2}{3} G_{2,x} + \frac{\eta_2 G_2}{2} h_x + \frac{\eta_2}{2} (D_2^2 h)_x + \zeta_x D_2^2 h \right] + O(\delta^2) \end{aligned} \quad (49)$$

$$\bar{u}_2 = \frac{Uh - \eta_1 \bar{u}_1}{\eta_2}, \quad (50)$$

$$\bar{v}_{1,t} + \bar{u}_1 \bar{v}_{1,x} + \mu \bar{u}_1 = 0 + O(\delta^2), \quad (51)$$

$$\bar{v}_{2,t} + \bar{u}_2 \bar{v}_{2,x} + \mu \bar{u}_2 = 0 + O(\delta^2), \quad (52)$$

$$285 \quad \zeta_t - (h_1 - \zeta) \bar{u}_{1,x} + \bar{u}_1 \zeta_x = 0. \quad (53)$$

where the \bar{v}_i -horizontal momentum equations (51) and (52) have been further simplified from (33)
and (34) by using the continuity equations (19) and (22). Eq. (19) has now been expressed in terms
of ζ for convenience. The other continuity equation (22) is no longer included explicitly since it is
already present via (50).

290

All in all, we have now five equations for five unknowns (\bar{u}_1 , \bar{u}_2 , \bar{v}_1 , \bar{v}_2 and ζ). The numerical
methods and schemes used to solve the model are described in *Appendix A*. The actual form of the
model equations as used in the numerical code is presented in *Appendix B*.

295 **3 Numerical experiments: Preliminaries**

Whilst not designed to represent a specific region of the world oceans, we aim to investigate in a general manner the conditions by which tidally generated solitons may evolve and, eventually, develop limiting amplitudes in ocean-like scenarios. It is then desirable that leading solitons can propagate towards a mature stage before overtaking preceding internal tides; otherwise, although being form-
 300 preserving features, the tracking of their wave properties becomes cumbersome. For this reason the environmental parameters that we describe in the following were selected to highlight the qualitative features of these nonlinear processes for a broad range of (mimicked) tidal forcing strength.

Although the model is solved in nondimensional form, we will discuss results from the numerical
 305 experiments with a dual view, dimensional/dimensionless, to ease the visualization of the ocean-like magnitudes being used.

3.1 The oscillating topography and the hydraulic state: the Froude number

We define the (dimensional) topography analytically following:

310
$$H(X) = \frac{H_T}{1 + (x/H_L)^2} \tag{54}$$

with x being the grid positions in space; and, H_T and H_L being the dimensional parameters which set the height and width of a symmetric sill, respectively. This manner we ensure perfectly smooth second and third derivatives of the dimensionless topography $h(X)$ in the model equations. Other analytical functions may be also used depending on the desired topography.

315

At this point it is worth while to recall that the oscillation of the topography is introduced in dimensionless form as $h = h(X)$ with $X(x, t) = x - U_0 \cos t$, where U_0 prescribes the strength (velocity amplitude) of the oscillating topography via $U = U_0 \sin(t)$, the mimicked barotropic tidal flow (see (43)–(47)). By increasing U_0 we enhance the forcing via U , which in dimensional form we introduce, respectively, as $\mathbf{U}_0 = c_0 U_0$ and $\mathbf{U} = c_0 U$.
 320

The topographic obstacle (ridge, sill, ...) is always centred in the x -axis and the length of the x -domain is chosen to be large enough to prevent waves from reaching the boundaries. In all experiments, fluid starts moving to the right at $t = 0$ (i.e., topography moving to the left); we start with
 325 a system at rest, i.e., $U = \bar{u}_1 = \bar{u}_2 = 0$ at $t = 0$. The waves are generated near the origin in x -axis due to the ‘tide-topography’ interaction; on the negative (positive) x -axis, waves travel to the left (right). Because the forcing enters in the simulation asymmetrically with fluid at rest moving to the right, it is expected that wave packets in the front appear rather different when comparing both sides (negative vs. positive x -domain). These fronts are the transients, which are influenced by the way

330 the experiment is started. A steady solution at both sides of the x -axis is reached after several tidal periods have passed away. In this regard, and to avoid transient effects generated at the start of each run, wave properties have been tracked systematically over the third leftward-propagating interfacial wave counting from the front, and after 9 tidal periods of forcing.

335 To characterize the hydraulic state where interfacial waves propagate we use the Froude number calculated as follows:

$$Fr = \frac{\mathbf{U}_0}{c_p} \quad (55)$$

where the velocity amplitude of the mimicked tidal flow acting as external forcing, \mathbf{U}_0 , is confronted to the linear long-wave phase speed for interfacial waves, c_p . The strength of \mathbf{U}_0 leads to
 340 three different regimes of interfacial wave generation (see e.g. in Vlasenko et al. (2005); Da Silva et al. (2015)). The hydraulic regime is *subcritical* ($Fr \ll 1$) when the strength of the external forcing is much less than the phase speed of the generated interfacial waves, which propagate as harmonic first-mode baroclinic interfacial waves (linear theory). When the phase speed of the generated waves is of the order of the external forcing, the flow is *critical* ($Fr \approx 1$). Then, nonlinear effects become
 345 evident and the linear baroclinic interfacial waves disintegrate into short nonlinear interfacial waves. A further increase of the external forcing leads to the most crucial regime when $Fr > 1$. Then, the flow becomes supercritical and, typically, a packet of strongly nonlinear short interfacial waves, or solitons, is generated from the longer quasi-linear interfacial wave. These solitons eventually may attain a ‘table-top’ form, the limiting scenario subject of this study. To account for the varying
 350 strength of the tidal forcing within a tidal cycle we introduce the instantaneous Froude number, defined as $Fr' = \mathbf{U}/c_p$.

Importantly, we also use the Froude number in *Appendix C* to discuss the applicability of our ‘non-inertial’ frame of reference, the oscillating topography, to the ocean case, where the topography is at
 355 rest. To this aim we compare the generation of interfacial waves from the (quasi-) linearized version the forced-MCC equations and the (quasi-) linearized version of the weakly nonlinear model derived in Gerkema (1996) (*GI996*), which works with actual tidal motion. The quasi-nonlinear case involves in both generation models neglecting the baroclinic interactions but retaining the nonlinear terms involving a combination of barotropic and baroclinic fields. The equations are then still linear
 360 with regard to the baroclinic fields, but the coefficients become time-dependent due to barotropic factors (which are prescribed), so that higher harmonics will be generated when the forcing is increased. For clarification, the (quasi-) linearization of the forced-MCC- f equations is presented in *Appendix B*.

365 Results from the above model intercomparison supports a semi-equivalence between both interfacial wave generation mechanisms within the framework of study, which we restrict to $0 < Fr < 1.6$

based on the cited analysis. This semi-equivalence encourages us to discuss our numerical results, henceforth, referring to the strength of the oscillating topography, U_0 , as the strength of the tidal flow. Similarly, we will refer to interfacial waves generated from the forced-MCC- f equations as internal tides.

3.2 Environmental parameters of study

We adopt a two-fluid layer system where the total water depth, D , is set to 100 m with the upper layer being always thinner than the lower layer ($H_1 < H_2$). The horizontal oscillation of the moving topography is always of semidiurnal frequency. Although the height of the topography varies between runs, its horizontal scale is kept constant and about 20 km ($H_L = 10$ km in (54)).

In Table 1 the varying environmental parameters are detailed. These vary between runs as indicated in underlined bold fonts, one at a time. The theoretical amplitude of the ‘table-top’ soliton predicted from Eq. (3.68) in Choi and Camassa (1999), and beyond which no solitary wave solution exists, is also indicated.

Comparison between runs A1, A2 and A3 account for the effect of varying the stratification via the reduced gravity, g' . Comparison between runs A1, B1 and B2 account for the effect of varying the topography ratio, $\varphi_T = H_T/D$, which measures the height of the topography relative to the total water depth. Lastly, comparison between runs A1, C1 and C2 account for the effect of varying the two-fluid layer thickness ratio, $\gamma = H_1/H_2$.

Run	g' (in m s^{-2})	$\varphi_T = H_T/D$	$\gamma = H_1/H_2$	$-A_m/H_1$	H_1, H_2 (in meters)	ρ_1, ρ_2 (kg m^{-3})
A1	<u>0.03</u>	0.4	0.43	0.67	30, 70	1022, 1025.15
A2	<u>0.02</u>	0.4	0.43	0.67	30, 70	1023.05, 1025.15
A3	<u>0.01</u>	0.4	0.43	0.67	30, 70	1024.1, 1025.15
B1	0.03	<u>0.35</u>	0.43	0.67	30, 70	1022, 1025.15
B2	0.03	<u>0.3</u>	0.43	0.67	30, 70	1022, 1025.15
C1	0.03	0.4	<u>0.33</u>	1	25, 75	1022, 1025.15
C2	0.03	0.4	<u>0.25</u>	1.5	20, 80	1022, 1025.15

Table 1. Summary of runs. Varying parameters are the reduced gravity, g' (m s^{-2}); the topography ratio, φ_T ; and, the two-fluid layer thickness ratio, γ . The theoretical maximum amplitude, A_m , as predicted from Eq. (3.68) in Choi and Camassa (1999) is also indicated.

In the literature, common values for g' where solitary waves have been observed range from
390 0.007 m s^{-2} in the Celtic Sea (Gerkema, 1996) to 0.027 m s^{-2} over the Oregon continental shelf
(Stanton and Ostrovsky, 1998). We approach this broad range of stratification in Sect. 4 to investi-
gate the generation of quasi-linear internal tides. Based on the results of this section, we will argue
by its end why in Sect. 5 we focus on a highly stratified regime ($g' = 0.03 \text{ m s}^{-2}$) for the study of
fully nonlinear waves.

395

For convenience in the discussion of our results, wave properties are scaled as follows. The interfa-
cial displacement, Z , the internal tide amplitude, A , and the soliton amplitude, A_s , are scaled to the
thickness of the upper layer, H_1 . The soliton phase speed, c_s , is scaled to the phase speed of linear
long-wave baroclinic interfacial waves, c_p . Horizontal distances along the x -direction and the soliton
400 width, L_s , are scaled to the wavelength of linear long-wave baroclinic interfacial waves, L_p . Lastly,
we use the scaled Coriolis parameter μ_p , which relates to μ in Sect. 2.1, following $\mu_p = \mu/(2\pi)$.

4 Numerical experiments: Quasi-linear internal tides

Tide-generated solitons emerge from nonlinear disintegration of the underlying internal tides and
405 may be, therefore, naturally subjected to their wave properties. For this reason, we find insightful to
investigate first the wave properties of the underlying internal tides, prior to its nonlinear disintegra-
tion, within the parameter space of this study.

Sandstrom and Quon (1993, 1994) already treated the generation and further disintegration of the
410 internal tide as separate problems. First, the generation was regarded as a hydrostatic process; sec-
ond, the resulting profile was used as a starting point of another, nonhydrostatic model in which
disintegration into solitary waves occurred. Although in nature these processes occur at once and,
hence, the connection between them must be treated carefully, we also believe this is a reasonable
approach to the phenomenon.

415

As described in Sect. 3, we recall that the quasi-linear case includes advective terms from the in-
teractions between the barotropic and baroclinic flows while interactions between baroclinic fields,
the genuinely nonlinear terms, are still absent. Therefore, higher harmonics are naturally generated
when the forcing is increased. The linear case, where advective terms are absent, is added here for
420 assessing potential departures from the quasi-linear case.

Accordingly, Fig. 2 presents an analysis of the internal tide response to the strength of the tidal forc-
ing for runs A1 to C2 (see Table 1). The minimum forcing strength for all cases is $\mathbf{U}_0 = 5 \text{ cm s}^{-1}$.




Fig. 2. Amplitude of the linear, L , and quasi-linear, QL , internal tide scaled to the thickness of the upper layer vs. the Froude number. Varying parameters between panels are: a) the strength of stratification, g' (runs A1, A2 and A3); b) the *topography ratio*, φ_T (runs A1, B1 and B2); and, c) the *two-fluid layer thickness ratio*, γ (runs A1, C1 and C2). The run time is 9 tidal periods. See Table 1 for further details.

In subsequent data-points, the increase of U_0 is of 10 cm s^{-1} from $U_0 = 10 \text{ cm s}^{-1}$ and onwards
 425 up to reaching a $Fr \sim 1.5$.

In the purely linear experiments, the amplitude of the internal tide increases linearly with the in-
 crease of the tidal forcing strength, as derived from theory. Interestingly, the quasi-linear internal
 tide exhibits a limiting amplitude in all runs as the tidal forcing increases well above $Fr = 1$, an
 430 unreported feature up to date. For weak forcing, the amplitude of the quasi-linear internal tides ap-
 proach the linear ones, especially for $Fr \ll 1$ as one would expect; the advective terms then become
 very small. This pattern indicates that the decisive factor on the amplitude saturation of quasi-linear
 internal tides lays on the barotropic advection, which is absent in the linear case.

435 Regarding the comparison between runs with different environmental parameters, we find the fol-
 lowing. In Fig. 2a the increase of stratification causes an earlier deviation between the amplitude
 growth of the quasi-linear and linear cases, hence occurring at a lower Froude number for runs with
 a higher stratification (c. f. runs A1, A2 and A3). The same effect is observed in Fig. 2b when the
 height of the topography is increased. The higher the topography, the earlier that a deviation from
 440 the linear case appears in the Froude number space (c. f. runs A1, B1 and B2). Lastly, no significant
 differences emerge regarding the rise of the quasi-linear departure in Fig. 2c, where the thickness of
 the upper layer varies within the study cases (c. f. runs A1, C1 and C2). These results indicate that
 the wave amplitude saturation, and hence the deviation from the linear case, becomes more impor-
 tant as either the strength of the stratification or the height of the topography increase.

445

Although not shown, it is worth mentioning that the wavelength of the quasi-linear tides does not
 deviate from the linear case in any of the settings of study and is independent of the strength of the
 tidal forcing (and hence of the Froude number) and of the height of the topography. On the contrary,

as predicted from linear theory for interfacial waves, a relative increase of g' or H_1 (with $H_1 < H_2$ and D being constant) generates longer internal tides.

The amplitude saturation described above is further illustrated in Fig. 3 for run A1, where snapshots of leftward-propagating quasi-linear internal tides are shown for various forcing strengths (see legend). This spatial view shows how the increase of the forcing transforms the wave from sinusoidal to an asymmetric shape, indicative of the generation of higher harmonics, but furthermore the amplitude becomes saturated.




Fig. 3. Snapshots of the interfacial displacement of leftward propagating quasi-linear internal tides for run A1 ($H_1 = 30$ m; $L_p = 35.49$ km). The amplitude saturation is evident as the tidal forcing is increased and the flow becomes supercritical (see legend). The run time is 9 tidal periods.

Findings in this section open an interesting question which regards to whether solitons emerging from disintegration of the quasi-linear internal tides may or may not be subjected to saturation amplitudes of the originating waves before they reach a limiting ‘table-top’ shape. For simplicity on answering the above question we focus in the next section on solving the full set of forced-MCC- f equations only for runs A1, B1 and C1, which account for varying the height of the topography and the thickness of the upper layer while preserving high the strength of the stratification. The latter allows us to investigate the broadest range of wave amplitudes, as suggested by Fig. 2a.

5 Numerical experiments: Fully nonlinear internal tides and solitons

The main question to address is whether tidally generated solitons are indeed subjected to the wave properties of the underlying quasi-linear internal tides, as we hypothesized in previous section, and whether they adequate qualitatively and quantitatively to predictions from classical eKdV and MCC theories.

5.1 Tide-generated ‘table-top’ solitons: Run A1

In Fig. 4a a spatial overview of leftward-propagating internal tides and solitons is shown after 9 tidal periods of run time. The tidal forcing is fairly strong and leads to the generation of ‘table-top’ solitons in a supercritical regime ($Fr = 1.13$, $U_0 = 90 \text{ cm s}^{-1}$). In subsequent panels, a set of snapshots zooms in on the space domain of panel (a) to highlight the different stages of the nonlinear disintegration of the internal tides.




Fig. 4. Snapshots of the interfacial displacement of nonlinear internal tides and solitons in run A1 for a supercritical regime ($Fr = 1.13$, $U_0 = 90 \text{ cm s}^{-1}$). (a) Overview of leftward-propagating internal tides and solitons. (b-e) Set of spatial zooms from (a) showing different stages of the nonlinear disintegration of the internal tides. Points Z_a (black dot), Z_b (grey dot), Z_c (black square) and Z_d (grey square) are shown to illustrate how the soliton amplitude, A_s , and width, L_s , are computed in this work (see the text in Sect. 5.1 for details). The run time is 9 tidal periods. For scaling purposes we recall that for run A1: $H_1=30 \text{ m}$ and $L_p = 35.49 \text{ km}$.

At a first stage, panel (b), the internal tide splits up into two different groups of rank-ordered solitons: a train of depressions on the leading edge; and a train of elevations, after the former packet, with initially smaller amplitudes. At a later stage, panel (c), the largest elevations have reached the smaller depressions in the train and three leading solitons at the front present almost equal amplitudes. Previous solitary wave packets, already propagating away from the generation area, are shown in panels (d) and (e) and correspond to preceding disintegrated internal tides. The ‘table-top’ soliton observed at the leading edge of every preceding internal tide emerged in all cases from the first of

the three solitons described previously in panel (c).

As the leading soliton evolves and reaches its maximum amplitude, it also broadens, as predicted by soliton wave theory (Helfrich and Melville, 2006), in comparison with subsequent solitons of smaller amplitude (Fig. 4d,e). The observed increase in the distance between the ‘table-top’ soliton and subsequent (smaller) solitons also indicates that, as expected from theory, the leading soliton moves (phase speed) faster than solitons in the tail.




Fig. 5. Wave evolution of leftward-propagating leading solitons in run A1 under different forcing strengths (see legend). In all panels the x -axis indicates the run time and soliton age (in brackets) in tidal periods. The (dimensionless) wave properties are: (a) soliton amplitude, A_s/H_1 ; (b) soliton width, L_s/D ; (c) instantaneous Froude number, $Fr' = U/c_p$; and, (d) soliton phase speed, c_s/c_p . Note that we take c_p to be negative (leftward propagation) to keep consistency with the physical meaning of the different sign in Fr' . For scaling purposes we recall that in run A1: $H_1 = 30$ m, $D = 100$ m and $c_p = -79$ cm s $^{-1}$.

Because tidally generated solitons propagate through the evolving internal tides, $z = 0$ cannot be used as a reference level to compute the amplitude down to the trough of the soliton (see Fig. 1 and Fig. 4). Similarly, the soliton width cannot be measured taking $z = 0$ as a reference level. A systematic criterion is required to adopt a suitable reference level which allows us to compute the soliton amplitude, A_s , and width, L_s . Here we introduce the reference level Z_a , which for every leftward-propagating soliton locates where the first spatial derivative of the interfacial displacement, Z , approaches zero while being above a certain threshold. This grid-point indicates the location of the front of the leading soliton connecting with the tail of the preceding interfacial tide. Accordingly, the soliton amplitude, A_s , is defined as the vertical distance between Z_a and the trough of the leading soliton, located at Z_b (see, e. g., in Fig. 4c-e). The soliton width, L_s , is defined as the horizontal

distance between Z_c and Z_d , which locate half-way of the vertical distance spanning A_s (see also, 505 e. g., in Fig. 4c-e). Lastly, the soliton phase speed, c_s , is computed by subtracting the velocity of the (mimicked) tidal flow, \mathbf{U} , to the velocity of the soliton embedded within the internal tide.

Using the above criteria Fig. 5 presents the wave evolution of leading solitons under different forcing strengths (see legend) towards a fully developed stage. Contrary to what one might expect, the 510 amplitude of the leading solitons decrease during its evolution (Fig. 5a). This can be ascribed to their tide-generated nature. At an early stage, the disintegration of the internal tide leads at its front to a large depression, and this subsequently evolves to a mature leading soliton propagating through the tail of the preceding internal tide (see Fig. 4c-e).




Fig. 6. Solitary wave solutions for mature leading solitons in run A1 from KdV (grey line), eKdV (black line) and MCC (red line) theories compared to numerical solutions from the forced-MCC equations (colored dots refer to the Froude number and strength of the tidal flow; see legend). (a) Soliton phase speed scaled to the linear long-wave phase speed of baroclinic interfacial waves (c_s/c_p) vs. soliton amplitude scaled to the thickness of the upper layer ($-A_s/H_1$). (b) Soliton width scaled to the total water depth (L_s/D) vs. soliton amplitude scaled to the thickness of the upper layer ($-A_s/H_1$).

515 The soliton reaches its maximum amplitude slightly before the flow becomes critical ($Fr = 0.88$) and attains the ‘table-top’ form in the supercritical regime when forced with a stronger tidal flow ($Fr = 1.13$). Unexpectedly, when the tidal forcing is increased even further, the soliton width starts to decrease while keeping its maximum amplitude (c. f. Fig. 5a and b). Because of the nature of classical eKdV and MCC theory, this feature could not arise before and it does indicate that limiting 520 factors related to the forcing may be acting.

Generally speaking, we distinguish between two types of solitons regarding their time-scales of growth (see Fig. 5a and b). First, the smaller and narrower solitons, generated in a subcritical regime and which present a nearly constant shape quickly after their generation ($Fr \leq 0.5$). Second, the 525 larger and broader solitons, generated in nearly critical and supercritical regimes and which evolve over longer time-scales ($Fr \geq 0.88$). We distinguish here three different states for strongly nonlinear

solitons, which are indicated with vertical dashed lines and labels in Fig. 5a and b. During State I emerging solitons evolve as transient waves which broaden linearly until they reach a fully developed form. Then, solitons propagate as mature waves, State II, which preserve their shape in time and, 530 occasionally, may overtake the preceding internal tide, State III, causing the oscillations observed in the width, amplitude and phase speed curves in Fig. 5a,b,d.

In agreement with the above description, the phase speed curves also reveal a clear distinction between the subcritical and critical/supercritical regimes (Fig. 5d). On the one hand, smaller solitons 535 present a nearly constant phase speed. They were generated with a small or moderate tidal forcing (subcritical flow). On the other hand, larger solitons present an oscillating phase speed which increases over time. They were generated with a relatively strong tidal forcing (critical and supercritical flow). The oscillation amplitude is in this latter case about 5% of its value and is the response to a governing flow where the accelerating and decelerating phases of the soliton are imposed by the 540 favouring and opposing phases of the strong tidal flow. This is visible by comparison of the instantaneous Froude number, Fr' , in Fig. 5c with the soliton phase speed in Fig. 5d. Crucial moments occur when $Fr' = -1$ and $Fr' < -1$. During the former, solitons cannot propagate against the tidal flow and remain stationary. During the latter, leftward-propagating solitons experience a rightward advection driven by the greater tidal flow.

545

Lastly, we compare in Fig. 6 the wave properties of mature forced-MCC solitons³ with KdV-type and MCC soliton solutions (Kakutani and Yamasaki, 1978; Ostrovsky and Stepanyants, 1989; Miyata, 1985, 1988; Choi and Camassa, 1999; Helfrich and Melville, 2006; Gerkema and Zimmerman, 2008). To this aim the soliton width for KdV-type and MCC theories is computed following the 550 same procedure as for the forced-MCC solitons, i. e. we use points Z_c and Z_d (see Fig. 4c-e).

As expected, small tide-generated solitons approach the linear long-wave phase speed for baroclinic interfacial waves ($c_s/c_p \approx 1$), while larger tide-generated solitons increase their phase speed following a curve as eKdV and MCC solutions do. However, because tide-generated solitons *ride* on 555 internal tides their wave properties are not simply the response to a settled two-layer fluid system, as it occurs for eKdV and MCC solitons, but they are also subjected to the forcing of the system and to a variable background flow (the internal tide). We suggest the above scenario might account for the slower phase speeds of the forced-MCC solitons when compared to their eKdV and MCC counterparts. Interestingly, this difference slightly decreases as the solitons grow (c. f. the length of 560 the colored dashed lines in Fig. 6a).

As regards to the relationship between the soliton width and amplitude, tide-generated solitons fol-

³These wave properties correspond to solitons of State II (mature solitons) after time averaging over a tidal cycle.

low a parallel behaviour to that predicted by eKdV and MCC soliton solutions, broadening as they approach their maximum amplitude. By this broadening, strongly nonlinear solitons develop the ‘table-top’ shape, although forced-MCC equations generate some larger and narrower solitons than their eKdV and MCC counterparts (Fig. 6b).

5.2 Growth limitation of tide-generated solitons: Runs B1 and C1




Fig. 7. Wave evolution of leftward-propagating leading solitons in run B1 under different forcing strengths (see legend). In all panels the x -axis indicates the run time and soliton age (in brackets) in tidal periods. The (dimensionless) wave properties are: (a) soliton amplitude, A_s/H_1 ; (b) soliton width, L_s/D ; (c) instantaneous Froude number, $Fr' = U/c_p$; and, (d) soliton phase speed, c_s/c_p . Note that we take c_p to be negative (leftward propagation) to keep consistency with the physical meaning of the different sign in Fr' . For scaling purposes we recall that in run B1: $H_1=30$ m, $D = 100$ m and $c_p = -79$ cm s $^{-1}$.

We use for runs B1 and C1 a similar range of Froude number as for run A1, however they present a more weakly nonlinear regime where a striking feature emerges. Leading solitons exhibit a maximum amplitude which is not related to a ‘table-top’ form and which cannot be exceeded by further increasing the tidal forcing (see Figs. 7a and 8a). They reach this limiting amplitude in both cases when the flow is supercritical (run B1: $Fr = 1.26$; and, run C1: $Fr = 1.33$). More importantly, above this limit, the strengthening of the tidal forcing leads to a narrowing and amplitude decrease of the leading solitons (Figs. 7a,b and 8a,b). We recall here that the decrease of the soliton width after reaching its maximum is also observed when the tidal forcing leading to limiting solitons in run A1 is increased (see Figs. 5a,b).




Fig. 8. Same as Figure 7 but for run C1. For scaling purposes we recall that in run C1: $H_1=25$ m, $D = 100$ m and $c_p = -75$ cm s $^{-1}$.

The above results support that tidally generated solitons may be conditioned in the real ocean to a limited growth which is beyond the classical view of KdV and MCC-type of models and which
 580 lays on the saturation of the underlying quasi-linear internal tide as the tidal forcing increases (see Sect. 4).

According to their phase speed, and in agreement with findings from run A1, two types of leading solitons also emerge in runs B1 and C1. The larger nonlinear solitons (critical and supercritical
 585 regime), which exhibit an oscillating speed in phase with the tidal flow and which increases over time. And, the smaller nonlinear solitons (subcritical regime), which exhibit a nearly constant phase speed (Figs. 7a,c,d and 8a,c,d).

From Figs. 9 and 10, we gain further insights on the different stages by which internal tides gener-
 590 erate saturated leading solitons in runs B1 ($Fr = 1.26$, $U_0 = 100$ cm s $^{-1}$) and C1 ($Fr = 1.33$, $U_0 = 100$ cm s $^{-1}$). By contrast to run A1 (Fig. 4), here the internal tides do not split up into two different groups of solitons but disintegrate into solitary wave packets of rank-ordered depressions. Also, the ‘table-top’ solitary waves that lead the long-life internal tides in run A1 (Fig. 4d,e) are not present in runs B1 and C1, as previously discussed from the wave property analyses. We




Fig. 9. Snapshots of the interfacial displacement of nonlinear internal tides and solitons in run B1 for a supercritical regime ($Fr = 1.26$, $U_0 = 100 \text{ cm s}^{-1}$). (a) Overview of leftward-propagating internal tides and solitons. (b-e) Set of spatial zooms from (a) showing different stages of the nonlinear disintegration of the internal tides. The run time is 9 tidal periods. For scaling purposes we recall that for run B1: $H_1 = 30 \text{ m}$ and $L_p = 35.49 \text{ km}$.

595 attribute this absence to the lower height of the topography in run B1 and the decrease of the upper layer thickness in run C1.

On the one hand, the smaller topography generates quasi-linear internal tides which are smaller than those in run A1 (see Fig. 2). With all other environmental parameters being the same, the smaller
600 internal tide in run B1 prescribes then a more weakly nonlinear disintegration. On the other hand, the thinner H_1 in run C1 requires a maximum amplitude to attain the ‘table-top’ form which is larger than for runs A1 and B1 (see A_m/H_1 in Table 1). In this context, the smaller quasi-linear internal tides generated in run C1, by comparison with run A1 (see Fig. 2c), do not lead to strongly nonlinear disintegration when the full forced-MCC equations are solved; not even in the supercritical regime
605 of $Fr > 1$. Indeed, although both run A1 and run C1 generate leading solitons of early stage with a relatively similar amplitude, the latter run exhibits mature leading solitons which are significantly smaller and narrower (c. f., Figs. 4d,e and 10d,e), suggesting that dispersive effects might overcome nonlinearities more noticeably when the upper layer is thinner.

610 When compared with solitary wave solutions from eKdV and MCC theories, the growth-limiting ef-




Fig. 10. Snapshots of the interfacial displacement of nonlinear internal tides and solitons in run C1 for a supercritical regime ($Fr = 1.33$, $U_0 = 100 \text{ cm s}^{-1}$). (a) Overview of leftward-propagating internal tides and solitons. (b-e) Set of spatial zooms from (a) showing different stages of the nonlinear disintegration of the internal tides. The run time is 9 tidal periods. For scaling purposes we recall that for run C1: $H_1 = 25 \text{ m}$ and $L_p = 33.54 \text{ km}$.

fect of the tidal forcing becomes a remarkable feature of forced-MCC solitons generated in runs B1 and C1 since they reach a limiting amplitude but do not attain a ‘table-top’ form (Fig. 11b,d). In this context it is also worth while noting that in run B1 saturated solitons present amplitudes larger than those predicted by eKdV and MCC theories whereas in run C1 saturated solitons present amplitudes well below those predicted by eKdV and MCC theories. Counterintuitively, it is also evident from both runs B1 and C1 that largest solitons decrease their amplitude and width as the tidal forcing increases above that which generates the saturated solitons, as previously noted from Figs. 7 and 8. Regarding the relationship between the soliton phase speed and amplitude, both runs B1 and C1 follow a similar curve to that predicted by eKdV and MCC theories (Fig. 11a,c), although the phase speed of forced-MCC solutions is slower in all cases, as it occurred for run A1 (see Fig. 6a). Also similar to run A1, the deviation in phase speed between MCC and forced-MCC solutions is observed to decrease as the solitons grow (c. f. the length of the colored dashed lines in Fig. 11a,c), suggesting that small solitons might be more subjected to effects related to the forcing system.

625




Fig. 11. Solitary wave solutions for mature leading solitons in run B1 (top row) and run C1 (bottom row) from KdV (grey line), eKdV (black line) and MCC (red line) theories compared to numerical solutions from the forced-MCC equations (colored dots refer to the Froude number and strength of the tidal flow; see legend). (a, c) Soliton phase speed scaled to the linear long-wave phase speed of baroclinic interfacial waves (c_s/c_p) vs. soliton amplitude scaled to the thickness of the upper layer ($-A_s/H_1$). (b, d) Soliton width scaled to the total water depth (L_s/D) vs. soliton amplitude scaled to the thickness of the upper layer ($-A_s/H_1$).

5.3 Effects of the Earth's rotation: Runs A1, B1 and C1

In Fig. 12 the effects of the Earth's rotation on the wave evolution of fully nonlinear tide-generated solitons are shown for runs A1, B1 and C1. The different colored lines refer to: rotationless case (black line); $\theta = 15^\circ$, $\mu_p = 0.27$ (green line); $\theta = 30^\circ$, $\mu_p = 0.52$ (blue line); and, $\theta = 45^\circ$, $\mu_p = 0.73$ (red line). 630

In agreement with previous studies we observe in all panels that an increase of the latitude leads to larger dispersive effects due to Coriolis dispersion, which prevents the nonlinear internal tide from disintegrating into strongly nonlinear solitons (Gerkema and Zimmerman, 1995; Gerkema, 1996). 635

This causes the long internal waves to envelope less solitary waves. Also, the internal tides are shown to travel faster as rotation becomes stronger due to rotation increases the phase speed of the linear internal tide, $c_f^2 = c_0^2 + f^2/k^2$, with k being the wavelength of the internal tide). Although the soliton speeds themselves are only very weakly affected by rotation, they appear traveling faster since they are embedded in the internal tide from which they emerge. As a consequence, leading 640 solitons overtake more quickly preceding internal tides.




Fig. 12. Effects of the Earth's rotation through a set of snapshots from runs A1 ($Fr = 1.13$; $U_0 = 90 \text{ cm s}^{-1}$), B1 ($Fr = 1.26$; $U_0 = 100 \text{ cm s}^{-1}$) and C1 ($Fr = 1.33$; $U_0 = 100 \text{ cm s}^{-1}$). The run time is 9 tidal periods. (a,c,e) Overview of leftward-propagating tide-generated internal tides and solitons. (b,d,f) Spatial zoom from the corresponding overview. In all panels the rotationless case ($\theta = 0$, $\mu_p = 0$) is shown as a black line. Rotating cases are: $\theta = 15^\circ$, $\mu_p = 0.27$ (green line); $\theta = 30^\circ$, $\mu_p = 0.52$ (blue line); and, $\theta = 45^\circ$, $\mu_p = 0.73$ (red line).

6 Discussion and conclusions

We investigate limiting amplitudes of internal tides and solitons using a generalization of the fully nonlinear MCC equations (Miyata, 1985, 1988; Choi and Camassa, 1999), extended here with forcing terms and Coriolis effects (forced-MCC- f). The focus is on the effects of adding a forcing, which represents a novelty in the existing literature and provides a closer view to an ocean-like scenario. The mechanism for internal tide generation is represented by a horizontally oscillating sill, mimicking a barotropic tidal flow over topography. Solitons are generated by a disintegration of the internal tide.

650

The application of an oscillating topography is not completely equivalent to the oceanic case of a tidal flow over a topography at rest. For this reason we have restricted our analyses to a parameter space where a semi-equivalence between both forcing systems was demonstrated (*Appendix C*). This agreement encourages us to conclude that our findings are not an artifact caused by the use of a mimicked barotropic tidal flow. Of course the findings presented here cannot describe the whole variety of the specific oceanic conditions. However, we believe that this study improves our understanding on the generation and evolution of tide-generated solitons.

Numerical solutions show that strongly nonlinear tide-generated solitons attain in some cases a limiting table-shaped form, in agreement with classical soliton theory. However, results also reveal that tide-generated solitons may alternatively be limited by saturation of the underlying quasi-linear internal tide. In the purely linear system the amplitude of the internal tide increases linearly with the strength of the barotropic tidal flow but as the forcing becomes stronger advective terms become stronger too and cannot be neglected. This is accounted for in the quasi-linear system, where products of barotropic and baroclinic fields in the advective terms are retained while still ignoring interactions of the baroclinic field with itself. As a result, a saturation in the amplitude of the internal tide occurs, generally, when the tidal forcing becomes supercritical; then, a further increase of the tidal flow does not produce a larger internal tide. This effect seems to have passed unnoticed in previous studies, but turns out to be a key factor in the subsequent disintegration of the internal tide into solitons. It implies that when one includes the genuinely nonlinear effects, i. e. products of baroclinic terms, resulting solitons may stay well below their formal limiting amplitude, no matter how strong the forcing. Interestingly, an increase of the tidal forcing above that which generates table-shaped solitons, or above that which *simply* generates solitons attaining a limited-growth, causes first its progressive narrowing and, subsequently, an amplitude decrease. The upshot is that increasing the tidal forcing above a certain strength does not lead to larger solitons but, counterintuitively, to smaller ones.

Motivated by the above finding we performed analogous runs using the full set of weakly nonlin-

ear equations derived in Gerkema (1996). Because these equations are built around the framework
680 of the classical KdV theory and Klein-Gordon equations, one should not expect that the amplitude
saturation of solitons could occur. Nevertheless, results (not shown) demonstrate that both the quasi-
linear internal tides and weakly nonlinear tide-generated solitons also exhibit a limiting amplitude.
Noting that this model works with an actual tidal flow over a topography at rest, it seems reasonable
to argue that the limiting factor is then related to the addition of a tidal forcing. This gives support
685 to conclude that findings from the forced-MCC- f equations represent an insightful extension to the
fully nonlinear frame of work where tide-generated solitons may attain limiting amplitudes with or
without reaching a ‘table-top’ form, then subjected to a saturation amplitude of the underlying internal
tide prior to its disintegration into solitary waves.

690 Another departure from classical theories is that strongly nonlinear tide-generated solitons may exhibit
larger maximum amplitudes than predicted from eKdV and MCC solutions, while soliton phase
speeds are always smaller. We attribute these differences to the fact that tide-generated solitons *ride*
on internal tides and, hence, their wave properties are not simply the response to a settled two-layer
fluid system, as it occurs for eKdV and MCC solitons, but are also subjected to the forcing of the sys-
695 tem and to a variable background flow and internal displacement imposed by the internal tide itself.
In this context, numerical results also show that solitons propagate *freely* from the source only when
the tidal flow is small (subcritical flow), while an increase of the tidal forcing generates accelerating
and decelerating phases of the soliton phase speed imposed by the favouring and opposing phases of
the strong tidal flow (critical and supercritical flow). Lastly, it is also worth while mentioning that in
700 all cases of study the time-scale of growth for large solitons (critical and supercritical flow) appears
to be longer than for small solitons (subcritical flow), which abandon its transient form sooner.

In relation to the rotational cases, and in agreement with previous studies (Gerkema and Zimmer-
man, 1995; Gerkema, 1996), numerical results from the forced-MCC- f equations show that when
705 rotation becomes stronger, the dispersive effect of the Coriolis force becomes stronger too and over-
comes nonlinearities, thus preventing the internal tide from disintegration into strongly nonlinear
solitons.

Before concluding we must note, as Ostrovsky and Grue (2003) previously did, that fully nonlinear,
710 weakly nonhydrostatic models entail a paradox to the effect that strongly nonlinear solitons appear
from a set of equations that have strong nonlinearity but weak dispersion, while the very existence
of solitons presume a balance between the two. In our case, the MCC-type model is used, involv-
ing only the lowest-order nonhydrostatic dispersive terms. Despite the small parameter featuring in
the nonhydrostatic terms, they may actually become large in practice (i.e., in the numerical runs)
715 if internal wave profiles are steepening, hence contradicting the original assumption. Indeed, there

is no guarantee that the higher-order dispersive terms, which were dropped from these equations, would always remain small. A suggestion for future work is, therefore, to check our results against a numerical computation with a fully nonlinear nonhydrostatic set of equations.

720 *Acknowledgements.* Financial support was provided by the Spanish government (Ministerio de Ciencia e Innovación) through a Ph.D. grant (FPU) awarded to the first author (AP2007-02307), and through the Netherlands Organization for Scientific Research (NWO), section Earth and Life Sciences (ALW), via the ZKO grant no. 839.08.431 awarded to the ‘Indian-Atlantic EXchange in present and past climate’ (INATEX) program. The authors also wish to thank three anonymous reviewers for their thoughtful comments to improve this paper.

725 **Appendix A**

Numerical strategy

We define a grid in time and space for discretization of the various derivatives of the system. Then,

$$t_n = n\Delta t \quad \text{and} \quad x_j = j\Delta x$$

are introduced for integer values of n (time-step) and j (spatial-step), where Δt and Δx are the magnitude of the steps. Time and spatial dependent variables are described as, e.g. $y(t_n, x_j)$, at any time and position. Thus, y_j^n means the value of the variable y at the current time and spatial-step, n and j , respectively. And, consequently, $n + 1$ represents the ‘next time-step’, and so $n - 1$ the ‘previous time-step’, what applies analogously for j in the space grid.

735 The various derivatives in the model are discretized with centered difference approximations (Durran, 1999) as follows

$$y_t(t_n, x_j) \cong \frac{y_j^{n+1} - y_j^n}{\Delta t}, \quad (\text{A1})$$

$$740 \quad y_x(t_n, x_j) \cong \frac{y_{j+1}^n - y_j^n}{\Delta x}, \quad (\text{A2})$$

$$y_{xx}(t_n, x_j) \cong \frac{y_{j+1}^n - 2y_j^n + y_{j-1}^n}{(\Delta x)^2}, \quad (\text{A3})$$

$$y_{xt}(t_n, x_j) \cong \frac{y_{j+1}^{n+1} - y_{j+1}^n - (y_{j-1}^{n+1} - y_{j-1}^n)}{2\Delta x\Delta t}, \quad (\text{A4})$$

745

$$y_{xxt}(t_n, x_j) \cong \frac{y_{j+1}^{n+1} - y_{j+1}^n - 2(y_j^{n+1} - y_j^n) + (y_{j-1}^{n+1} - y_{j-1}^n)}{(\Delta x)^2\Delta t}. \quad (\text{A5})$$

(A6)

Initiailly the system is at rest with mean horizontal velocities, \bar{u}_i and \bar{v}_i , and displacement of the interface, ζ , being all zero at the first two time levels ($n - 1, n$), what represent the initialization
750 fields. The thickness of the upper, h_1 , and lower layer, h_2 , together with the topography, $h(X)$, draw the scenario where the two-layer system runs. At the next time-step ($n + 1$), we start to move the topography to the right creating the effect of a tidal motion flowing to the left. For given U , i. e. scaled velocity of moving topography (Eq. (47)), and time-step, the excursion of the topography is a known quantity which is used to shift (first, second and third) spatial derivatives of $h(X)$ at every
755 new time-step.

The time derivatives of the \bar{v}_i -momentum and continuity equations (51), (52) and (53) are solved numerically using the third-order Adams-Bashforth approximation (Durrant, 1999), for which \bar{v}_1, \bar{v}_2 and ζ at the next time-step ($n + 1$), and at all j positions, are determined in terms of the known
760 quantities at the previous two time-steps ($n - 1, n$).

However, solving numerically \bar{u}_1 from Eq. (49) is not straightforward as we deal with three different time derivatives of \bar{u}_1 accompanied with space-time-dependent coefficients. Thus, after collecting the various time derivatives involving \bar{u}_1 on one side and remaining terms on the other side, the
765 horizontal momentum equation of \bar{u}_1 evolves to a generic expression in the form of

$$a \bar{u}_{1,t} + b \bar{u}_{1,xt} + c \bar{u}_{1,xxt} = Y(t_n, x_j) \quad (\text{A7})$$

where a, b and c collect spatial derivatives of space-time dependent variables ($\zeta(x, t)$ and $h(x, t)$); and, $Y(t_n, x_j)$ represents a collection of known quantities whose values may be dependent on time and/or space. In the remainder, we describe the numerical strategy we follow to solve this
770 problem.space-time-dependent partial differential equations. If now we operate the time derivative as a common factor in the left-hand side, the result leads to

$$(a \bar{u}_1 + b \bar{u}_{1,x} + c \bar{u}_{1,xx})_t = Y(t_n, x_j) + (a_t \bar{u}_1 + b_t \bar{u}_{1,x} + c_t \bar{u}_{1,xx}) \quad (\text{A8})$$

what helps us to introduce a new variable, \bar{U}_1 , which groups coefficients a, b, c and time derivatives of \bar{u}_1 and turns our problem into a numerically solvable expression in the form of

$$775 \quad \bar{U}_{1,t} = Y(t_n, x_j) + (a_t \bar{u}_1 + b_t \bar{u}_{1,x} + c_t \bar{u}_{1,xx}) \quad (\text{A9})$$

It is important to recall here that $Y(t_n, x_j)$ and the spatial derivatives of \bar{u}_1 are both evaluated at the current time-step (n); and, the time derivatives of a, b and c , which involve values of ζ at the current (n) and new time-step ($n + 1$), have been previously achived with Eq. (53) via Adams-Bashforth approximation. This allows to rewrite the above expression as

$$780 \quad \bar{U}_{1,t} = R(t_n, x_j) \quad (\text{A10})$$

by grouping all known quantities on the right-hand side under the variable $R(t_n, x_j)$. Next we need to discretize the time derivative of \overline{U}_1 but before doing that, we work out and discretize its spatial derivatives using Eqs. (A2) and (A3), what results in

$$\overline{U}_1 = \left(a_j - \frac{2c_j}{2\Delta x}\right) \overline{u}_{1j} + \left(\frac{-b_j}{2\Delta x} + \frac{c_j}{(\Delta x)^2}\right) \overline{u}_{1j-1} + \left(\frac{b_j}{2\Delta x} - \frac{c_j}{(\Delta x)^2}\right) \overline{u}_{1j+1}$$

785 which we rewrite by introducing factors d , e and f as follows

$$\overline{U}_{1j} = d_j \overline{u}_{1j} + e_j \overline{u}_{1j-1} + f_j \overline{u}_{1j+1}. \quad (\text{A11})$$

If we now discretize the time derivative of \overline{U}_1 and apply Adams-Bashforth, we obtain a numerically solvable expression for \overline{U}_1 at the next time step, which reads

$$\overline{U}_{1j}^{n+1} = \overline{U}_{1j}^n + \frac{\Delta t}{12} \left(23R_j^n - 16R_j^{n-1} + 5R_j^{n-2}\right), \quad (\text{A12})$$

790 where \overline{U}_{1j}^{n+1} actually includes

$$\overline{U}_{1j}^{n+1} = d_j^{n+1} \overline{u}_{1j}^{n+1} + e_j^{n+1} \overline{u}_{1j-1}^{n+1} + f_j^{n+1} \overline{u}_{1j+1}^{n+1}. \quad (\text{A13})$$

To close our system we still need to obtain \overline{u}_{1j}^{n+1} for all j terms. To that end, the equation above is more complicated to solve and gives rise to implicit equations, as we have not only the unknown \overline{u}_{1j}^{n+1} , but also $\overline{u}_{1j-1}^{n+1}$ and $\overline{u}_{1j+1}^{n+1}$, which come from the mixed second and third derivatives of u_1

795 in Eq. (A7). However, this is a well-known problem that can be solved using the tridiagonal matrix algorithm (TDMA), also known as the Thomas algorithm (Logan, 1987).

Following the numerical strategy described above, the model resolution is closed for every new time level $n + 1$ and the model equations can be solved successfully.

800

The choice of the space-time steps Δt and Δx is based on two main requirements. Firstly, the resolution in x (Δx) must be sufficiently fine to resolve third-derivative terms and ensure that any short, solitary-like waves are properly resolved. Nevertheless, dealing with equivalent equations to Miyata (1988) and Choi and Camassa (1999), as we do in our model, Kelvin-Helmholtz instabilities are not filtered out. In this regard, Jo and Choi (2002) found that solitary waves of sufficient amplitude could be unstable at high wave numbers to Kelvin-Helmholtz instability. Thus, if the grid resolution is too fine, unstable short waves will emerge near the wave crest and ultimately overwhelm the calculations and explode numerically (Jo and Choi, 2002; Helfrich and Melville, 2006; Helfrich and Grimshaw, 2008). In some cases, the instability can be controlled by filtering out wavenumbers above a threshold (W. Choi 2007, personal communication cited in Helfrich and Grimshaw (2008)).

810

For our numerical experiments we consider a Δx course enough to prevent the problem. A second condition follows from the requirement of stability. Then, for a given spatial step one may take the Courant-Friedrichs-Lewy condition for the linearized equations as an indication of the required time step. The criterion implies that $\Delta x/\Delta t$ should be larger than the phase speed of the wave; taking

815 special care where the advection by the barotropic tidal flow (here mimicked with the moving topography) should be added to the phase speed to apply the criterion properly (Gerkema, 1994).

For the simulations we present, it was not needed to filter out wavenumbers above a threshold to control Kelvin-Helmholtz instabilities as we designed the space-time grid to avoid this problem
 820 following previous conditions. However, in some cases, specially in the simulations where the forcing was fairly strong, an additional trick was needed to retain stability around the generation area (Gerkema, 1994). In those cases averages were taken in the vicinity of the top of the ridge (around the steepest part of the topography), where the instabilities arised. At one particular point (x_j, t_n) in space-time, new values of \bar{u}_i , \bar{v}_i and ζ were calculated by taking the average of the old values at
 825 x_{j-1} , x_j and x_{j+1} , and subsequently in time between t_n and t_{n-1} . The disturbance provoked by this procedure was tested and found to be a minor effect only, as it was only applied over the closest region to the top of the topography.

Appendix B

830 B1 forced-MCC- f model equations

In Appendix A, the numerical scheme used to solve the model is explained using a generic expression (A7) for the \bar{u}_i horizontal momentum equation (49). Here we present the full set of nondimensional forced-equations actually used for the numerical solving of the model. The procedure to that end is as follows.

835

Firstly, all terms of the \bar{u}_i horizontal momentum equation (49) are worked out and grouped according to their physical effects (i. e. *linear*, *nonlinear* and *dispersive effects* from the upper and lower layer, and from topography), leaving unknown quantities involving time derivatives of \bar{u}_1 on the left-hand side. The resulting expression (31) resembles (A7), where coefficients a , b and c involve derivatives
 840 of space-time dependent variables and $Y(t_n, x_j)$ is represented here by the sum of all terms on the right-hand side,

$$\begin{aligned}
 a \bar{u}_{1,t} + b \bar{u}_{1,xt} + c \bar{u}_{1,txt} = & \text{linear} + \text{nonlinear} + \text{dispersive}_1 + \text{dispersive}_2 + \text{dispersive}_{topo} \\
 & + \delta_2 \left[(\eta_2 h_x - \eta_2 \zeta_x) \phi_x - \frac{\eta_2^2}{3} \phi_{xx} + \phi \left(\frac{\eta_2}{2} h_{xx} + \zeta_x h_x \right) \right],
 \end{aligned}
 \tag{B1}$$

845

$$\bar{u}_2 = \frac{Uh - \eta_1 \bar{u}_1}{\eta_2}, \tag{50}$$

$$850 \quad \bar{v}_{1,t} = -\mu\bar{u}_1 - \bar{u}_1\bar{v}_{1,x} + O(\delta^2), \quad (51)$$

$$\bar{v}_{2,t} = -\mu\bar{u}_2 - \bar{u}_2\bar{v}_{2,x} + O(\delta^2), \quad (52)$$

$$855 \quad \zeta_t = (h_1 - \zeta)\bar{u}_{1,x} - \bar{u}_1\zeta_x. \quad (53)$$

with

$$\phi = \frac{1}{\eta_2} \left[hU_t + U^2h_x + (\bar{u}_1 - \bar{u}_2)(\eta_1\bar{u}_{1,x} - \bar{u}_1\zeta_x) + \bar{u}_2Uh_x \right] \quad (B2)$$

$$860 \quad a(\zeta, h) = 1 + \frac{\delta\eta_2}{1-h} \left[(\eta_2h_x - \eta_2\zeta_x)(\eta_1/\eta_2)_x - \frac{\eta_2^2}{3}(\eta_1/\eta_2)_{xx} + \frac{\eta_1}{\eta_2} \left(\frac{\eta_2}{2}h_{xx} + \zeta_xh_x \right) \right], \quad (B3)$$

$$b(\zeta, h) = \delta \left(1 - \frac{\eta_1}{1-h} \right) \eta_1\zeta_x + \frac{\delta\eta_2}{1-h} \left[\frac{\eta_1}{\eta_2}(\eta_2h_x - \eta_2\zeta_x) - \frac{2\eta_2^2}{3}(\eta_1/\eta_2)_x \right], \quad (B4)$$

865

$$c(\zeta, h) = -\delta \left(1 - \frac{\eta_1}{1-h} \right) \frac{\eta_1^2}{3} - \frac{\delta\eta_2}{(1-h)} \frac{\eta_1\eta_2}{3}, \quad (B5)$$

$$870 \quad \text{linear} = \mu\bar{v}_1 + \zeta_x + \frac{1}{1-h} \left[hU_t + U^2h_x + \bar{u}_2h_t \right], \quad (B6)$$

$$\text{nonlinear} = -\bar{u}_1\bar{u}_{1,x} + \frac{1}{1-h} \left[(\bar{u}_1 - \bar{u}_2)\zeta_t + \bar{u}_1\eta_1\bar{u}_{1,x} + \bar{u}_2\eta_2\bar{u}_{2,x} - \mu(\eta_1\bar{v}_1 + \eta_2\bar{v}_2) - \eta_1\zeta_x \right], \quad (B7)$$

$$875 \quad \text{dispersive}_1 = \delta \left(1 - \frac{\eta_1}{1-h} \right) \left[-\eta_1\zeta_x(\bar{u}_1\bar{u}_{1,xx} - (\bar{u}_{1,x})^2) + \frac{\eta_1^2}{3}(\bar{u}_1\bar{u}_{1,xxx} - \bar{u}_{1,x}\bar{u}_{1,xx}) \right], \quad (B8)$$

$$\text{dispersive}_2 = \frac{\delta\eta_2}{1-h} \left[-\eta_2\zeta_x(\bar{u}_2\bar{u}_{2,xx} - (\bar{u}_{2,x})^2) - \frac{\eta_2^2}{3}(\bar{u}_2\bar{u}_{2,xxx} - \bar{u}_{2,x}\bar{u}_{2,xx}) \right], \quad (B9)$$

880

$$\begin{aligned} \text{dispersive}_{topo} &= \frac{\delta\eta_2}{(1-h)} \left[\bar{u}_2h_x(\eta_2\bar{u}_{2,xx} + \zeta_x\bar{u}_{2,x}) \right. \\ &+ \frac{\eta_2}{2}(U_t h_{xx} + U^2 h_{xxx} + 2U\bar{u}_{2,x}h_{xx} + 2\bar{u}_2Uh_{xxx} + 3\bar{u}_2\bar{u}_{2,x}h_{xx} + \bar{u}_2^2h_{xxx}) \\ &\left. + \zeta_x(U_t h_x + U^2 h_{xx} + 2\bar{u}_2Uh_{xx} + \bar{u}_2^2h_{xx}) \right]. \end{aligned} \quad (B10)$$

885 **B2 Linear and quasi-linear forced-MCC- f model equations**

The quasi-linear forced-MCC- f model follows from neglecting the purely nonlinear terms and weakly nonhydrostatic dispersive terms in (B1) and (50)–(53). The equations are linear with regard to the baroclinic fields, but the coefficients become time-dependent due to barotropic factors (which are prescribed) and, therefore, higher harmonics will be generated when the forcing is increased. Then, the quasi-linear forced-MCC- f equations
 890 read,

$$\bar{u}_{1,t} = \mu \bar{v}_1 + \zeta_x + \frac{1}{1-h} \left[hU_t + U^2 h_x + \bar{u}_2 h_t - \mu(h_1 \bar{v}_1 + ((h_2 - h)\bar{v}_2) - h_1 \zeta_x \right] \quad (\text{B11})$$

$$895 \quad \bar{u}_2 = \frac{Uh - h_1 \bar{u}_1}{h_2 - h}, \quad (\text{B12})$$

$$\bar{v}_{1,t} = -\mu \bar{u}_1, \quad (\text{B13})$$

$$900 \quad \bar{v}_{2,t} = -\mu \bar{u}_2, \quad (\text{B14})$$

$$\zeta_t = h_1 \bar{u}_{1,x}. \quad (\text{B15})$$

905

We notice that the linear runs were actually done somewhat indirectly by taking the quasi-linear forced-MCC- f equations above, (B11)–(B15), and reducing the forcing by a factor of 100 since the quasi-linear terms cannot be removed explicitly in this model setting without also removing the forcing. Afterwards we enhance the amplitude in the plots accordingly. By reducing the forcing, we effectively enter the linear regime.

910

Appendix C

Oscillating topography vs. tidal flow

A Galilean transformation involves two frames of reference which move with constant and rectilinear speed with respect to each another. Hence, observations made in one frame can be converted to another, as physical
 915 laws are identical. However, our oscillating topography is not an inertial frame since it is accelerated with respect to a situation where the topography is at rest (as in the ocean). It is, therefore, not evident that the results from the two frames are equivalent.

We use the generation model of weakly nonlinear, weakly nonhydrostatic interfacial waves derived in Gerkema
 920 (1996) (*G1996*), which works with tidal motion over a fixed topography, as a benchmark for testing the impact of our ‘non-inertial’ frame of reference. If we compare interfacial waves generated from the nonlinear version of both models, differences are expected to arise from the fact that forced-MCC equations are fully nonlinear. For this reason we restrict the comparison to the linear and quasi-nonlinear model versions. If the results between the models turn out to be similar, it thus seems reasonable to accept that within the parameter space of

925 study we can compare our present setting to that in the ocean setting.

In Fig. C.1, interfacial waves generated from both models are presented for various numerical experiments under a fairly strong forcing, i. e. when both models may be expected to deviate more noticeably from each other. Our interest focus then on the upper limit of the supercritical regime ($Fr > 1$) that we can reach while preserving a good agreement between both generation mechanisms. The different settings in Fig. C.1 differ
 930 in the strength of stratification from top to bottom panels, while the thickness of the upper and lower layer ($H_1 = 30$ m, $H_2 = 70$ m) and the height and width of the sill are kept fixed ($H_T = 40$ m and $H_L = 10$ km in Eq. 54).

935 Results from Fig. C.1 indicate that in all cases a close correspondence exists between numerical solutions from *GI996* (gray line) and the forced-MCC equations (black line), suggesting only a minor impact of the non-inertial nature of our frame of reference when reaching up to a $Fr \sim 1.5$. These results encourage us to approach in our study the strength (velocity amplitude) of the oscillating topography as the ‘strength of the tidal flow’ within the parameter space of study.

940




Fig. C.1. Linear (left panels) and quasi-linear (right panels) interfacial waves generated via a tidal flow over a sill from the model equations derived in (Gerkema, 1996) (grey line) and via a horizontally oscillating sill from the model equations derived in this study (black line). The Froude number and corresponding strength of the (mimicked) tidal flow are indicated in the upper-right corner of each panel. For scaling purposes one must note that the wavelength of the linear long-wave interfacial wave, L_p , varies from top to bottom panels as: $L_p = 35.5$ km ($g' = 0.03$ m s $^{-1}$) in (a, b); $L_p = 29$ km ($g' = 0.02$ m s $^{-1}$) in (c, d); and, $L_p = 20.5$ km ($g' = 0.01$ m s $^{-1}$) in (e, f). The run time is 9 tidal periods.

References

- Apel, J. R., Ostrovsky, L. A., Stepanyants, Y. A., and Lynch, J. F.: Internal solitons in the ocean, Tech. rep., Woods Hole Oceanographic Institution, 2006.
- Choi, W. and Camassa, R.: Fully nonlinear internal waves in a two-fluid system, *J. Fluid Mech.*, 396, 1–36,
945 1999.
- Da Silva, J. C. B., Buijsman, M. C., and Magalhaes, J. M.: Internal waves on the upstream side of a large sill of the Mascarene Ridge: a comprehensive view of their generation mechanisms and evolution, *Deep-Sea Res. I*, 99, 87–104, 2015.
- Durrant, D.: Numerical methods for wave equations in geophysical fluid dynamics, vol. 32, Springer Verlag,
950 1999.
- Gerkema, T.: Nonlinear dispersive internal tides: generation models for a rotating ocean, Ph.D. thesis, Utrecht University, 1994.
- Gerkema, T.: A unified model for the generation and fission of internal tides in a rotating ocean, *J. Mar. Res.*, 54, 421–450, 1996.
- 955 Gerkema, T. and Zimmerman, J. T. F.: Generation of nonlinear internal tides and solitary waves, *J. Phys. Oceanogr.*, 25, 1081–1094, 1995.
- Gerkema, T. and Zimmerman, J. T. F.: An introduction to internal waves: Lecture Notes, R. Neth. Inst. for Sea Res., Den Burg., 2008.
- Helfrich, K. and Grimshaw, R.: Nonlinear disintegration of the internal tide, *J. Phys. Oceanogr.*, 38, 686–701,
960 2008.
- Helfrich, K. R.: Decay and return of internal solitary waves with rotation, *Phys. Fluids*, 19, 026 601–1,026 601–9, 2007.
- Helfrich, K. R. and Melville, W. K.: Long nonlinear internal waves, *Annu. Rev. Fluid Mech.*, 38, 395–425, 2006.
- 965 Jo, T.-C. and Choi, W.: Dynamics of strongly nonlinear internal solitary waves in shallow water, *Stud. Appl. Math.*, 109, 205–227, 2002.
- Kakutani, T. and Yamasaki, N.: Solitary waves on a two-layer fluid, *J. Phys. Soc. Japan*, 45, 674–679, 1978.
- Logan, J. D.: Applied mathematics: a contemporary approach, Wiley, 1987.
- Miyata, M.: An internal solitary wave of large amplitude, *La mer*, 23, 43–48, 1985.
- 970 Miyata, M.: Long internal waves of large amplitude, in: Proceedings of the IUTAM Symposium on Nonlinear Water Waves, pp. 399–400, 1988.
- Ostrovsky, L. A. and Grue, J.: Evolution equations for strongly nonlinear internal waves, *Phys. Fluids*, 15, 2934–2948, 2003.
- Ostrovsky, L. A. and Stepanyants, Y. A.: Do internal solitons exist in the ocean?, *Rev. Geophys.*, 27, 293–310,
975 1989.
- Sandstrom, H. and Quon, C.: On time-dependent, two-layer flow over topography. I. Hydrostatic approximation, *Fluid Dyn. Res.*, 11, 119–137, 1993.
- Sandstrom, H. and Quon, C.: On time-dependent, two-layer flow over topography: II. Evolution and propagation of solitary waves, *Fluid Dyn. Res.*, 13, 197–215, 1994.
- 980 Stanton, T. and Ostrovsky, L.: Observations of highly nonlinear internal solitons over the continental shelf,

Geophys. Res. Lett., 25, 2695–2698, 1998.

Vlasenko, V., Stashchuk, N., and Hutter, K.: Baroclinic tides: theoretical modeling and observational evidence, Cambridge University Press, 2005.