

REPLY TO REVIEWER

I wish to thank the reviewer for his suggestions and for specifying what he meant in the first review. Reviewer observations are in italics.

1) I would agree that the phase portraits suggest a singly periodic behavior is dominant, although the small amplitude departures from a single limit-cycle trajectory do appear to show some short timescale periodicity which might be dynamically significant - though this component does not seem to show up in the spectra in Fig. 2. For the 23 day forcing period case the oscillations look to have a period near roughly a 1 day(?) period? But this is probably not a major issue and I am happy to leave this for the author to answer at his discretion.

1) Actually, I think that in this model, small differences are not dynamically important.

2) the additional tests carried out are useful in establishing the robustness of the bifurcation that is reported.

2) First, we must agree on a definition of the concept of robustness. Is it the maintenance of some desired system characteristics despite parameter fluctuations, or is it defined by how the model responds to changes in the architecture of the system (for instance solstitial vs equinoctial configurations)? Because the concept of robustness can be confusing (e.g., Jen 2003), I discussed sensitivity (in order to avoid being misleading) experiments for some parameters in the paper. In particular, I changed the values of Δ_H , which may alter the solution when the changes are large.

3) I was simply concerned that the model imposed a symmetry in the circulation about the equator which would only really apply either at equinoxes or for a planet whose obliquity was zero. So one could ask whether the same bifurcation would occur if this symmetry were broken e.g. to represent the circulation close to a solstice? Also, how significant is it that the forcing pushes the stratification precisely towards neutral stability at its extreme? What would happen if the minimum stability being forced was either weakly stable or even weakly unstable? These points are again concerned with establishing how robust the behavior might be and therefore how applicable generally the results might be to the real Earth (or other planet).

3) I performed simulations that depart from the symmetric conditions, and the results are discussed in the paper. Concisely, we can say that the breakdown of symmetry with respect to the equator changes the model response, especially for larger values of t_0 . Because the circulation is stronger in the solstitial configuration, static stability neutrality is not that important in creating spikes when compared to equinoctial experiments. However, the circulation is still large. Large changes in the static stability limit with respect to neutrality can be influential too, but small perturbations do not change the model's response.

All these new results were added to the section "Sensitivity to some parameters," where the results obtained with different vertical viscosity values were also described.

Jen, E. Stable or Robust? What's the Difference? Complexity, 8, 12-18.2003.

[<http://onlinelibrary.wiley.com/doi/10.1002/cplx.10077/pdf>]

Dynamics of the Hadley circulation in an axisymmetric model undergoing stratification periodic forcing

Nazario Tartaglione

Uni Research Climate and Bjerknes Centre for Climate Research, Bergen, Norway

Correspondence to: Nazario.Tartaglione@uni.no

Abstract. The time-dependent response of the Hadley circulation to a periodic forcing is explored via a simplified nonlinear axisymmetric model. Thermal forcing towards a given equilibrium potential temperature drives the model atmosphere. The vertical stratification of this temperature is forced to become periodically neutral with a period t_0 . Simulations performed with values of t_0 ranging from 10 to 90 days exhibit stronger circulation compared to the results of a constant thermal forcing experiment. As the period increases, a transition occurs first from a stationary regime, obtained when forcing is constant, to a periodic (and possibly quasi-periodic) regime, and then to an intermittent regime, albeit one with a strong periodic component. The stream-function response to periodic forcing is generally a periodic oscillation, with two main frequencies dominating, one with a period equal or close to the forcing period and another with a period that is half of the forcing period. The former is dominant for values of t_0 larger than 30 days, whereas the latter is prevalent for t_0 smaller than 30 days. The periodic oscillations obtained in this model may be associated with the periodic oscillations observed in the tropical regions. In this case the periodic charge and discharge of moisture in the tropical atmosphere, with consequent change of stratification, may be linked to those oscillations. In the model, at forcing periods of over 63 days the response of the stream-function periodically enters into a quasi-intermittent regime, exhibiting high frequency chaotic oscillations that are modulated by the slow timescale of forcing. ~~The vertical viscosity plays a role in determining the evolution of the Hadley circulation under the conditions established by the forcing~~ Sensitivity experiments for model parameters and configuration were performed to check whether results obtained are still valid under different conditions. Although for small changes of parameters the results are still valid, when parameters depart from the prescribed ones, we can observe change in the Hadley circulation dynamics.

1 Introduction

The Hadley cell is a specific and observable feature of tropical circulation (Dima and Wallace, 2003). The main driving process of this feature can be seen as a combination of low-latitude convective processes and interactions of tropical circulation with higher-latitude eddies (Charney, 1969). Although tropical circulation follows a seasonal cycle due to the asymmetry of solar forcing during a solar year (Fang and Tung, 1999), it is also subjected to alternating intra-seasonal strong and weak periods (Goswami and Shukla, 1984). The cycles modulate precipitation activity in the equatorial regions.

It is well known that the rainfall distribution over Indian tropical regions varies considerably from day to day with an alternating behavior associated with a few periodic or quasi-periodic cycles. Cycles of 3–7, 10–20 and 30–60 days are present

in the atmospheric dynamics of tropics and subtropics. The 3–7 day cycle is related to synoptic-scale convective systems generated over the Bay of Bengal, while the others have been associated with the cycles of monsoon rainfall over the Indian region as discussed by Kripalani et al. (2003) who showed a prevalence of the 10–20 day cycle during a normal monsoon year and of the 30–60 day cycle during a drought year.

5 The latter cycle is known as the Madden-Julian oscillation (Madden and Julian, 1972). Observational analyses and modeling studies revealed that there are dominant periods in the tropics and subtropics. Since the publication of that paper, tropical oscillations have been extensively studied with observations (Yasunari, 1979, 1980, 1981; Sikka and Gadgil, 1990, Yoneyama et al., 2013) as well as by means of models (Goswami and Shukla, 1984, Zhu and Hendon, 2015, Wang et al., 2016). The Madden-Julian oscillation has a significant impact on the Indian (Murakami 1976; Yasunari 1979) and Australian monsoons (Hendon
10 and Liebmann 1990). Kessler and McPhaden (1995) for instance suggested that it could play an important role in the onset and development of El Niño events. The relationship between this oscillation and tropical cyclogenesis has also been posited in some works (Maloney and Hartmann, 2000; Mo 2000; Maloney and Hartmann, 2001). Moreover, He et al. (2011) showed that Hadley circulation variability is closely related to Madden-Julian oscillation convection. Goswami and Shukla (1984) suggested that this oscillation in the Hadley circulation is due to the interaction between the internal dynamics of tropical circulation with
15 moist convection; in fact, with constant latent heat included in their model, the quasi-periodic oscillation vanished. This led them to consider that latent heat released during the moist processes can play a fundamental role in the dynamics of this cycle.

Tropical atmosphere dynamics can be explored via complex general circulation models as well as rather simpler models like the axisymmetric ones. Axisymmetric models focus on the main processes occurring in the tropospheric region, capturing the central process of the Hadley circulation. In such models, for example, eddies are not allowed, and all the processes involved
20 in the higher latitudes are not considered. In this paper, an axisymmetric model (Cessi, 1988) is used to perform an analysis of Hadley cell behavior when periodic forcing is applied. The model spin up takes less than 100 days. When the imposed forcing is not variable, the developing circulation results in a steady state with the Hadley cell representing a fixed point solution within the phase space. In this kind of configuration, other atmospheric processes, acting on longer or shorter spatial and temporal scales, like those mentioned earlier, are essentially excluded.

25 The Hadley circulation is the meridional overturn that develops in response to a temperature that is in radiative-convective equilibrium, and so it is worth analyzing how the model atmosphere behaves when the vertical stratification of equilibrium temperature becomes neutral. Although the equilibrium temperature towards which the model atmosphere evolves is essentially stable, a less stable temperature stratification may occur, example, due to tropospheric heating caused by condensation of water vapor, resulting in a less stable atmosphere. However, it is reasonable to assume that this condition occurs only periodically.

30 In Section 2 we describe the model we used and how we set up the stratification by modulating the vertical distribution of the temperature so as to periodically reach neutral static stability. In Section 3, we describe how modulation of the vertical stratification by a periodic function has an impact on the time evolution of the stream-function, which exhibits a periodic response of two main frequencies, suggesting that a periodic response is intrinsic to this model when stratification changes periodically. In this section, we also investigate [~~the role of the vertical viscosity~~model sensitivity for some parameters and
35 model configuration \(solstitial vs equinoctial\)](#). In Section 4 we present our conclusions.

2 Model equation and applied forcing

This paper employs the full model described in Cessi (1998), who studied its analytic and numerical solutions via a power series expansion in the Rossby number R of the prognostic variables, the zonal momentum M , the potential temperature θ , and the stream-function ψ . The model follows the same hypothesis of the classical axisymmetric models such as those used by

5 Held and Hou (1980) and Fang and Tung (1999).

The horizontal coordinate is defined as $y = a \sin \phi$, from which we have

$$c(y) = \cos \phi = \sqrt{(1 - y^2/a^2)}, \quad (1)$$

where a is the radius of a planet having a rotation rate Ω , with an atmosphere height H . The model is similar to the Held and Hou model, but it prescribes a horizontal and vertical viscosity, respectively equal to $\nu_H \nu_V$. The prognostic variables are

10 the angular momentum $M = \Omega a c^2 + u c$, where u represents the zonal velocity; the zonal vorticity ψ_{zz} where the meridional stream-function ψ is defined by

$$\begin{aligned} \partial_y \psi &\equiv w, \\ \partial_z \psi &\equiv -cv, \end{aligned} \quad (2)$$

and the potential temperature θ that is forced towards a radiative–convective equilibrium temperature θ_E .

15 Starting from the dimensional equations of the angular momentum, zonal vorticity and potential temperature, we obtain a set of dimensionless equations. The new equations are non-dimensionalized using a scaling that follows Schneider and Lindzen (1977), but the zonal velocity u is scaled with Ωa . A detailed description can be found in Cessi (1998).

The non-dimensional model equations are

$$M_t = \frac{1}{R} \left\{ M_{zz} + \mu \left[c^4 (c^{-2} M)_y \right]_y \right\} - J(\psi, M), \quad (3a)$$

$$\begin{aligned} 20 \quad \psi_{zzt} &= \frac{1}{(R^2 E^2)} y c^{-2} (M^2)_z - \frac{1}{c^{-2}} J(\psi, c^{-2} \psi_{zz}) \\ &+ \frac{1}{(R E^2 c^{-2})} \theta_y + \frac{1}{(R c^{-2})} [c^{-2} \psi_{zzzz} + \mu \psi_{zzyy}], \end{aligned} \quad (3b)$$

$$\theta_t = \frac{1}{R} \left\{ \theta_{zz} + \mu \left[c^2 \theta_y \right]_y + \alpha [\theta_E(y, z) - \theta] \right\} - J(\psi, \theta). \quad (3c)$$

where the term $J(A, B) = A_y B_z - A_z B_y$ is the Jacobian.

The thermal Rossby number R , the Ekman number E , the ratio of the horizontal to the vertical viscosity μ and the parameter α are defined as

25

$$\begin{aligned} R &\equiv g H \Delta_H / (\Omega^2 a^2); \quad E \equiv \nu_V / (\Omega H^2); \\ \mu &\equiv (H^2 / a^2) \nu_H / \nu_V; \quad \alpha \equiv H^2 / (\tau \nu_V). \end{aligned} \quad (4)$$

The term α is the ratio of the viscous timescale across the depth of the model atmosphere to the relaxation time τ toward radiative-convective equilibrium.

The boundary conditions for the set of Eq. (3) are

$$\begin{aligned} M_z &= \gamma (M - c^2), \quad \psi_{zz} = \gamma \psi_z, \\ \psi &= \theta_z = 0 \text{ at } z = 0, \\ M_z &= \psi_{zz} = \psi = \theta_z = 0 \text{ at } z = 1, \end{aligned} \tag{5}$$

- 5 where $\gamma = \frac{CH}{\nu_\nu}$ is the ratio of the spin-down time due to drag to the viscous timescale; the bottom drag relaxes the angular momentum M to the local planetary value $\Omega a c^2$ through a drag coefficient C . The model forcing is represented by an equilibrium potential temperature θ_E defined by

$$\theta_E = \frac{4}{3} - y^2 + \frac{\Delta_V}{\Delta_H} \left(z - \frac{1}{2} \right). \tag{6}$$

- A simulation with the forcing given by Eq. 6 leads to a stationary situation where the Hadley cell is a fixed point for the system of Eq. (3). We refer to this simulation as the control experiment. The parameters used in the simulations we describe in this paper are shown in Table 1.

In order to change vertical stratification we can define θ_E as an exponential function of z :

$$\theta_E = \frac{4}{3} - |y|^n + \frac{\Delta_V}{\Delta_H} \left(z^k - \frac{1}{2} \right).$$

- This approach was used by Tartaglione (2015), who used constant k values, to investigate the role of vertical stratification on the strength and position of the Hadley circulation. The response of the model atmosphere to vertical stratification will be defined by variations of the exponent k in the forcing equation as a function of time. In general, the vertical stratification can change the intensity of the Hadley cell only when the forcing is concentrated on the equator, whereas it loses importance when the forcing is represented by a weak meridional gradient in the equilibrium temperature (Tartaglione, 2015).

- Thus, we analyze the response of our model to an equilibrium temperature that moves periodically from a stable stratification to a neutral one; i.e. we allow exponent k of Eq. (6) as a function of time and we assume that $n = 2$. Thus Eq. (6) takes the form:

$$\theta_E = \frac{4}{3} - |y|^n + \frac{\Delta_V}{\Delta_H} \left(z^{1+\sin \frac{2\pi t}{t_0}} - \frac{1}{2} \right). \tag{7}$$

- When the exponent is zero the thermal forcing has an adiabatic stratification with no variation of the potential temperature along the vertical direction. The parameter t_0 takes a value less than or equal to 90 days, and defines the time in which the forcing becomes adiabatic, forcing the model atmosphere to become neutral. When t_0 is small, the main mechanisms involved in the change of the static stability are associated with short time convective events, whereas large values of t_0 can be related to longer-lasting meteorological events like monsoons. Whatever process is involved in changing the vertical stratification, we assume in our model that it is periodic.

- A cautionary note is necessary. The main role of convection is to bring the atmosphere into a state of neutral vertical stratification, deleting the effects of the unstable layer created by radiative only processes. Thus the dry unstable condition is

Table 1. Values of the parameters used in this work

Parameter	Value	Formula
H	$8 \cdot 10^3 m$	
Δ_H	$1/3$	
Δ_V	$1/8$	
C	$5 \cdot 10^{-3} m^2 s^{-1}$	
ν_H	$5 m^2 s^{-1}$	
ν_V	$1.86 m^2 s^{-1}$	
τ	$1.728 \cdot 10^6 s$	
a	$6.4 \cdot 10^6 m$	
g	$5 \cdot 10^{-3} m s^{-2}$	
Ω	$2\pi/(8.64 \cdot 10^4) s^{-1}$	
R	0.121	$gH\Delta_H/(\Omega^2 a^2)$
E	$1.07 \cdot 10^{-3}$	$\nu_V/(\Omega H^2)$
α	19.9	$H/(\tau \nu_V)$
Δ	$3/8$	Δ_V/Δ_H
γ	25500	CH/ν_V
μ	$4.2 \cdot 10^{-6}$	$(H^2/a^2) \cdot \nu_H/\nu_V$

almost never met in the real atmosphere as there is always an overturn that leads the atmosphere to be statically stable and the most prevalent instability is the conditional stability due to the presence of water vapor. However, the neutral condition of θ_0 represents for this model a less stable condition compared with the θ_0 prescribed in the control simulation (Eq. 6). We parameterize the contribution of the water cycle in the model by means of a time function of θ_0 in such a way that the model

5 atmosphere becomes alternately more or less stable.




Figure 1. Temporal evolution of the stream-function at 3° North and 2400 m (a,c,e) and its power spectral density (b,d,f) when $t_0 = 10$ (a,b), 23 (c,d) 60 days (e,f) and 90 days(g,h). The figures on the bars are the cycle periods in days.

3 Results

3.1 Model response to periodic forcing

Using Eq. (7) as forcing, we obtain a vertical stratification of the equilibrium temperature that becomes periodically neutral. The values of t_0 are chosen so as to explore the rate of vertical stratification change, and consequently, the effects on the static stability. The value of t_0 chose lies in the range of 10 to 90 days. The aim of these simulations is to relax the atmosphere to neutral stability, but leave the time average of vertical lapse rate equal to that used in the literature, i.e. equal to one (see Eq. 6).

We now take a closer look at the details in a specific point of the domain, that corresponding to 3° N and 2400 m of altitude. The choice of this specific point is based on the fact that the dynamics of this model are essentially equatorial, and that point is in the ascending branch of the cell and its signal in term of spectrum power is higher than that obtained where the value of the stream-function is a maximum, at about 9° N and 1600 m of altitude.

The time evolution of the stream-function, at 3° N and 2400 m, and its power spectrum for $t_0 = 10$ days are shown in Figs. 1a,1b. It is easy to see that the evolution resembles a periodic solution with more peaks in the spectrum; however, the dominant frequency is one corresponding to the period of 5 days, i.e., half of the forcing period. As we will see, as the time period increases, the second component related to the time forcing becomes more important. Figures 1c and 1d show the temporal evolution of the stream-function at a point in the model domain, and the spectral density, for $t_0 = 23$ days. When the forcing period is a prime number, the quasi-periodicity is formally present, as for this value of t_0 the two main frequencies are incommensurate. When $t_0 = 30$ days the solution is still periodic with a main period of 30 days, although there is a signal even at 15 days. This behavior is the same at least when $t_0 = 60$ days (Figs. 1e,1f). The peak in the spectrum corresponding to a period of 60 days is higher than that corresponding to the half forcing period.

If we look at the phase space evolution and the plot of couples($\psi(t); \psi(t+t_0)$), we can see that for both $t_0 = 23$ days and $t_0 = 60$ days (Fig. 2), the solution does not lie on the same curve, but rather assumes a behavior that appears quasi-periodic. As the solution is essentially numeric, this behavior could be related to rounding errors rather than a real quasi-periodic solution.

Time evolution of the solution, for $t_0 = 90$ days, is shown in Figs. 1h and 1g. Looking at the time evolution of the stream-function, the motion is chaotic only when the time is close to multiples of $t_0 = 90$ days, whereas it returns to the steady solution when the stratification θ_E is far away from the condition of neutral stability. Thus, the slow variation with time, because of the high value of t_0 , triggers a fast response in the model that allows instabilities to grow faster than those obtained by using small values of t_0 . Therefore, we observe that there is an unusual sort of slow-fast dynamic. This phenomenon can be viewed as a sort of intermittency with the slow component, which modulates the fast process.

Thus, if the adiabatic forcing is reached with a relatively fast change of the stratification, the solution follows adiabatic forcing, increasing the strength of the circulation (Fig. 3a), which leads to strong subtropical jet streams and stronger easterly winds in the equatorial region (Fig. 3b). As in Fang and Tung (1999), who found stronger circulation when they replaced a fixed sun (equinoctial Hadley cell) with a moving sun, here it is the time-varying stratification stability of θ_0 that causes a stronger circulation with respect to a fixed lapse rate.

Periodicity with two dominant cycles in the model response is interesting in light of the oscillations observed in the tropical atmosphere. Madden and Julian (1971,1972) were the first to show the existence of an oscillation in pressure and winds with a predominant peak in the spectrum at a period of 40-50 days. They also showed that the amplitude of this peak was greater in the tropical station and was weaker in the sub-tropical stations. Yasunari (1979) demonstrated by means of spectral analysis that cloudiness fluctuations have two dominant periodicities: one of about 15 days and another of 40 days. Other studies documented 15-day oscillations within the tropical regions related to monsoons (e.g. Krishnamurti and Bhalme, 1976; Krishnamurti and Ardanuy, 1980; Krishnamurti and Subrahmanyam, 1982). Yasunari (1981) showed that even the 40 day oscillation has some relation to the Asian summer monsoon. Anderson and Rosen (1983) found similar results by using zonally averaged zonal winds. Thus, the features of these oscillations suggest that it may be possible to understand them with a zonally averaged model. Goswami and Shukla (1984) used a symmetric model with hydrology to study the Hadley circulation and found that it has well-defined strong and weak episodes. These oscillations of the Hadley circulation occurred in their model in two dominant ranges of periodicities: one with a period of between 10 and 15 days and another with a period of between 20 and 40 days. Since our model does not include hydrology, this double period has to be related to the internal dynamics of the system. In fact, if periodicity is expected with a time period equal to the forcing time period t_0 , the response with cycles with more frequencies has to come from the interaction between changing static stability and the internal dynamics of the model. As the changing stratification stability implies a way to simulate the moist convection, our result seems to be in agreement with the findings of Goswami and Shukla (1984). They found that quasi-periodic oscillations of the Hadley circulation were seen only when the moist convective heating was turned on in their model.

The time series of the vertically-averaged stream-function shows that the observed periodic response involves mainly low-level processes. Figure 4a shows the time series of the stream-function averaged over the lower 3200 m, while Fig. 4b shows it averaged over all the domain height. In both cases strong and weak patterns in the stream-function are present, and the strong




Figure 2. Representation of the evolution of the stream-function at 3° North and 2400 m $\psi(t)-\psi(t+t_0/8)$ (left panels) for $t_0 = 23$ days (upper panels) and $t_0 = 60$ days experiments (lower panels).




Figure 3. Difference between the stream-function (in dimensionless units) of the experiment with $t_0 = 50$ days, and the control experiment (a), the zonal wind U (in ms^{-1}) averaged on the height as a function of latitude y (b), for the experiment $t_0 = 50$ days (in red) and the control experiment (in black) after 500 simulation days.




Figure 4. Time evolution of the vertically averaged stream-function (non-dimensional unit) over lower levels, up to 3200 m (a) and over the entire height of the domain (b), for the experiment with $t_0 = 60$ days.

episodes of the stream-function at lower levels do not dump suddenly, but persist for a while. However, when averaged over the entire domain time series shows strengthening and weakening of the stream-function that seem to be periodic. This periodic behavior tends to become intermittent with increasing t_0 . Higher values of t_0 mean that the model atmosphere takes more time to become neutral, while at the same time remaining close to the neutral stability, which induces a perturbation that produces a dramatic increase in the stream-function values.

The interaction of slow parameter variation with the fast rate of motions in the phase space causes a phenomenon known as "dynamic bifurcation" (Guckenheimer and Holmes, 2002). Figure 5 shows a one-dimensional bifurcation diagram, i.e., the differences between two stream-functions at the same point of the domain (3° N and 2400 m of altitude) having a time lag of t_0 days, plotted for each value of t_0 . The outliers, indicating chaotic behavior, start to appear at $t_0 = 64$ days, indicating the presence of spikes in the stream-function values associated with a chaotic solution. Thus, the periodic behavior starts losing force by allowing chaos to emerge when the z exponent approaches zero.

Periodicity is still present, in the sense that chaotic behavior of the model appears periodically. This occurs, for instance, for $t_0 = 90$ days (Figs. 1g, 1h). In some respects we can say that the low variability is still governed by periodic oscillation, with the presence of chaotic behavior associated with the neutral static stability of the forcing. Since the transition occurs at $t_0 = 64$ days, which is in the 50-60 day range of the Madden and Julian oscillation, the question arises whether results of this work imply that the Madden-Julian oscillation also has chaotic components. Unfortunately, this is not easy to determine with real data as high frequency components are usually removed in order to study the Madden-Julian oscillations, and because of high non-linearity of the atmosphere where chaotic features are normal.




Figure 5. The 1-D bifurcation diagram as a function of t_0 . At a period of 63 days, deviations from periodic behavior start to appear. The plot shows the differences of two stream-functions (in dimensionless units) at the same point of the domain, but with a time lag of t_0 days. The value of the vertical viscosity is set to $5 \text{ m}^2 \text{ s}^{-1}$.

It seems that a bifurcation delay might be active when t_0 is longer than 63 days. The system fails to notice the onset of instability. This is quite evident in Figs. 1h and 1g, where it is clear that, most of the time, the low frequency (period of 90 days) component is the dominating one; however, a fast component is observed when the chaotic behavior is triggered, a result obtained even with simpler models (e.g. Zaks et al., 2002). Having reached a chaotic state, the system wanders along this state for a time, until it finds itself with the exponent k , which defines the slow component of the system, having a value "far enough" from zero that the system goes back to steady solution. At this time, self-modulation is temporarily switched off until the cycle is repeated.





Figure 6. As in Fig. 1g, but halving the time step, and increasing the spatial resolution, i.e. halving the horizontal (b), vertical (c) and both (d) grid steps..

We can say that identification of dynamic bifurcations caused by slow variation can, in general, be a problematic task, because of transitions. During these transitions two situations could occur: bifurcations with abrupt change of the attractor size (and in such a case the dynamic bifurcation could be visible), or transitions occurring with changes in the geometry of the chaotic attractor. In the latter case it may be difficult to observe these variations in the short time that the system is in the

5 chaotic state.

3.2 Model Sensitivity to time and grid steps

One can argue that the amplification seen in the model solution when t_0 equals 90 days may be an artifact of the model. In order to eliminate this possibility, some simulations were performed in order to assess the model sensitivity to time and grid steps. Figure 6 shows the temporal evolution of the stream-function at 3° North and 2400 m (as in Fig. 1), obtained by halving the time step (Fig. 6a), doubling the horizontal (Fig. 6b) and the vertical (Fig. 6c) resolutions and halving all the spatial grid steps (Fig. 6d). In all these cases, albeit with some slight differences, there is an explosive increase of the stream-function value at some point, with the most remarkable difference seen when the vertical resolution is increased. There is still periodic strengthening of the Hadley circulation, but its variability is essentially reduced with the negative spikes vanishing. Although not shown here, these results hold when the time step and the spatial grids are even further reduced.






Figure 8. Temporal evolution of the stream-function (in dimensionless units) at 3° North and 2400 m of altitude for a simulation where t_0 changes with time.

To investigate whether history affects the value of the internal state of our model we performed a simulation where we changed the value of t_0 during the run from 10 to 80 days, and back again to 10 days (Fig. 8). The model adjusted its response, which is the same of that obtained starting from an initial condition at rest, almost immediately, even though the change of t_0 was sudden. This result shows that the model response does not depend on history and on the initial conditions. Moreover the quick adjustment of the model suggests that the fluctuations observed in the simulation are an intrinsic characteristic of it and not an artifact, suggesting that the model, at the least in this form is resilient.

3.3 ~~Role of the vertical viscosity~~Model Sensitivity to some parameters

Although a parameter uncertainty analysis similar to that presented by Knopf et al. (2006) is outside of the aims of this paper, a sensitivity analysis for some parameters was performed to establish whether found features persist under different conditions.




Figure 9. The time evolution of stream-function (in dimensionless unit) in the ascending branch of the circulation for $t_0 = 30$ (a) and 90 (b) days with the model in solstitial configuration.

In previous experiments, static stability is forced toward a specific minimum limit, such as neutrality, but whenever it is forced toward values that differ slightly from the neutral condition, the solution behavior remains unaltered.

When the change is significant (e.g., 20% higher or lower than the neutral condition), we need to distinguish the model response as a function of t_0 . With low values of t_0 , the periodic solution is independent from the static stability limit. As t_0 increases, the model becomes more sensitive to the minimum static stability to which the system is forced. In such a case, a more stable minimum prevents the appearance of spikes in the model solution; however, an unstable static stability limit leads to completely chaotic behavior with the appearance of secondary cells far from the equator.

The sensitivity to the parameter Δ_H is also tested, and the results show that, independently on t_0 , model response depends on this value. If the parameter is close to $1/3$ as set in the previous experiments, we cannot find any difference, but when it




Figure 10. The time evolution of stream-function (in dimensionless unit) at 3° North and 2400 m for $t_0 = 10$ (a), 30 (b), 60 (c) and 90 (d) when $\nu_V = 0.5$.

departs significantly from that value we notice important changes in the solution. The model response is more stable for low values of Δ_H and exhibits strongly periodic behaviors. Large values of Δ_H make the model unstable with a completely chaotic response.

- The model configuration has symmetry about the equator. To determine how the model responds to the loss of this
- 5 symmetry, we performed solstitial experiments with the same parameters used for the equinoctial (symmetric) configuration with maximum heating located in a hemisphere at 6° from the equator. Figure 9 shows the stream function value at one point (2400 m of altitude) in the ascending branch of the single cell forms for experiments with t_0 values of 30 and 90 days. For low values of t_0 , the solution is still periodic. When $t_0 = 90$ days, the spikes present in the equinoctial configuration disappear, but the stream function, although noisier, is similar, in term of evolution, to that at $t_0 = 30$ days. The main reason for this
- 10 change may be the strength of the circulation in the solstitial configuration, which is much greater than that of the equinoctial configuration.




Figure 11. [The 1-D bifurcation diagram as a function of \$\nu_v\$ for simulations with \$t_0 = 30\$ days.](#)

The periodic chaotic behavior is [even](#) related to vertical viscosity. It appears even for time periods less than 63 days, when the vertical viscosity is close to zero. When the vertical viscosity is $0.5 \text{ m}^2 \text{ s}^{-1}$, i.e. one tenth of that used previously, the appearance of the chaotic behavior is immediate for small values of t_0 , as we can see when $t_0 = 10$ days (Fig. [9a10a](#)). When $t_0 = 30$ days, the solution is chaotic for almost each of the 30 days (Fig. [9b10b](#)), and the same occurs when $t_0 = 60$ days (Fig. [9c10c](#)). In such a case, the evolution of the stream-function will have a steady solution with intermittent chaotic behavior when the static stability gets close to the neutral condition, especially for higher t_0 . Figure [9d-10d](#) illustrates the situation for $t_0 = 90$ days. This representation is only intended to show the effect of the vertical viscosity which must be taken into account when we consider the development and evolution of the stream-function. The high values of the non-dimensional stream-function appear to be unrealistic, but we cannot exclude the possibility transport of momentum and heat from the equator could occur by means of bursts (Majda and Stechmann, 2008). If we wish to estimate for a specific value of t_0 at which level of ν_v this

occurs Fig. 10-11 gives us insight as to what happens for different values of vertical viscosity, for example when $t_0 = 30$ days. For values less than $2 \text{ m}^2 \text{ s}^{-1}$ the systems begin to exhibit some spikes in the solutions associated with a chaotic behavior of the system. For higher values of t_0 , the amplification of the model response occurs for higher values of ν_V . On the other hand, high values of vertical viscosity kill the chaotic behavior, making the evolution periodic (not shown).

5 The time evolution of stream-function (in dimensionless unit) at 3° North and 2400 m for $t_0 = 10$ (a), 30 (b), 60 (c) and 90 (d) when $\nu_V = 0.5$.

The 1-D bifurcation diagram as a function of ν_V for simulations with $t_0 = 30$ days.

4 Conclusions

We have used a dry axisymmetric model to simulate the Hadley circulation and to investigate the role of a changing stratification of the thermal forcing, which simulates moist convection that alters the static stability of the model atmosphere. The bi-dimensionality of the model prevents the generation of any eastward traveling wave. Hence, in our discussion, the influence of eddies' momentum fluxes is not taken into account. We have shown that the stream-function representing the Hadley circulation in an axisymmetric model can exhibit a periodic behavior when the vertical stratification of the thermal forcing is periodically forced to become neutral. The question that arises is whether this oscillation can be linked with the observed atmospheric fluctuations within the tropical region. Although Madden-Julian oscillation is a three dimensional phenomenon with the development of a Kelvin-Rossby wave (Gill, 1980), it is certainly associated with the evolution of convective anomalies (Hendon and Salby, 1994). It has already been suggested that quasi-periodic oscillation seems to be an intrinsic characteristic of the tropical atmosphere, in accordance with the results of Goswami and Shukla (1984). Bi-dimensional models, while they can give us a framework of basic physics underlying atmospheric processes are quite limited as the real atmosphere is naturally three-dimensional. The findings of this work suggest that if a cyclic process perturbs tropical stratification, the Hadley circulation strength may be periodic with possible bursts when the perturbation time is larger than 60 days. These results must necessarily be compared with observations of the real atmosphere or to the results of three-dimensional models.

If the forcing period is up to 63 days the stream-function evolution shows a periodic behavior. For period forcing longer than 63 days, the slow frequency associated with the forcing period modulate a fast response in the system, generating a chaotic motion that persists for a period of time before returning to a non-chaotic solution. It is not clear whether these high frequency characteristics are actually present in the meridional circulation of our planet. In fact, when we look for periodicities of the order of tens of days in observations, the higher frequency signals are usually removed. Moreover, even though we can detect such signals, it is not easy to associate them with the oscillations caused by a change of static stability, as opposed to other processes. The chaotic dynamics observed in the model could exist in other planets where the vertical stratification takes longer to become neutral. The change of vertical stability that in our model simulates the cycle of large-scale convection might be equivalent to the recharge and discharge of moisture that supports the Madden-Julian oscillation (Zhu and Hendon, 2015). This change of stability can be imagined to control the aggregation process of convection, which allows a bistable equilibrium between moist and dry situations (Raymond and Zeng, 2000, Zhang et al., 2003, Arnold and Randall, 2015). An important

aspect is the rate at which this process occurs. As we have shown in the model we consider, when this rate is over 63 days it results in the short term in a chaotic impact on the Hadley circulation strength.

Many parameters control the numerical solution for the model presented in this paper. We have analyzed model sensitivity to a few of these parameters with imposed stratification periodic forcing. The forcing imposes static stability to move toward neutrality, and the model's response does not change if static stability remains close to neutral, but when it departs from neutrality, there is no change in the model solution for low values of t_0 . For higher values of t_0 , however, the model response is strongly affected by static stability. Low static stability prevents chaotic behavior, but higher static stability triggers chaotic motion. Moreover, even Δ_H is important for higher values of t_0 and has the same effects discussed earlier with regard to static stability: low values of Δ_H minimize chaotic behaviors, whereas high values of Δ_H tend to trigger very chaotic responses.

The model in the equinoctial configuration shows intermittent chaotic behavior when the forcing period is large because of the importance of the stratification forcing imposed on the model atmosphere. However, this static stability becomes less important in creating large spikes in the stream-function when a solstitial configuration is adopted. With a single strong cell, the circulation is more stable.

The role of friction in the symmetric circulation, driven by a meridional thermal gradient of a fast rotating planet like the Earth, is contradictory. On the one hand it is an essential ingredient to allow a meridional overturn, instead of a strong zonal wind in cyclostrophic balance only. On the other hand, the value must be close enough to zero to allow angular momentum conservation. Although these conditions are met in the model we consider, the presence of a time-varying stratification alters the classic view of a stably stratified vertical temperature gradient. Other than the meridional thermal gradient, this imposed time-varying stratification represents another nonlinear forcing, which amplifies the model response when the vertical viscosity is small, itself representing a source of amplification of the model response in the inviscid case.

Acknowledgements. The author wishes to thank two anonymous reviewers who were willing to review this paper. One of these provided useful suggestions to remarkably improve the paper.

References

- Anderson, J.R., and Rosen, R.D.: The latitude–height structure of 40–50 day variation in atmospheric angular momentum. *J. Atmos. Sci.*, 41, 1584–1591, 1983.
- Arnold, N. P., and Randall, D.A: Global-scale convective aggregation: Implications for the Madden-Julian oscillation, *J. Adv. Model. Earth Syst.*, 7, 1499–1518, doi:10.1002/2015MS000498, 2015.
- Charney, J.G.: "The Intertropical Convergence Zone and the Hadley Circulation of the Atmosphere" Proceedings of the WMO-IUGG Symposium on Numerical Weather Prediction, Tokyo, Japan, November 26–December 4 , 1968, Japan Meteorological Agency, Tokyo, 111–73–111–79, 1969.
- Cessi, P.: angular momentum and temperature homogenization in the symmetric circulation of the atmosphere, *J. Atmos. Sci.*, 55, 1997–2015, 1998.
- Dima, I. M. and Wallace, J. M.: On the seasonality of the Hadley cell, *J. Atmos. Sci.*, 60, 1522–1527, 2003.
- Fang, M. and Tung, K. K.: Time-dependent nonlinear Hadley circulation, *J. Atmos. Sci.*, 56, 1797–1807, 1999.
- Goswami, B.N., and Shukla, J.: Quasi-periodic oscillations in a symmetric general circulation model. *J. Atmos. Sci.*, 41, 4120–4137, 1984.
- Guckenheimer, J., and Holmes, P.: *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields*, Springer, 1983.
- He, J., Lin, H., and Wu, Z.: Another look at influences of the Madden-Julian Oscillation on the wintertime East Asian weather. *J. Geophys. Res.* 116, D03109. doi:10.1029/2010JD014787, 2011.
- Held, I.M., and Hou, A.Y.: Nonlinear axially symmetric circulation in a nearly inviscid atmosphere. *J. Atmos. Sci.* 37, 515–533, 1980.
- Hendon H.H., Liebemann, B.: The intraseasonal (30–50 day) oscillation of the Australian summer monsoon. *J. Atmos. Sci.*, 47, 2909–2923, 1990.
- Hendon, H.H., and Salby, M.L.: Life cycle of the the Madden Julian oscillation. *J. Atmos. Sci.*, 51, 2225–2237, 1994.
- Kessler, W.S., and McPhaden, M.J.: Oceanic equatorial waves and the 1991–93 El Niño. *J. Climate*, 8, 1757–1774, 1995.
- [Knopf, B., Flechsig, M., and Zickfeld, K.: Multi-parameter uncertainty analysis of a bifurcation point, *Nonlin. Processes Geophys.*, 13, 531–540, doi:10.5194/npg-13-531-2006, 2006.](#)
- Krishnamurti, T.N., and Ardanuy, P.: The 10–20–day westward propagating mode and breaks in the monsoon. *Tellus*, 32, 15–26, 1980.
- Krishnamurti, T.N., and Bhalme, H.N.: Oscillations of a monsoon system. Part I. Observational aspects. *J. Atmos. Sci.*, 33, 1937–1954, 1976.
- Krishnamurti, T.N., and Subrahmanyam, D.: The 30–50 day mode at 850mb during MONEX. *J. Atmos. Sci.*, 39, 2088–2095, 1982.
- Kripalani, R.H., Ashwini Kulkarni, A., Sabade, S.S., Revadekar, J.V., Patwardhan S.K., and Kulkarni, J.R.: Intra–seasonal oscillations during monsoon 2002 and 2003 *Current Science*, 87, 325–331., 2004.
- Madden, R., and Julian, P.: Detection of a 40–50 day oscillation in the zonal wind in the tropical Pacific. *J. Atmos. Sci.*, 28, 702–708, 1971.
- Madden, R., and Julian, P.: Description of global-scale circulation cells in the tropics with a 40–50 day period. *J. Atmos. Sci.*, 29, 1109–1123, 1972.
- Madja, A.J., and Stechmann S.N.: Stochastic models for convective momentum transport. *PNAS*, 105, 17614–17619, 2008.
- Maloney, E.D., and Hartmann, D.L.: Modulation of eastern north Pacific hurricanes by the Madden-Julian oscillation. *J. Climate*, 13, 1451–1460, 2000.
- Maloney E.D., and Hartmann D.L.: The sensitivity of intraseasonal variability in the NCAR CCM3 to changes in convective parameterization. *J. Climate*, 14, 2015–2034, 2001.
- Murakami T.: Analysis of summer monsoon fluctuations over India. *J. Meteorol. Soc. Japan*, 54, 15–31, 1976.

- Mo K.C.: Intraseasonal Modulation of Summer Precipitation over North America. *Mon. Wea. Rev.* 128, 1490–1505, 2000.
- Raymond, D. J., and Zeng X.: Instability and large-scale circulations in a two-column model of the tropical troposphere, *Q. J. R. Meteorol. Soc.*, 126, 3117–3135, 2000.
- Sikka, D.R., and Gadgil, S.: On the maximum cloud zone and the ITCZ over Indian longitudes during the south-west monsoon. *Mon. Wea. Rev.* 108, 1840–1853, 1980.
- 5 Tartaglione, N.: Equilibrium temperature distribution and Hadley circulation in an axisymmetric model. *Nonlinear Proc. Geoph.*, 22, 173–185, 2015.
- Yasunari, T.: Cloudiness fluctuations associated with the Northern hemisphere summer monsoon. *J. Meteorol. Soc. Japan*, 57, 227–242, 1979.
- Yasunari, T.: A quasi-stationary appearance of 30 to 40 day period in the cloudiness fluctuations during the summer monsoon over India. *J. Meteorol. Soc. Japan*, 58, 225–229, 1980.
- 10 Yasunari, T.: Structure of the Indian monsoon system with around 40-day period. *J. Meteorol. Soc. Japan* 59, 336–354, 1981.
- Yoneyama, K., Zhang, C., and Long, C.N.: Tracking pulses of the Madden–Julian oscillation. *Bull. Am. Meteorol. Soc.*, 1871–1891, doi:10.1175/BAMS-D-12-00157.1, 2013.
- Zaks, M.A., Park, E.-H., Kurths, J.: Self-induced slow-fast dynamics and swept bifurcation diagrams in weakly desynchronized systems. *Phys Rev E*, 65, 026212 1–5, 2002.
- 15 Zhang, C., Mapes, B. E. and Soden B. J.: Bimodality in tropical water vapour, *Q. J. R. Meteorol. Soc.*, 129, 2847–2866, 2003.
- Zhu H., and Hendon, H.H. : Role of large scale moisture advection for simulation of the MJO with increased entrainment, *Q. J. R. Meteorol. Soc.*, 141, 2127–2136, doi:10.1002/qj.2510, 2015.