

## Responds to Anonymous Referee 1

A few tel-conferences of all co-authors have been held to discuss the comments from reviewer #1. All authors converge to the point that all the comments are very important and useful for authors to improve the quality of this manuscript (MS). Therefore, all comments from reviewer #1 have been fully  
5 addressed in the revision.

Now we will reply to each comment point by point as following:

1 the title does not fit. You are not discussing the “Impact of Optimal Observational Time Window’ but  
10 “Impact of Observational Time Window”. It may be “Impact of Observational Time Window in Coupled Data Assimilation with a Simple Climate Model”.

RE: Thank you very much for your generous comment and we fully agree with that. We have corrected the tile as “Impact of Observational Time Window on Coupled Data Assimilation: Simulation with a Simple Climate Model”.


15

2 The concept of OTW in this study validates the observations in a time window to the center of the window (analysis time). It is very useful technique for data assimilation (DA). More or less it has been applied in the assimilation with real observations. Here the OTW in this study is applied in 3-dimension DA but not 4-dimension DA, which need address in the introduction. The citations of OTW are not very  
20 relevant (Page 2, line 13-15). I could not find the clear concept of OTW and how they applied in data assimilation. So you should address the technique details in the paper. I guess that you treat all the observations independently and assimilate then sequentially using their original error scales. another approach is to assimilate the average values of the observations in the OTW. It is worth to compare these two approach in the study.

25 RE: As Line 19-23 of Page 2, we have rewritten this part and addressed that in this study the OTW validates the observations including in a time window to the center of the window and are sequentially assimilated using their original error scales. The irrelevant citations have been deleted and we have added some ideas in a relevant one to introduce the concept of OTW using in this study.

As Line 8-18 of Page 12 in the revised manuscript, we have added some discussion about the comparison of sequential assimilation and averaged-observational assimilation approaches. Thanks.

Here the standard deviation of the observations will be  $1/\sqrt{N}$  of the original error scales (N represents the number of the observations including in the OTW). And the results of comparing of their two methods are shown as follow:



Here the Sqe-CDA-OTW represents the experiment that sequentially assimilates the observations including in the OTWs using their original error scales. And the Ave-CDA-OTW denotes the experiment that assimilates the average of the observations including in the OTWs. But the standard deviation of the observational error will be  $1/\sqrt{N}$  of the original error scales. Here all the experiments will use the biased model setting and the uni-variate adjustment scheme.

We can see that the results of the Ave-CDA-OTW is almost same as those of the Sqe-CDA-OTW. As follow we will illustrate this phenomenon from the perspective of the calculation of the EAKF method.

Here we assume the OTW including two observation  $a$  and  $b$ ,

- 1) Sequentially assimilate these two observations using their original error scales:

$$\Delta Z_{i,j} = \frac{Cov(Z_j, Y_k)}{(\sigma_{k,k}^p)^2} \left[ \left( \frac{\bar{y}_k^p}{1+r_k^2} + \frac{y_k^o}{1+r_k^{-2}} + \frac{\Delta y_{k,i}^p}{\sqrt{1+r_k^2}} \right) - y_{k,i}^p \right]$$

in this study there is no inflation scheme, thus after many assimilation steps, the ensemble spreads of the model states will greatly decreased will be greatly smaller than the observational error. Thus we can

assume  $\bar{y}_k^p \approx y_{k,i}^p$  and  $\frac{Cov(Z_j, y_k)}{(\sigma_{k,k}^p)^2} \approx \text{constant } r$ . For simplicity and convenience, we assume at a assimilation time the  $\bar{y}_k^p$ ,  $\sigma_{k,k}^p$  and  $\Delta y_{k,i}^p$  will keep as constants during the sequential data assimilation process. And we can use constants  $C_1$ ,  $C_2$  and  $C_3$  to represent the  $\bar{y}_k^p$ ,  $r_k^2$  and  $\Delta y_{k,i}^p$ , respectively.

$$Z_{i,j} = \frac{1 + (1-r)C_2}{1 + C_2} \left[ \frac{1 + (1-r)C_2}{1 + C_2} C_1 + \frac{rC_2}{1 + C_2} a + \frac{r}{\sqrt{1 + C_2}} C_3 \right] + \frac{rC_2}{1 + C_2} b + \frac{r}{\sqrt{1 + C_2}} C_3$$

$C_2$  is very small and we can assume that  $C_2^2 \approx 0$ . Thus

$$Z_{i,j} \approx \frac{1 + 2(1-r)C_2}{1 + 2C_2} C_1 + \frac{rC_2}{1 + 2C_2} (a + b) + \frac{2r + 2rC_2 - r^2C_2}{\sqrt{(1 + C_2)^3}} C_3$$

5 2) assimilate the average of the observations but  $C_2' = (r_k')^2 = 2C_2$ ; thus

$$Z_{i,j}' = \frac{1 + 2(1-r)C_2}{1 + 2C_2} C_1 + \frac{2rC_2}{1 + 2C_2} \frac{(a + b)}{2} + \frac{r}{\sqrt{1 + 2C_2}} C_3$$

And we can see that the difference will be the third term  $\frac{2r + 2rC_2 - r^2C_2}{\sqrt{(1 + C_2)^3}} C_3$  and  $\frac{r}{\sqrt{1 + 2C_2}} C_3$ .

$$\text{Method 1: } \frac{2r + 2rC_2 - r^2C_2}{\sqrt{(1 + C_2)^3}} C_3 \approx \frac{2r + 2rC_2 - r^2C_2}{1 + C_2} C_3$$

$$\text{Method 2: } \frac{r}{\sqrt{1 + 2C_2}} C_3 \approx \frac{r}{1 + C_2} C_3$$

Method 1-Method 2, thus:

$$\begin{aligned} \frac{2r + 2rC_2 - r^2C_2}{1 + C_2} C_3 - \frac{r}{1 + C_2} C_3 &= \frac{r + 2rC_2 - r^2C_2}{1 + C_2} C_3 \\ &= \frac{-C_2(r - 1)^2 + r + C_2}{1 + C_2} C_3 > 0 \text{ and } < C_3 \end{aligned}$$

10 Here  $C_2$  and  $C_3$  are also very small, thus the results of these two methods are almost same. And we can extend that to longer OTWs and more observations cases in this study.


And we can see that for a CDA system in the CGCM with a suitable inflation scheme, the assimilation approach with the time-averaged observations will have more advantages over that which sequentially assimilate the observations. It is very interesting to comparing these two approaches, but using either  
15 assimilation approach will have no influence on the essential of this study.

3 From your result I can not connect the de-correlation time with the optimal OTW directly. The  
optimal OTW is much short than the de-correlation time. Like in figure7, the correlation is 0.995 for the  
optimal OTW. Please revise the abstract and conclusion. The time scales of the variables really can not  
5 provide the useful information to quantify the optimal OTW. I suggest you remove or shorten the  
discussion and analysis related to the time scales of the variables.

RE: As Line 19-27 of Page 10, We have shortened this part. On the one hand, we have deleted the  
discussion about the relationship between the length of the optimal OTWs and the de-correlation time  
scales. On the other hand, we reserved the discussion about the high correlation between the  
10 observations including in the optimal OTW and the model state at the analysis time, which makes sense  
to the main idea of this study. Thanks.

4 The time (s) of observations in the OTW are different from the assimilation time. The temporal offsets  
introduce the represent errors for the observations, which need to be consider during analysis. If not,  
15 EnKF overweights the observations The analysis will not be optimal and the analysis ensemble spread  
will has negative biased. Please address this part and justify your assimilation.

RE: This is a very important and necessary comment. As Line 19-33 of Page 12, we have added some  
discussion about the temporal offset introduced the representation errors for the observations that are  
different from the analysis time. We use the de-correlation coefficients (As the Panel a and b of the  
20 following figure) to weight the observations including in the OTWs and avoid the overweighting. The  
comparison of the results of the weighted and no-weighted experiment are as following:



We can see that the lengths of the optimal OTWs obtained by these two assimilation schemes are the same except that the RMSEs of the weighted observation experiment will be lower than that of the non-weighted observation experiment when using the longer OTWs (when the length of ATM-OTW is greater than 4 and that of the OCN-OTW is larger than 50). This is owing to the high correlation between the observation including in the optimal OTWs and models states at the analysis time (exceed 0.995). Thus the influence of the temporal offset can be ignored and the results obtained by these two scheme will be almost same when using the lower OTWs (no or less greater than the lengths of the optimal OTWs). When we use the longer ones, the correlation will decrease and influence of the temporal offsets will be obvious that the results of the weighted observation experiment will better.

For the CGCMs, owing to the complex physics and dynamics, the influence of the time offsets will be obvious and the weights of the observations will be very necessary. But from this simple model case, we can see that whether or not using the weighted observations, the relationship between the characteristic variability time scales and the optimal OTWs will be firm and the essence of this study is established. Thanks.

5 It is hard to find the assimilation intervals for different components. Please address them together in experiment setup (2.3). I noticed that the OTW is longer than the assimilation interval, which means that you assimilate the same observations multiple times. In the perfect framework, it should degrade the filter performance from my understand and the information theory. Can you explain why you can  
5 achieve a good analysis under that circumstance. For a biased model framework, people do use observation multiple times to compensate the negative bias of forecast ensemble spread. So it make sense to see the optimal OTW in biased model cases are longer than that in perfect model case.

RE: As Line 21-22 of Page 7, in this study the assimilation interval of the atmospheric and slab oceanic states are 5 and 20 time steps, respectively. But the optimal OTWs obtained for the atmospheric and  
10 oceanic component in the perfect experiment framework are 1 and 10, which shows that no observation will be assimilated for multiple times. When the length of the OTW increases beyond that of the optimal one, the filter performance degrades, which satisfy your understanding and the information theory. Thanks.

15 6 The section 2.2 of “Ensemble coupled data assimilation” is incomplete. People will not understand your equations without trace back to your references. Please provide the completed two steps of EAKF.

RE: This comment is necessary. As in the section 2.2 in the revised manuscript we have provided the completely two steps of EAKF. Thanks.

20 7 Page 9 the last paragraph. The discussion of OTW with 4D-var is very confusion and hand-waving. The OTW in 4D-var is very different concept than yours. You should remove i

RE: Have removed it. Thanks.

8 Figure 4(a & b) show that the lower bound of RMSE reaches to 0. That does not make sense. Please  
25 check.

RE: Figures 4 and 7 are re-plotted. Thanks.

9 Figure 4 and Figure 5 show that the ensemble spread is significantly smaller than the RMSE. Please address the reasons and the effect on the assimilation.

5 RE: As Lines 14-18 of Page 10 of the revised manuscript, in this study, no inflation scheme was used in all the assimilation experiments. And only the data obtained in the last 5000 TUs will be used to conduct the error statistics for evaluation. Thus after first 5000 TUs assimilation, the ensemble spread has been greatly reduced and is significantly smaller than the corresponding RMSE. Although in theory the reduced ensemble spread will degrade the quality of state estimation, in this study with a one-dimensional conceptual coupled model, for simplicity and computational convenience as well as  
10 convenience for comparison, we insist not using the inflation scheme. Thanks.

10 P11 Lin13 I can not buy the statement of “since more observational information is needed to compensate the model bias and recover the characteristic variability.” The reason should be “the forecast ensemble in a biased model underestimates the forecast error, which results the EnKF  
15 underweight the observations. Therefore one can improve filter performance by using observation multiple times”

RE: We fully agree with your point. In the biased model experiment framework, the length of the optimal OTWs is larger than those in the perfect experiment case. The only reason will be the influence of the bias in the biased model. As Line 21-27 of Page 11, we think that we need stronger observational  
20 constraints to reduce the mean discrepancy between the model and observation induced by the influence of the bias. Also we fully agree that it needs to assimilate the observations for multiple times to compensate the underestimation of the forecast error by the bias in the biased model. We think these two interpretations are identical that using larger OTWs to compensate the influence of the bias except from two different aspects. Thus we think these two interpretations are not contradictory. Here we  
25 reserve our original interpretation and add your point into the revised manuscript. Thanks.

11 A reading proof is required to improve the manuscript. There are too many typos and grammar errors. For example Page 2 Lin 33 The “ensemble filter” should be “ensemble Kalman filter” Page 6 Lin 13 “but they are started from different initial states.” can be deleted Page 6 Lin 15 “observation” should be

“truth” Page 6 Lin 33 “the coupling co- efficient C1 is sensitive to model stability” should be “the model stability is sensitive to the coupling coefficient C1” Page 7 Lin3-5 The whole sentence need rephrase. You can just mention that “Following Zhang and Anderson (2003) an ensemble size of 20 is applied in all experiments in this study.” Page 7 Lin9-11 “absolute error” is not understandable. I guess that is “the absolute values of error”. The experiments are not described clearly. Please rewrite. Page 7 Lin28 the sentence is not completed. Page 7 Lin 30 Please also give the equivalent time of atmosphere (1-2 model month?). The deep ocean should be 50 model year.

RE: Thank you very much for your indication. All of these are fixed.

Here the “absolute error” just represents the “absolute value of the difference between the estimated model state value and corresponding truth”.

Second in the section 2.3 of the revision manuscript, the experiments has been rewritten and described clearly.

As Line 28-29 of Page 8, the equivalent time of atmosphere is 1-2 model month and that of the deep ocean should be 5 model decades.

Thank you very much again for your generous help and comments!  
We hope that the revised manuscript meets your requirement.

Sincerely yours,  
Shaoqing Zhang and Co-author



Responds to O. Martinez Alvarado:  
Dear O. Martinez Alvarado,

5 A few tel-conferences of all co-authors have been held to discuss yours comments. All authors converge to the point that all the comments are very important and useful for authors to improve the quality of this manuscript (MS). Therefore, all comments of yours have been fully addressed in the revision.

Now we will reply to each comment point by point as following:

10 1 It would be useful to have more background information on the CDA strategy studied in the paper. In particular I was wondering why the observational data should be assumed at the assimilation time when 4DVar would avoid that problem. It was only in Section 4 that 4DVar was mentioned as an alternative.

RE: As Line 10-16 of Page 2, we have added some background information on the CDA strategy in the introduction part of the revised manuscript. A 4-dimensional variation (4D-Var) scheme implements state estimation through minimizing a distance measure between the model and observations defined on all individual observational times within an OTW. Thus the minimized distance reflects the averaged effect in the OTW. As in Hunt et al. (2004), we expand the EnKF to include a time window in which the observations are treated as the exact assimilation times, even though their times are different in the window. Namely, we just assume that all the collected data sample the “truth” variation at the assimilation time and will be sequentially assimilated with their original error scales. Thus in this study the observational data included in the OTWs should be assumed at the assimilation time while 4DVar can relax that problem. Thanks.

15  
20

2 P2, L20-22: The two questions posed are circular in the sense that the answer to the first would depend on the effects of varying the OTW on the quality of CDA and the answer to the second depends on the existence of an optimal OTW. I would suggest the following rearrangement: ‘1) What is the impact of varying OTWs for each coupled component within the coupled model framework on the quality of CDF for climate estimation and prediction initialization? 2) Based on this impact, Is there an

25

optimal OTW so that assimilation fitting has maximum observational information, but minimum variability distortion?’ The point is subtle, but it might be worth doing for clarity.

RE: Thank you very much for your generous comment and we fully agree with that. We have corrected that as Line 27-29 of Page 2 in the revised manuscript.

5

3 Equation (1): Should there be a dot over eta? Is the dot over C6 correct?

RE: Fixed. Thanks.

4 It is not necessary to describe the Runge-Kutta method in Section 2.1.

10 RE: As Line 3-4 of Page 5, we have deleted the discussion and equation of the Runge-Kutta method. Thanks.

5 P5, L3: What is ‘localization and imbalance’?

15 RE: Because of the sampling error from a finite ensemble size, the ensemble-evaluated background variance is usually underestimated, and spurious correlations exist between a state variable and remote observations. To remove the long-distance spurious correlations and increase the reliability of ensemble-evaluated background co-variance, the localization technique was introduced into ensemble-based filters.

20 Given a biased CGCMs and the atmospheric and oceanic observing system (also including the land and sea ice), maintaining a balanced and coherent climate estimation is of critical importance for producing accurate climate analysis and prediction initialization. Given the assimilation model bias of warmer atmosphere and colder ocean, the atmospheric-only (oceanic only) data constraint produces an overcooling (overwarming) ocean through the atmosphere–ocean interaction will create imbalanced and incoherent oceanic (atmospheric) states estimation contrast to the observational model. The imbalanced  
25 atmosphere-to-ocean flux tends to decrease the assimilation quality and produce different climate features and variability from the real world.

But this one-diamond conceptual coupled model is too simple to simulate the complex physics such as the imbalance between different components and have no need to apply the localization scheme and consider the imbalance.

5 6 P5, L26: Please define  $Y_k$ , or is it the same as  $Y_{\{k,t\}}$ .

RE: In this study the  $Y_k$  is same as  $Y_{k,t}$ , as Line 24 of Page 5. Thanks.

7 P7, L3-5: Move this paragraph to the end of section 2.2. I was actually wondering whether twenty cases were sufficient to compute statistics.

10 RE: This simple conceptual coupled model is same as that used in Zhang et al. 2012 (A study of  
enhance parameter correction with coupled data assimilation for climate estimation and prediction  
using a simple coupled model). In that study, based on the trade-off between cost and assimilation  
quality, after a series of sensitivity tests on ensemble sizes of 10, 20, 50 and 100, no significant  
difference on assimilation quality is found when the ensemble size is greater than 20. Thus we chose the  
15 ensemble size of 20 in this study. And in order to avoid causing confusion, we correct that as Line 28-29  
of Page 7 of the revised manuscript. Thanks.

8 P7, L30: How are model years defined?

RE: A model year is defined as 10 non-dimensional time units, as Line 24 of Page 6. Thanks.

20

9 P8, L9-11: What do you mean by the strong nonlinearity and smoothness of the Runge-Kutta method?  
Can you elaborate on this point? Perhaps a different method would be more appropriate to obtain the  
solution of system (1).

RE: Usually the Leap-frog method and Runge-Kutta method will be used as the time difference time  
25 scheme to obtain the solution of the climate models. In the paper named “Mitigation of coupled model  
biases induced by dynamical core misfitting through parameter optimization: simulation with a simple  
pycnocline prediction model”, Fig.3 showed the power spectra of this simple coupled model’s states

derived by the Leap-frog and RK4 time difference scheme. We can see that the characteristic variability time scale of each coupled components obtained by both methods is a little larger than that set in Eq. (3) and varies for different time difference time scheme selected. The main idea of this study just illustrates that a closely relationship between the length of optimal OTWs and the corresponding characteristic variability time scales exists, which is independent from the time difference scheme we selected in this study. Owing to the simplicity and directness, we choose the RK4 as the time difference scheme used in this study. But its strongly nonlinearity and smoothness of the Runge-Kutta method will smooth the solution of the time series of the solutions, the characteristic variability time scale will be extended and larger than that set originally. Thanks.

10

10 P8, L22: It is not clear why the spread should be small. Please explain this point.

RE: In this study each assimilation experiment will be repeated for 20 times starting from 20 different ensemble initial conditions. The mean value and uncertainty (standard deviation) of the 20 cases will be evaluated. Here the spread just represents the standard deviation of the 20 cases. Thus the spread will be smallest when using the optimal OTW, as Line 17-19 of Page 9. Thanks.

15

11 P8, L28: What is ‘convexity with respect to OCN-OTWs’?

RE: Sorry for this mistake and confusion. And the “convexity” has been replaced by “sensitive variation” which just represents the variation of the  $\omega$ -RMSE curves is sensitive with respect to the OTWs. As Line 25 in Page 9 of the revised manuscript we have corrected. Thanks.

20

12 P11, L1: Which one is the curve of X2-RMSE in the OCN-OTW space? The figure (Fig. 9b) only shows one set of lines as a function of OCN-OTW and it corresponds to omega-RMSE.

RE: In this study the curve of X2-RMSE in the OCN-OTW space is not shown and the reason has been added as Line 25- 27 in Page 9 in the revised manuscript. Thanks.

25

13 P11, L12-17: The information in these lines is a repetition from the previous paragraphs. They can be deleted.

RE: Corrected. Thanks.

5 14 P12, L3-8: The discussion about the influence of the coupling term in the optimal OTWs is interesting, but it is hard to see what the implications for the real world or for a realistic CDA system are. Please, also discuss these points.

RE: As we have shown that changing that changing coupling strength (controlled by the coupling coefficients  $\mathcal{C}_1$  and  $\mathcal{C}_2$  in this case) between the atmosphere and upper ocean may have some influence on the characteristic variability time scales of coupled media, so as on the optimal OTWs. On the one hand these experiments can further illustrate the idea that a close relationship between the length of the optimal OTW and the corresponding characteristic variability time scale exists. On the other hand, for a realistic CDA system, the coupling physics could be very complicate and affected by many factors. The results of this simple model give the insights that we can only consider the factors which have obvious influence on the characteristic variability time scales when determining the length of the optimal OTWs for a realistic CDA system. And the factors which will have litter influence on the characteristic variability time scales can be just ignored. Through this way the process of determining the optimal OTWs for the realistic CDA system will be greatly simplified and make it possible to apply the method of using the optimal OTWs to the realistic CDA system. As Line 22-28 of Page 13 and Line 20-23 of Page 14, we have added some discussion on these points. Thanks.

15 Figure 5: What does the blue shading under the lower bound mean?

RE: As Line 8-9 of Page 10 and Line 3-4 of Page 20 of the revised manuscript, the blue shading below the ensemble spread represents the range of the uncertainty of state estimation in each assimilation experiments. Thanks.

16 Figure 10c is not discussed and therefore could be dismissed.

RE: From panel c) we can clearly see that the time series of the  $\omega$  also obviously varies with changing  $\mathcal{C}_2$ , which also corresponds to the results shown in panel b) in Figure 10. Thus we think that panel c) is also necessary and reserve it. Thanks.

- 5 17 Figures 11a and specially Fig. 11b, which only shows a horizontal line, are not needed as they can be simply described in the text.

RE: We fully agree with this comment and in the revised manuscript the panel a) and b) of the Figure 11 have been deleted. Thanks.

- 10 Technical corrections:

P1, L21-23: Some of the main results are given in these lines. However, the sentence is written in a very confusing style. I would encourage the authors to rewrite it to make the abstract, and the paper, more accessible.

RE: This sentence has been rewritten, as Line 21-22 of Page 1. Thanks.

- 15 P1, L30: Delete 'the' between 'by' and 'coupled'.

P1, L32: Add 'to sub-grid processes' after 'approximation'.

P2, L11: Delete 'in each medium here' and add 'for each medium' after '(OTW)'.

P2, L20: Delete 'exist' between 'not' and 'an' and add 'exists' after 'OTW'.

P2, L21: Expand 'What's' to 'What is'.

- 20 P3, L25: Change 'frequent' to 'frequency'.

P4, L7: Delete 'Namely'.

P5, L6: Either change 'In the words of Zhang (2011a)' to 'Following Zhang (2011a)' or clearly indicate the quote from that paper using quotation marks.

P6, L15: Change 'including' to 'included'.

P6, L19 and L20: Change 'integrates' to 'is integrated'

P6, L24: Delete 'etc.' or perhaps change it to 'and other parameters'.

P7, L4: Change 'will obtain' to 'was obtained'.

5 P7,L11: Omega and w are two different letters.

Figure 3 presents the power spectrum of what?

P8, L23: Change 'keeps' to 'is kept'.

P9, L26 and L27: Change 'including' to 'included'.

P10, L15: Delete 'gained'.

10 P10,L11: I think where it says 'only limits' should say 'is limited'.

Caption to Fig. 9: Change 'panels' to 'panel'.

RE: Corrected. Thanks.

P2, L24: Change 'identify' to 'identifies'. Also, the end of the sentence, after 'medium'

is not clear. Please, rewrite.

15 RE: Corrected. (The sentence has been corrected as Line 30-31 of Page 2 in the revised manuscript).  
Thanks

P4, L8: Define TU.

RE: As Line 9-11 Of Page 4, TU is the time unit. And in this study 1TU equals 100 time steps. Thanks.

P4, L12: I think the sentence is incomplete. What does 'C2' does in contrast?

RE: In this study  $\mathcal{C}_1$  denotes the upper slab oceanic forcing on the atmosphere while  $\mathcal{C}_2$  in contrast. Here ‘ $\mathcal{C}_2$  in contrast’ just represents that  $\mathcal{C}_2$  denotes the atmospheric forcing on the upper slab oceanic, as Line 19 of Page 4. Thanks

P4, L16-17: As it is written, the sentence starting with ‘Where’ and ending with ‘scale’ makes no sense.  
5 Please, rewrite.

RE: This sentence has been rewritten as Line 24-26 of Page 4 in the revised manuscript. Thanks.

P4, L21: Do you mean ‘In summary’ rather than ‘Summarily’?

RE: Yes. Corrected. Thanks.

P4, L24: The method’s name is Runge-Kutta not Runger-Kutta. Since it is not used in the text the  
10 acronym RK4 is not necessary.

RE: In the revised manuscript we have corrected as ‘Runge-Kutta’ and deleted the acronym RK4.  
Thanks.

P4, L25: Change notation as ‘k0-k3’ looks like ‘k0 minus k3’.

RE: According to a reviewer’s generous comment, we have deleted the discussion about the RK4.  
15 Thanks.

P5, L14: The sentence does not make sense. Please rewrite.

RE: This sentence has been rewritten as Line 18-22 of Page 5 of the revised manuscript. Thanks.

P7, L28: The sentence is incomplete. Figure 3 presents the power spectrum of what?

RE: Corrected as Line 26 of Page 8 of the revised manuscript. Thanks.

20 Figure 10: The value of  $C_1$  in the legend is constant and therefore it is not needed there.

RE: We have deleted the  $C_1$  in the legend in Figure 10. Thanks.

Thank you very much again for your generous help and comments!



We hope that the revised manuscript meets your requirement.

Sincerely yours,

5 Shaoqing Zhang and Co-author

# Impact of Observational Time Window on Coupled Data Assimilation: Simulation with a Simple Climate Model

Yuxin Zhao<sup>1,2</sup>, Xiong Deng<sup>1,2</sup>, Shaoqing Zhang<sup>\*3,4</sup>, Zhengyu Liu<sup>5,6</sup>, Chang Liu<sup>1,2</sup>, Guijun Han<sup>7</sup>, Xinrong Wu<sup>7</sup>

<sup>1</sup>College of Automation, Harbin Engineering University, Harbin, 150001, China

5 <sup>2</sup>GFDL-Wisconsin Joint Visiting Program, Princeton, NJ08540, USA

<sup>3</sup>Key Laboratory of Physical Oceanography, MOE.China, Ocean University of China, Qingdao, 266003, China

<sup>4</sup>National Laboratory for Marine Science and Technology, Qingdao, China

<sup>5</sup>Laboratory for Climate and Ocean-Atmosphere Studies (LaCOAS), Department of Atmospheric and Oceanic Sciences, School of Physics, Peking University, Beijing, 100871, China

10 <sup>6</sup>Center for Climate Research and Department of Atmospheric and Oceanic Sciences, University of Wisconsin-Madison, Madison, WI 53706, USA

<sup>7</sup>National Marine Data and Information Service, Tianjin, 300171, China

*\*Correspondence to: Shaoqing Zhang ([szhang@ouc.edu.cn](mailto:szhang@ouc.edu.cn))*

**Abstract.** Climate signals are the results of interactions of multiple time scale media such as the atmosphere and ocean in the coupled earth system. Coupled data assimilation (CDA) pursues balanced and coherent climate analysis and prediction initialization by incorporating observations from multiple media into a coupled model. In practice, an observational time window (OTW) is usually used to collect measured data for an assimilation cycle to increase observational samples **that are sequentially assimilated with their original error scales**. Given different time scales of characteristic variability in different media, what are the optimal OTWs for the coupled media so that climate signals can be most accurately recovered by CDA? With a simple coupled model that simulates typical scale interactions in the climate system and “twin” CDA experiments, we address this issue here. **Results show that in each coupled medium, an optimal OTW can provide maximal observational information that best fits characteristic variability of the medium during the data blending process**. Maintaining correct scale interactions, the resulted CDA improves the analysis of climate signals greatly. This simple model results provide a guideline when the real observations are assimilated into a coupled general circulation model for improving climate analysis and prediction initialization by accurately recovering important characteristic variability such as sub-diurnal in the atmosphere and diurnal in the ocean.

## 1 Introduction

Currently, the interactions between the earth climate system’s major components, such as the atmosphere, ocean, land and sea ice, have been reasonably simulated by coupled climate models, which can also give the evaluation of climate changes (Randall et al. 2007). However, because of the uncertainties and errors in models (e.g., parameterization is only an approximation **to sub-grid processes** and dynamical core is imperfect), models always tend to produce different climate features and variability from the real world (e.g. Delworth et al. 2006; Collins et al. 2006; Zhang et al. 2014). Due to the significant importance of preserving the balance and coherence of different model components (or media) during the coupled

model initialization, data assimilation for state estimation and prediction initialization should be performed within a coupled climate model framework (e.g. Chen et al. 1995; Zhang et al. 2007; Chen 2010; Han et al. 2013). The characteristic variability time scales of different media within the coupled frameworks are usually different. When the observed data included in one or more components of the coupled system framework are assimilated, the observational information will be able to be transferred among different media through the coupled dynamics so that all media gain consistent and coherent adjustments. Such an assimilation procedure is called coupled data assimilation (CDA), which can sustain the nature of multiple time-scale interactions during climate estimation and prediction initialization (e.g. Zhang et al. 2007; Sugiura et al. 2008; Singleton 2011), thus producing better climate analysis and prediction initialization and therefore improve the coupled models' predictability (e.g. Yang et al. 2013). Zhang et al. (2007) developed the first CDA system in a fully coupled general circulation model, the version 2 of Geophysical Fluid Dynamics Laboratory Coupled Model (GFDL CM2). The National Centres for Environmental Prediction (NCEP) also started using coupled models to generate first-guess forecasts for their Climate Forecast System Reanalysis (CFSR, Saha et al. 2010). Despite the enormous benefits and demand for CDA, it remains both theoretical and technical challenges to implement strong CDA in fully-coupled models, including the estimation of the coupled model error covariance matrix and the huge computational costs (Han et al. 2013; Lu et al. 2015; Liu et al. 2016).

During the coupled data assimilation process, usually an observational time window (OTW) is used to collect measured data in each medium for an assimilation cycle (e.g. Pires et al., 1996; Hunt et al., 2004; Houtekamer and Mitchell, 2005; Laroche et al., 2007) to increase observational samples. As in Hunt et al. (2004), we expand the EnKF to include a time window in which the observations are treated as the exact assimilation times, even though their times are different in the window. Namely, we just assume that all the collected data sample the "truth" variation at the assimilation time and will be sequentially assimilated with their original error scales. Thus the OTW is applied in a 3-dimensional data assimilation fashion rather than a 4-dimensional one. Apparently, while a large OTW provides more observational samples at the assimilation time, the assimilation process blends more data from different times and may distort variability being retrieved. Given the fact that climate signals are the results of interactions of multiple time scale media, correct variability retrieved for each medium so as correct scale interaction maintained in CDA is particularly important for climate analysis and prediction initialization. In this study we attempt to answer the following two questions: 1) What is the impact of varying OTWs for each coupled component within the coupled model framework on the quality of CDA? 2) Based on this impact, does an optimal OTW exist so that assimilation fitting has maximum observational information but minimum variability distortion? With a simple conceptual coupled climate model and a sequential implementation of the ensemble Kalman filter, this study first analyses the characteristic variability time scale of each coupled medium and identifies the corresponding optimal OTW. Then the impact of optimal OTW on the quality of CDA and its linkage with the corresponding time scale of characteristic variability are investigated. The simple coupled model consists of three typical components, including the synoptic atmosphere (Lorenz 1963) and the seasonal-interannual slab upper ocean (Zhang et al. 2012) coupling with the decadal deep ocean (Zhang 2011a,b). Although the simple conceptual coupled model does not share the similar complex physics with a

coupled general circulation model (CGCM), it does reasonably simulate the typical interactions between multiple time-scale components in the coupled climate system (see Zhang et al. 2013). The simple coupled model helps us understand the essence of the problem by revealing the relationship between the optimal OTWs and corresponding time scales of characteristic variability as well as their impact on CDA. The low-cost nature of the simple model also provides convenience for a large number of CDA experiments with different OTWs in optimal OTW detection. The ensemble Kalman filter (e.g. Evensen 1994; 2007; Whitaker and Hamill 2002; Anderson 2001; 2003) used in this study is the ensemble adjustment Kalman filter (EAKF, e.g. Anderson 2001; 2003; Zhang and Anderson 2003). Using the EAKF with the simple coupled model, we first establish a twin experiment framework. Within such a framework, the degree by which the state estimation based on a certain OTW recovers the truth is an assessment of the influence of the OTW on the quality of CDA. By such a way, the optimal OTW of each medium is detected and the impact of optimal OTWs on CDA is evaluated. We also discuss the influence of model bias on optimal OTW through biased twin experiment setting.

This paper is organized as follows. Section 2 briefly describes the simple conceptual coupled model, the ensemble adjustment Kalman filter, as well as the twin experiment framework including perfect and biased settings. With a simplest case, we first show the influence of OTWs on assimilation quality and its linkage with the time scale of characteristic variability in section 3. Then section 4 presents results on detection of the optimal OTWs for different media and the impact of optimal OTWs on CDA. The influence of realistic assimilation scenarios on optimal OTWs is discussed in section 5. Finally, summary and discussions are given in section 6.

## 2 Methodology

### 2.1 The model

Due to the complicate physical processes and huge computational cost involved, it is inconvenient to use a CGCM to investigate the impact of the different OTWs on the analysis of climate signals so as to detect each coupled medium's optimal OTW. Instead, here we employ a simple coupled "climate" model developed by Zhang (2011a). This simple model is based on the Lorenz's 3-variable chaotic model (Lorenz 1963) that couples with a slab upper ocean (Zhang et al. 2012) and a simple pycnocline predictive model (Gnanadesikan 1999). Although very simple with low computational cost, in terms of multi-scale interaction inducing low-frequency climate signals, this model shares fundamental character with a CGCM and it is very suitable for addressing the problem that is concerned here. And for the readers' convenience, here we simply review some key aspects of this conceptual coupled model. With all quantities being given in non-dimensional units, the governing equations are:

$$\begin{aligned}
\dot{X}_1 &= -\sigma X_1 + \sigma X_2 \\
\dot{X}_2 &= -X_1 X_3 + (1 + C_1 \omega) k X_1 - X_2 \\
\dot{X}_3 &= X_1 X_2 - b X_3 \\
O_m \dot{\omega} &= C_2 X_2 + C_3 \eta + C_4 \omega \eta - O_d \omega + S_m + S_s \cos(2\pi t / S_{pd}) \\
\Gamma \dot{\eta} &= C_5 \omega + C_6 \omega \eta - O_d \eta
\end{aligned} \tag{1}$$

where  $X_1, X_2$  and  $X_3$  represent the atmospheric model states while  $\omega$  and  $\eta$  denote those for upper and deep ocean, respectively. A dot above the variable denotes the time tendency. The atmosphere model states are the high frequency variables while the slab oceanic variable is of a lower frequency. For sustaining the chaotic nature of the atmosphere in reality, the standard values of the parameters including in the atmospheric component ( $\sigma, k$  and  $b$ ) are set as 9.95, 28 and 8/3, respectively. In the equation of  $\omega$ , the parameters  $O_d$  and  $O_m$  denote the damping coefficient and heat capacity of the upper slab ocean, respectively. Due to the lower frequency of  $\omega$  than that of the model states in the atmospheric components, the time scale of the upper slab ocean variable must be much slower than that of the atmospheric model states. Thus the damping rate parameter ( $O_d$ ) should be much smaller than the heat capacity, namely,  $O_d \ll O_m$ . Here following Lorenz's idea (Lorenz 1963), the atmospheric time scale is defined as the typical time by which the atmosphere goes through an attractive lobe as one non-dimensional time unit (TU)  $\sim O(1)$ . We set the parameters ( $O_m, O_d$ ) as (10, 1), which show that the slab oceanic variable's time scale is  $\sim O(10)$ , i.e. 10 times of that of the atmospheric model states. While the  $S_m + S_s \cos(2\pi t / S_{pd})$  represents the external forcing, the parameter  $S_{pd}$  denoting the model seasonal cycle is set as 10 to make sure that the period of the external forcing is comparable with the upper slab ocean variables' time scale. In this simple coupled model, the seasonal cycle is set as 10TUs and thus a model year (decade) equals to 10 (100) TUs. And the parameter  $S_s$  and  $S_m$ , denoting the magnitudes of the external forcing's seasonal cycle and annual mean, are insensitive to the coupled model and set as (1, 10). The coefficients  $C_1$  and  $C_2$  in the equations of  $X_2$  and  $\omega$  are used to implement the coupling between the fast atmosphere model states and the upper slab oceanic variable and set as (0.1, 1), with that  $C_1$  denotes the upper slab oceanic forcing on the atmosphere while  $C_2$  denotes the atmosphere forcing on the ocean. In addition,  $C_3$  and  $C_4$  represent the deep oceanic forcing and the nonlinear interaction between the upper and deep ocean. In order to make sure that the atmospheric forcing plays a dominant role in the upper slab ocean, the magnitudes of  $C_3$  and  $C_4$  should be lower than that of  $C_2$  and both set as 0.01. Same as in Zhang (2011a), the deep ocean model state variable  $\eta$ , denoting the anomaly of pycnocline depth in the deep ocean, is derived from the two-term balance model of the zonal-time mean pycnocline (Gnanadesikan, 1999). Within the equation of  $\eta$ , the parameter  $\Gamma$  keeps as a constant and the ratio of  $\Gamma$  and  $O_d$  denotes the deep ocean variable's time scale. The time scale of deep ocean variable is longer than that of slab ocean, defined by the relative magnitude of  $\Gamma$  to  $O_d$  ( $\Gamma$  is set as 100). Similar to the equation of  $\omega$ , the coefficients  $C_5$  and  $C_6$  denote the linear slab oceanic forcing and the nonlinear interaction between upper and deep ocean. Also for guaranteeing that the linear interaction is dominant and the nonlinear interaction is weaker than that in the deep ocean model,  $C_5$  and  $C_6$  are set as (1, 0.001). In summary, in this study the standard values of the parameters including in this simple model

$(\sigma, k, b, c_1, c_2, O_d, O_m, S_m, S_s, S_{pd}, \Gamma, C_3, C_4, C_5, C_6)$  are set as (9.95,28,8/3,0.1,1,1,10,10,1,10,100,0.01,0.01,1,0.001) (e.g., Zhang 2011a,b; Zhang et al., 2012; Han et al., 2013,2014).

As the study of Han et al. (2014), the fourth-order Runge-Kutta time-differencing scheme is used in this paper to resolve this simple coupled model and the time step equals to 0.01TU (1 TU=100 time steps).

- 5 Zhang (2011b) illustrated that, given the model parameters described above, the constructed simple coupled model can effectively simulate a fundamental feature of the real world climate system in which different time scales interact with each other to develop climate signals. Namely, the synoptic to decadal time-scale signals are produced by the interactions between the transient atmosphere attractor, the slow slab ocean and the even-slower deep ocean (see Zhang 2011a; Han et al. 2014). Again, although the simple coupled model does not have complex physics and cannot consider the issue of impact of  
 10 localization and imbalance as in a CGCM, it can help us investigate the fundamental issue we want to address here more directly and clearly.

## 2.2 Ensemble coupled data assimilation

- Following Zhang (2011a), during the state estimation, the error statistics evaluated from ensemble model integrations, such as the error covariance between model states, will be used in an ensemble filter to extract observational information to adjust  
 15 the model states (e.g., Evensen, 1994, 2007; Anderson, 2001; Hamill et al., 2001; Zhang, 2011a,b; Zhang et al., 2012; Han et al., 2014). In this study, a derivative of Kalman filter (Kalman 1960; Kalman and Bucy 1961) called ensemble adjustment Kalman Filter (EAKF, Anderson 2001; 2003; Zhang and Anderson 2003; Zhang et al. 2007) which is a sequential implementation of ensemble Kalman Filter under an “adjustment” idea is used to implement the CDA scheme. The  
 20 assumption of independence of observational error allows the EAKF to sequentially assimilate observations into corresponding model states (Zhang and Anderson 2003; Zhang et al. 2007). While the sequential implementation provides much computation convenience for data assimilation, the EAKF maintains much of the non-linearity of background flows as possible (Anderson 2001; 2003; Zhang and Anderson 2003).

- Based on the two-step implementation of EAKF scheme (Anderson, 2001; 2003), the observational increment at an observation location is first computed. The observation is denoted as  $\mathcal{Y}$  at time  $t$  (simply  $\mathcal{Y}$  instead of  $\mathcal{Y}_t$ ) which has the  
 25 observation value  $\mathcal{Y}^o$  and standard deviation  $\sigma_{\mathcal{Y}}^o$  (assumed to be Gaussian). Firstly, the reshaping of the model ensemble at the observation location,  $\Delta\mathcal{Y}'$  is formulated as:

$$\Delta\mathcal{Y}'_i = \frac{\Delta\mathcal{Y}_i^p}{\sqrt{1+r_k^2}} \text{ and } r_k = \frac{\sigma_{k,k}^p}{\sigma_{k,k}^o} \quad (2)$$

- where  $i$  represent the ensemble index and  $k$  denotes the observation index.  $\sigma_{k,k}^o$  and  $\sigma_{k,k}^p$  are the standard deviation of observation error and its prior estimated ensemble standard deviation, respectively, while  $r_k$  is the corresponding ratio. If  
 30  $r_k > 1$ , the ensemble spread is largely reduced by the observation; otherwise, the ensemble remains close to the prior. The shift of the ensemble mean induced by the observation is computed by:

$$\bar{y}^u = \frac{\bar{y}^p}{1 + r_k^2} + \frac{y^o}{1 + r_k^{-2}} \quad (3)$$

We can see that if the prior estimated ensemble standard deviation is greater than that of the observation error, the ensemble mean shifts toward the observation value; otherwise the ensemble mean remains close to the prior model ensemble mean  $\bar{y}^p$ . Then the observational increment induced by the observation value  $y^o$  for the  $i$ th ensemble member at the  $k$ th observation location is computed as:

$$\Delta y_{k,i}^o = (\bar{y}_k^u + \Delta y_{k,i}^u) - y_{k,i}^p = \left( \frac{\bar{y}_k^p}{1 + r_k^2} + \frac{y_k^o}{1 + r_k^{-2}} + \frac{\Delta y_{k,i}^p}{\sqrt{1 + r_k^2}} \right) - y_{k,i}^p \quad (4)$$

- 5 Once we get the observational increment at the observation location, then a least square fit is used to distribute the increment over the relevant grid points impacted by the observation using the covariance between the grid index  $j$  and the observation  $k$ ,  $c_{j,k}^p$ , using:

$$\Delta Z_{i,j} = \frac{c_{j,k}^p}{(\sigma_{k,k}^p)^2} \Delta y_{k,i}^o = \frac{Cov(Z_j, y_k)}{(\sigma_{k,k}^p)^2} \Delta y_{k,i}^o \quad (5)$$

- Where  $Z$  represents a certain state variable at the grid point  $j$ . The term  $\Delta Z_{i,j}$  is the contribution of the  $k$ th observation to  $i$ th ensemble member of the model state estimated at grid point  $j$ . When an observation is available, the Eq.(5) will be applied to  
 10 implement CDA for state estimation in a straight forward manner (Zhang et al., 2007; Zhang 2011a;).

- Although many sophisticated inflation algorithms (e.g. Anderson 2007;2009; Li et al. 2009; Miyoshi 2011) exist for atmosphere data assimilation, the inflation scheme for a coupled model is a new subject due to the multiple time-scale nature of the system. Furthermore, trial-and-error experiments show that the usual form of inflation (e.g. only inflate the atmosphere model states or inflate all the model states equally) will lead the analysis to become unstable. Thus, in this paper,  
 15 for simplicity and computational convenience as well as convenience for comparison, no inflation is used in our assimilation experiments, just as in Han et al. (2014).

### 2.3 Perfect and biased twin experiment setups

- In this study, a perfect twin experiment framework and a biased twin experiment framework are designed, respectively. In both perfect and biased twin experiments, a “truth” model using the standard parameter values listed in section 2.1 is used to  
 20 generate the “true” solution of the model states and produce the observations sampling the “truth”. Starting from the initial condition (0,1,0,0,0), the “truth” model is firstly integrated forward 10000TUs (i.e. 1000 model years) for sufficient spin-up and then integrated forward for another 10000TUs to generates the “truth” model states. The observations are produced by sampling the “truth” solution of the model states at an observational interval and superimposing with a white noise simulating the observational errors. As schematically shown in **Fig. 1**, all the observational intervals using in this study are  
 25 assumed to be 1 time step (0.01TU). Although in the real climate system, the oceanic observations are usually available less frequently than those in the atmosphere (namely the oceanic observation interval is larger than that we set here), for this

proof-of-concept study we will set the time interval of the oceanic observations as small as possible. The standard deviations of the observational errors are 2 for  $\mathcal{X}_1, \mathcal{X}_2, \mathcal{X}_3$  and 0.5 for  $\omega$ . Also, although the deep ocean lacks observations in the real world, we also conduct some observation simulation experiments for  $\eta$  (the standard deviations of the observational error is 0.06 for  $\eta$ ) in this conceptual study.

5 We first want to learn some basics from the perfect experiment which represents an idealized data assimilation regime. In the perfect twin experiment framework, the assimilation model also uses the standard parameter values, but starts from different initial conditions. Using the Gaussian white noises with the same standard deviation as observational errors (2 for  $\mathcal{X}_1, \mathcal{X}_2$  and  $\mathcal{X}_3$ , 0.5 for  $\omega$  and 0.06 for  $\eta$ ) added on the model states at the different time in spin-up run to form the ensemble initial conditions for each ensemble filtering data assimilation experiment. Each assimilation experiment is integrated for  
10 10000TUs and only the data obtained in the last 5000TUs is used to conduct error statistics for evaluation. We choose the model states between 9000TUs and 10000TUs during the spin-up at an interval of 50TUs being perturbed to form 20 cases of ensemble initial conditions for each assimilation experiment analysed in section 4 and 5. In this way, we attempt to minimize the dependence of the results of optimal OTWs on ensemble initial states. Then each assimilation experiment will be repeated for 20 times starting from these 20 independent ensemble initial conditions and we will analyse the mean value  
15 and uncertainty evaluated from these 20 cases.

Then we use the biased experiment setting to simulate the real world scenario. The biased twin experiment framework is similar as the perfect one except that the assimilation model in the biased twin experiment framework has a systematic discrepancy from the observations. Thus, in the biased twin experiment framework, the parameters included in the assimilation model will have 10% errors relative to the standard values. The errors in the parameters will be the only model  
20 error source.

**Figure. 1** also illustrates the assimilation update intervals (the assimilation intervals are 5 time steps for atmosphere, 20 time steps for slab ocean in all assimilation experiments and 100 time steps for deep ocean if using the  $\eta$  observations) as well as the length of observational time window (OTW), which will be used throughout the study. In addition, the coupling strength between the atmosphere and ocean may have influences on the characteristic variability time scale of each coupled medium,  
25 so as on the optimal OTW. We discuss this issue through changing the values of coupling coefficients  $\mathcal{C}_1$  and  $\mathcal{C}_2$ . In this simple model case, the model stability is sensitive to coupling coefficient  $\mathcal{C}_1$  (Zhang et al., 2012) and changing  $\mathcal{C}_1$  only influences on the chaotic component, so here we just change  $\mathcal{C}_2$  to investigate the impact of coupling coefficient between the atmosphere and upper ocean on the optimal OTWs. As in Zhang and Anderson (2003), an ensemble size of 20 is applied in all assimilation experiments in this study.



### 3 The length of observational time window (OTW) and retrieval of characteristic variability

#### 3.1 Influence of OTW on the accuracy of CDA

In order to exhibit the influence of OTW on the quality of climate analysis, we show 3 simple assimilation experiments (the time series of  $\omega$ 's absolute errors) in **Fig. 2**: 1) standard CDA (green) (assimilating the observations rights at the analysis times without any atmospheric/oceanic OTW); 2) OCN-OTW(5) CDA (red) (assimilating all 11 ocean observations in an oceanic OTW with a half-width of 5, defined as the length of the OTW hereafter, but no atmospheric OTW is considered) and 3) OCN-OTW(100) CDA (assimilating all 201 ocean observations in an oceanic OTW with the length of 100, but no atmospheric OTW is considered). Three assimilation experiments above are all conducted using the perfect model setting (all the parameters use their standard values) and the uni-variate adjustment scheme. And the atmospheric and oceanic update intervals are 0.05 TU and 0.2 TU, respectively. While the standard CDA does NOT use the atmospheric and oceanic OTWs and only assimilates the observations right at the analysis time, the OCN-OTW CDA incorporates all the valid observations collected in the oceanic OTW. All three assimilation experiments above do not use an atmospheric OTW.

From **Fig. 2**, we can see that a small OCN-OTW (total 11 observations in the oceanic OTW) can make a much better ocean analysis than the standard CDA (comparing the red line with the green line). We can understand that this is because an OTW can provide more observational information thus enhancing the observational constraint so as to improve the accuracy of climate analysis. However, comparing the blue line to the green/red line, it is clear that a too large OTW degrades the quality of the ocean analysis. The results of these simple assimilation experiments tell us, if an appropriate OTW is used, we can gain optimal climate analysis. How can we determine such an optimal OTW? Next, starting from analysing characteristic variability of each coupled medium, we will discuss the methodology how to determine an optimal OTW for each medium in a coupled climate system.

#### 3.2 The time scale of characteristic variability and optimal OTW

The key to improve the accuracy of climate analysis in CDA is accurately recovering characteristic variability of different media in the coupled system. Thus we can assume that the length of optimal OTW for each medium will have some relationship with the corresponding characteristic variability time scale. Then, we should first analyse the time scale of characteristic variability in each medium.

**Fig. 3** presents the power spectrum of  $\mathcal{X}_2$ ,  $\omega$  and  $\eta$  based on the model states with a 4800-TU length (totally 480000 data) after the spin-up described in section 2.3. From **Fig. 3**, we learned that in this simple model, the characteristic variability time scales of atmosphere ( $\mathcal{X}_2$ ), upper ocean ( $\omega$ ) and deep ocean ( $\eta$ ) are about 1-2 TUs (1-2 model months), 50-100 TUs (5-10 model years) and 500 TUs (5 model decades), respectively. Namely, the characteristic variability time scale of the slab ocean is much larger than that of the atmosphere but smaller than that of the deep ocean.

An optimal OTW aims to provide maximal observational information that best samples characteristic variability of that medium during the data blending process. Thus the length of the optimal OTW should be smaller than the corresponding

characteristic variability time scale, which means that the optimal OTW in the atmosphere must be much smaller than 1 TU (100 time steps), and in the ocean, the optimal OTW must be much smaller than 50 TUs (5000 time steps). If we will take observations for  $\eta$ , the optimal OTW for  $\eta$  must be much smaller than 500 TUs (50000 time steps). From **Fig. 3**, we also see that the characteristic variability time scales of different coupled media are a little larger than the corresponding ones set in the Eq.(1). This is owing to the strong nonlinearity and smoothness of the fourth-order Runge-Kutta time-differencing scheme that prolongs the characteristic variability time scales of the simple coupled model. But they do not change the essence of the problem we address in this study. Given different time scales of characteristic variability in different media, in the following section we will further detect the optimal OTWs based on the corresponding characteristic variability time scales and examine their impact on the quality of climate analysis in CDA.

#### 10 4 Detection of the optimal observational time window

In this section, with the perfect model framework described in section 2.3, we first conduct a series of CDA experiments with different ATM-OTWs and different OCN-OTWs to detect the optimal OTW for each medium. The assimilation scheme is the simple uni-variate adjustment scheme serving as a proof-of-concept study. To eliminate the dependency of results on initial states, each experiment is repeated for 20 independent initial conditions described in section 2.2. Then the mean value of 20 case RMSEs and the spread are plotted in **Fig. 4**.

**Fig. 4a** shows that the optimal ATM-OTW is 1, i.e., the optimal ATM-OTW includes only 3 atmosphere observations, with which the assimilation produces the lowest RMSE of the atmosphere and the smallest spread (**In this study each assimilation experiment will be repeated for 20 times starting from 20 different independent initial ensemble conditions. Here the spread just represents the standard deviation of these 20 cases. Thus it will be smallest when using the optimal OTW.**). In these experiments for detecting the optimal ATM-OTW, the ocean assimilation is kept in the standard setting (i.e. no OTW, 0.2 TU update interval). Then we keep the ATM-OTW as 1 and change the length of OCN-OTW to produce **Fig. 4b**.

From **Fig. 4b**, we can see that the optimal OCN-OTW is about 10 (i.e., each OTW includes total 21 observations), with which the lowest  $\omega$ -RMSE and the smallest RMSE spread are produced. Compared to the case of standard CDA (denoted as CDA\_NOTW), the uses of optimal ATM-OTW and OCN-OTW make the RMSEs of  $\mathcal{X}_{1,2,3}$  and  $\omega$  significantly reduced. When the RMSE of  $\omega$  has a distinguishable **sensitive variation** with respect to OCN-OTWs, the RMSE of  $\mathcal{X}$  (**Because we just choose the optimal OCN-OTW from the figure of  $\omega$ -RMSE. Thus in this study the variation of  $\mathcal{X}$ -RMSEs in the OCN-OTW space is not shown.**) does not show such a sensitivity to the optimal OCN-OTW. This means that in this simple system, due to the strong nonlinearity and chaotic nature of the “atmosphere”, the improved accuracy for  $\omega$  from optimal observational constraint is not sufficient to impact the “atmosphere” (this point will be expended in section 5.3). Similar to the characteristic variability time scale of the slab ocean vs. that of the “atmosphere” shown by **Fig. 3**, the optimal OCN-OTW is much larger than that of ATM-OTW.

To further understand the relationship between the optimal OTW and characteristic variability time scale, we also examine the  $\eta$  –RMSEs in the space of  $\eta$ -OTWs. The assimilation interval of the pycnocline depth is set as 1 TU (100 time steps), which is much larger than that of the slab ocean. When we change the  $\eta$ -OTW, the optimal ATM-OTW and OCN-OTW detected from **Figs. 4ab** are used. As shown in **Fig. 4c**, the optimal  $\eta$ -OTW is about 100 (i.e. total 201 observations), which is much larger than that of OCN-OTW and smaller than the characteristic variability time scale of the deep ocean pycnocline depth. With the optimal  $\eta$ -OTW, the RMSE of  $\eta$  is reduced about 77.4% from the level of the CDA\_NOTW.

We also check the variation of the 20-case mean ensemble spread in the space of OTWs as shown in **Fig. 5**. The mean and standard deviation of the ensemble spreads of  $\mathcal{X}_{1,2,3}$  and  $\omega$  (the uncertainty of the state estimation in each assimilation experiment is shown as the blue shadow in **Fig. 5**) gradually decrease when the ATM-OTW and OCN-OTW become larger.

When the OTWs are set too large (here the ATM-OTW and OCN-OTW are greater than 20 and 250, respectively), the ensemble spreads of  $\mathcal{X}_{1,2,3}$  and  $\omega$  decrease dramatically. While increasing the length of the OTWs, more observations will be assimilated into the corresponding model states, which can function as a smoother. The larger the OTW is, the stronger the smoother will be. Under this circumstance, the too strong smoother will distort characteristic variability of the model states, which explains the green line of **Fig. 2**. Also from **Fig. 4** and **Fig. 5**, we can see that the mean of ensemble spread is significantly smaller than that of the corresponding RMSE. It is owing to that no inflation scheme is applied in this study. And the statistics for evaluation are conducted from the data obtained in last 5000TUs. Thus after first 5000TUs' assimilation in each assimilation experiment, the ensemble spreads of model states have been greatly reduced due to no inflation. Then the mean ensemble spread is significantly smaller than the mean RMSE.

To understand the essence of optimal OTWs, we show the auto-correlation for each model state and marks the time correlation coefficients at the time scales of optimal OTWs for  $\mathcal{X}_2$  (panel a),  $\omega$  (panel b) and  $\eta$  (panel c) detected from **Fig. 4** in **Fig. 6**. The result is the mean of 20 cases. In each case, the number of data is 10000 (steps) (100TUs), which are chosen from the period of 5000TUs to 9000TUs in the truth run after spin-up. From **Fig. 6** we can see that all auto-correlation at the optimal OTW length are located around 0.995. This means that the observations including in an optimal OTW are extremely highly correlated with the model state at the analysis time. This can be understood since in this sequential assimilation scheme all the observations including in an OTW are assumed to be sampled at the analysis time so that the difference among them must be in a negligible arrange. Under such a circumstance, the optimal OTWs provide maximal observational information that best fits characteristic variability and minimizes the analysis error.

## 5 Influences of realistic assimilation scenarios on optimal OTW

In this section, we first show the impact of the multi-variate adjustment scheme on the optimal OTWs in perfect model setting. Then we discuss the influence of model bias through a biased model framework. We will also investigate the impact of coupling strength on the optimal OTWs.

## 5.1 Influence of multi-variate adjustment on optimal OTWs

While the experiments with the uni-variate adjustment scheme provide us a direct understanding of the influence of the OTWs on CDA, we want to check whether or not it also applies to the multi-variate adjustment scheme. So we repeat the experiments described in section 4 but with the multi-variate adjustment scheme. The results are shown in **Fig. 7**. Here the multi-variate adjustment scheme is only limited to the atmospheric observations (i.e., only the cross-covariances among  $\mathcal{X}_1, \mathcal{X}_2, \mathcal{X}_3$  are used) (as indicated in Han et al. 2013, the multi-variate adjustment scheme using the coupling cross-covariance between different coupled media involves complex scale interactions and may complicate the investigation of the problem we are addressing here). The results shown in **Fig. 7** are similar with **Fig. 4**, suggesting the multi-variate adjustment scheme has little influence on the optimal OTWs, **since it does not change the characteristic variability time scales (especially in this simple model)**.

The perfect experiment framework provides a direct guideline for the relationship between the optimal OTW and the corresponding characteristic variability time scale. However, in reality, the numerical model has errors and is biased with the observation. It is necessary to investigate the influence of model bias on optimal OTWs so as on the quality of CDA.

## 5.2 Influence of model bias on optimal OTWs

With the biased model experiment framework described in section 2.3, we repeat all the experiments above for detection of the optimal OTWs. The results are shown in **Fig. 8**. Compared to the results in the perfect model setting, the results in the biased model setting have 2 differences. First, the optimal ATM-OTW and OCN-OTW are larger than their counterparts in the perfect model setting, becoming 3 and 20 (namely the total observations are 7 and 41, respectively). Second, the RMSE curves in the space of OTWs show more **concavity and sensitive variation**. This is more distinguishable in the curve of  $\omega$ -RMSEs in the OCN-OTW space. All these phenomena can be explained by the influence of model bias on the assimilation quality. **On the one hand, due to the existence of model bias, the assimilation not only needs observations to fit the observed variability but also needs observations to reduce the mean discrepancy between the model and observation.** This requires stronger observational constraints. An optimal OTW that makes the smallest RMSE of model states must include more observed data. **On the other hand, the forecast ensemble in a biased model underestimates the forecast error, which results in that the EAKF under-weighs the observations. Therefore the optimal OTWs are larger than those in the perfect experiment case that the observations including in the optimal OTWs will be assimilated for multiple times, which results in improving of filter performance.** The test experiment for the optimal  $\eta$ -OTW is also consistent with this point (in **Fig. 8c**): the optimal  $\eta$ -OTW in the biased model setting is larger than that in the perfect model setting. Then we also investigate the influence of OTWs on the quality of CDA with the multi-variate adjustment scheme in the biased experiment framework (not shown here). Results are the same as the perfect model setting case, i.e., multi-variate adjustment scheme does not change the optimal OTWs.

Comparing the results from two experiment frameworks, we can see that regardless of perfect or biased model setting used in the assimilation experiments, the optimal OTW must be associated with the corresponding characteristic variability time scale in the medium. It is clear that while using observations in an OTW increases observational information, a too large OTW can distort the characteristic variability of coupled media during the information blending process. Therefore choosing an optimal OTW that is much smaller than the medium's characteristic variability time scale is very important. The simple model results suggest that the length of an optimal OTW is about 1-5% of the medium characteristic time scale, with which characteristic variability of the medium can be retrieved most accurately.

In this study, the OTW validates the observations in a time window to the analysis time and all the observations including in the OTW are sequentially assimilated with their original error scales. Another general approach is to assimilate the average of the observations including in the OTW but the observational errors decrease as  $1/\sqrt{N}$  of their original error scales ( $N$  represents the number of the observations including in the OTW). From the comparison of these two methods (not shown), we can see that the results obtained by them is almost same. From the perspective of the calculation process of the EAKF method, owing to that no inflation scheme was used, after many assimilation steps the ensemble spreads of the models states have been greatly reduced and significantly smaller than the corresponding observational error scales. And the prior ensemble member will be very close to the prior ensemble mean. Thus the analysis adjustments obtained by these two methods will be almost same. It is worth mentioning that although the resulted RMSEs obtained by these two assimilation schemes will be different when using the suitable inflation schemes, the lengths of the optimal OTWs are still same and the essence of this study still firms and does not change.

Also among above assimilation experiments in this study, we have not considered the temporal offset induced by the difference between the time of observations in the OTW and the analysis time. Here we can use the de-correlation coefficients to weight the observations including in the OTW and avoid overweighting them. The comparison of these two assimilation approaches (non-weighted and weighted) have been conducted (the results are not shown). From the comparison we learn that the lengths of the optimal OTWs obtained by these two assimilation schemes are similar except that the RMSEs in the weighted observation experiment will be lower than that in the non-weighted one when using longer OTWs (when the length of ATM-OTW is greater than 4 and/or that of the OCN-OTW is larger than 50). This is owing to the high correlation between the observation included in the optimal OTWs and model states at the analysis time (exceed 0.995). Thus the influence of the temporal offset can be ignored and the results obtained by these two schemes shall be almost same when using the shorter OTWs. And when we use the longer ones, the correlation will decrease and influence of the temporal offsets will be obvious that the results of the weighted observation experiment will be better. For the CDA systems in the CGCMs, owing to the complex physics and dynamics, the influence of the time offsets will be obvious and the weights of the observations will be very necessary. But from this simple model case, we can see that whether or not using the weighted observations, the relationship between the characteristic variability time scales and the optimal OTWs will be firm and the essence of this study is established.

### 5.3 Influence of coupling strength on optimal OTWs

Changing coupling strength (controlled by the coupling coefficients  $\mathcal{C}_1$  and  $\mathcal{C}_2$  in this case) between the atmosphere and upper ocean may have some influence on the characteristic variability time scales of coupled media, so as on the optimal OTWs. Test experiments show that changing the coupling coefficient  $\mathcal{C}_1$  has little influence on the characteristic variability time scales of  $\mathcal{X}_{1,2,3}$  and  $\omega$ . This is because the characteristic time scale of  $\mathcal{X}$  is determined by the chaotic nature of the Lorenz equations, not by the oceanic forcing associated with the coupling coefficient  $\mathcal{C}_1$ . Therefore, here we just change  $\mathcal{C}_2$  to investigate the coupling coefficient between the atmosphere and upper ocean on the optimal OTW of  $\omega$ . Setting the values of  $\mathcal{C}_2$  as 1.5, 1.25, 1.0, 0.8, 0.5 and 0.1 and keeping  $\mathcal{C}_1$  as 0.1, we repeat all the biased CDA experiments with the multi-variate adjustment scheme. The results are shown in **Fig. 9**, which presents the power spectrum of  $\mathcal{X}_2$  and  $\omega$  (panel a and b) of six cases above based on the model states between 5000 TUs and 9800 TUs, as well as the time series of model states between 5000 TUs and 5100 TUs (panel c) after the spin-up described in section 2.3. We can see that changing  $\mathcal{C}_2$  does not influence on the characteristic variability time scale of the atmosphere but strongly influences on variability of the slab ocean. From the equation of  $\omega$ , the characteristic variability time scale of  $\omega$  is determined by the combination of the atmospheric forcing and the periodic external forcing. When  $\mathcal{C}_2$  is small, the forcing of atmosphere to ocean is weak and then the periodic external forcing plays a dominant role on determining the characteristic variability time scale of ocean component.

Then we examine the difference of the optimal OTW of  $\omega$  in the six cases above, as shown in **Fig. 10**. **The results show that changing  $\mathcal{C}_2$  does not have any influence on the optimal ATM-OTW (not shown)**. From Panels *a* and *b* we can see that when  $\mathcal{C}_2$  is smaller, the optimal OCN-OTW is larger. This can be explained by the increasing role of the periodic external forcing on determining variability of the slab ocean, for which data assimilation needs more observational information to recover the periodic variation of  $\omega$ , determined by the time scale defined by  $\mathcal{S}_{pd}$  (10 TU). When  $\mathcal{C}_2$  is larger than 1.0, changing it has little influence on characteristic variability of the  $\omega$ , so as on the optimal OCN-OTW.

**On the one hand these experiments can further illustrate the idea that a close relationship between the length of the optimal OTW and the corresponding characteristic variability time scale exists. On the other hand, for a realistic CDA system, the coupling physics could be very complicate and affected by many factors. The results of this simple model give the insights that we can only consider the factors which have obvious influence on the characteristic variability time scales when determining the length of the optimal OTWs for a realistic CDA system. In this way the process of determining the optimal OTWs in a realistic CDA system can be greatly simplified and make it possible to apply the method of using the optimal OTWs into the realistic CDA system.**

## 6 Summary and discussions

With a simple conceptual climate model and the EAKF method, the impact of OTWs on the quality of CDA has been investigated in this study. This simple conceptual coupled model consists of a synoptic atmosphere (Lorenz 1963) and

seasonal-interannual slab upper ocean (Zhang et al. 2012) coupling with a decadal deep ocean (Zhang 2011a,b), and reasonably simulates the typical interactions between multiple time-scale components in the climate system. Determined from the characteristic variability time scale in each coupled medium, an optimal OTW provides maximal observational information to best fit characteristic variability of the medium during the data blending process. With correct scale interactions within the coupled system, CDA can recover the climate signals most accurately through incorporating all observations in the optimal OTWs into the coupled model. Although in an idealized and simple model circumstance, the conclusion addressing best fitting characteristic variability in each medium with the optimal OTW is comprehensive and therefore provides a guideline for improving climate analysis and prediction initialization when real observations are assimilated into a CGCM. For example, learned from the simple model results, we may consider to improve the quality of climate analysis and prediction initialization through accurately recovering some important characteristic variability in the atmosphere (sub-diurnal variations, for instance) and ocean (diurnal cycle in the tropical oceans, for instance).

However, the current work only can serve as a proof-of-concept study. Although CDA with the optimal OTWs has shown promising improvement in this simple model, serious challenges still exist for detecting optimal OTWs in the real world with a CGCM for improving climate analysis and predictions. First, the characteristic variability time scales in different media of the real world are complex and it remains great challenges to identify the characteristic variability of the different component models and the real atmosphere, upper and deep ocean, which need to be further studied. Also in a real ocean model, the upper and deep ocean is inseparable, which bring some troubles to use different OTWs for different parts of the same ocean model. Second, due to model biases, characteristic variability in a CGCM may be different from the real world. The combination of variability of the real world and that of the model may further complicate the problem. Therefore, model bias and its influence on model variability need to be thoroughly analysed before an optimal OTW is determined. **Thirdly, the coupling physics between different coupled components are very complicate and impacted by many factors for a realistic CDA system. Even though we only consider the factors which will obviously impact the characteristic variability time scales when determining the length of OTWs for different coupled components, it remains as heavy workload.** In addition, in this study we assume that all observations in the OTWs have equal weighting to contribute to the observational constraint. In the real observation case, the observation far away from the assimilation time should have less contribution to the state estimation at the assimilation time. How to take the time correlation into account in a sequential algorithm needs to be studied before implementing optimal OTWs into the assimilation with CGCM and real observations.

## Acknowledgements

Special thanks to Drs. Liwei Jia, Wei Zhang, Xuefeng Zhang, Wei Li, Lianxin Zhang, Shuo Yang for their comments and suggestions at the early version of this manuscript. This work was supported by the NSFC (No.51379049, 41676088), the Fundamental Research Funds for the Central Universities of China (No.HEUCFX41302, HEUCFD1505, HEUCF160410), the Young College Academic Backbone of Heilongjiang Province (No.1254G018), the Scientific Research Foundation for

the Returned Overseas Chinese Scholars, Heilongjiang Province (No.LC2013C21) and China Scholar Council (awarded to Xiong Deng for two years' study abroad at UW-Madison – NOAA/GFDL Joint Visiting Program).


## References

- Anderson, J. L.: An ensemble adjustment Kalman Filter for data assimilation. *Mon. Wea. Rev.*, 129, 2884-2903, doi:10.1175/15200493(2001)129<2884:AEAKFF>2.0.CO;2, 2001.
- Anderson, J. L.: A local least squares framework for ensemble filtering. *Mon. Wea. Rev.*, 131, 634-642, doi:10.1175/15200493(2003)131<0634:ALLSFF> 2.0.CO;2, 2003.
- Anderson, J. L.: An adaptive covariance inflation error correction algorithm for ensemble filters. *Tellus A*, 59(2), 210-224, doi:10.1111/j.16000870.2006.00216.X, 2007.
- 10 Anderson, J. L.: Spatially and temporally varying adaptive covariance inflation for ensemble filter. *Tellus A*, 61(1), 72-83, doi:10.1111/j.16000870.2008.00361.X, 2009.
- Chen, D., Zebiak, S. E., Busalacchi, A. J., and Cane, M. A.: An improved procedure for El Niño forecasting: implications for predictability. *Science*, 269, 1699-1702, 1995.
- Chen, D.: Coupled data assimilation for ENSO prediction. *Adv. Geosci.*, 18, 45-62, 2010.
- 15 Collins, W. D., Blackman, M. L., Hack, J., Henderson, T. B., Kiehl, J. T., Large, W. G. and Mckenna, D. S.: The community climate system model version 3 (CCSM), *J. Climate*, 19, 2122-2143, doi:10.1175/JCLI3761.1, 2006.
- Delworth, T. L., and Coauthors: GFDL's CM2 Global Coupled Climate Models. Part I: Formulation and simulation characteristics. *J. Climate*, 19(5), 643-674, doi:10.1175/JCLI3629.1, 2006.
- Evensen, G.: Sequential data assimilation with a nonlinear quasi-geostrophic model using Monte Carlo methods to forecast error statistics. *J. Geophys. Res.*, 99, 10143-10162, doi:10.1029/94JC00572, 1994.
- 20 Evensen, G.: *Data assimilation: The Ensemble Kalman Filter*. Springer, 187 pp, 2007.
- Gnanadesikan, A.: A simple predictive model for the structure of the oceanic pycnocline. *Science*, 283, 2077-2079, 1999.
- Hamill, T. M. and Snyder, C.: A hybrid ensemble Kalman filter-3D variational analysis scheme. *Mon. Wea. Rev.*, 128(8), 2905-2919, doi:10.1175/15200493 (2000)128<2905:AHEKFF>2.0.CO;2, 2000.
- 25 Han, G., Wu, X., Zhang, S., Liu, Z., and Li, W.: Error covariance estimation for coupled data assimilation using a Lorenz atmosphere and a simple pycnocline ocean model. *J. Climate*, 26(24), 10218-10231, doi: 10.1175/JCLI-D-13-00236.1, 2013.
- Han, G., Zhang, X., Zhang, S., Wu, X. and Liu, Z.: Mitigation of coupled model biases included by dynamical core misfitting through parameter optimization: simulation with a simple pycnocline prediction model. *Nonlinear Processes in Geophysics*, 21, 357-366, doi:10.5194/npg-21-357-2014, 2014.
- 30 **Hunt B. R., Kalnay E., Kostelich E. J. and Co-authors: Four-dimensional ensemble Kalman filtering. *Tellus A*, 56(4), 273-277, doi:10.1111/j.1600-0870.2004.00066.x,2004.**




- Houtekamer, P. L. and Mitchell, H. L.: Ensemble kalman filtering. *Quarterly Journal of the Royal Meteorological Society*, 131(613), 3269-3289, doi:10.1256/qj.05.135, 2005.
- Hunt, B. R., Kalnay, E., Kostelich, E. J. and Co-authors: Four-dimensional ensemble Kalman filtering. *Tellus A*, 56(4), 273-277, doi:10.1111/j.1600-0870.2004.00066.x, 2004.
- 5 Kalman, R.: A new approach to linear filtering and prediction problems. *Trans. ASME. Ser. D. J. Basic Eng.* 82, 35-45, doi:10.1115/1.3662552, 1960.
- Kalman, R. and Bucy, R.: New results in linear filtering and prediction theory. *Trans. ASME. Ser. D. J. Basic Eng.* 83, 95-109, doi:10.1115/1.3658902, 1961.
- Laroche, S., Gauthier, P., Tanguay, M., Pellerin, S. and Morneau, J.: Impact of the different components of 4DVAR on the  
10 global forecast system of the Meteorological Service of Canada. *Mon. Wea. Rev.*, 135(6), 2355-2364, doi:10.1175/MWR3408.1, 2007.
- Li, H., E. Kalnay, and Miyoshi, T.: Simultaneous estimation of covariance inflation and observation errors within an ensemble Kalman filter. *Quart. J. Roy. Meteor. Soc.*, 135, 523-533, doi:10.1002/qj.371, 2009.
- Liu H., Lu F., Liu Z., Liu Y. and Zhang S.: Assimilating Atmosphere Reanalysis in Coupled Data Assimilation. *Journal of  
15 Meteorological Research*, 30(4), 572-583, doi:10.1007/s13351-016-6014-1, 2016.
- Lorenz, E. N.: Deterministic non-periodic flow. *J. Atmos. Sci.*, 20, 130-141, 1963.
- Lu F., Liu Z., Zhang S., and Liu Y.: Strongly Coupled Data Assimilation Using Leading Averaged Coupled Covariance (LACC). Part I: Simple Model Study. *Mon. Wea. Rev.*, 143(9), 3823-3837, doi:10.1175/MWR-D-14-00322.1, 2015.
- Miyoshi, T.: The Gaussian approach to adaptive covariance inflation and its implementation with the local ensemble  
20 transform Kalman filter. *Mon. Wea. Rev.*, 139, 1519-1535, doi:10.1175/2010MWR3570.1, 2011.
- Pires, C., Vautard, R. and Talagrand, O.: On extending the limits of variational assimilation in nonlinear chaotic systems. *Tellus A*, 48(1), 96-121, doi:10.1034/j.1600-0870.1996.00006.x, 1996.
- Randall, D. A., and Coauthors: Climate models and their evaluation. *Climate Change 2007: The physical Science Basis*, S. Solomon et al., Eds., Cambridge University Press, 589-662pp, 2007.
- 25 Saha, S., and Coauthors: The NCEP Climate Forecast System Reanalysis. *Bull. Amer. Meteor. Soc.*, 91, 1015-1057, doi:10.1175/2010BAMS3001.1, 2010.
- Singleton, T.: Data Assimilation experiments with a simple coupled ocean-atmosphere model. Ph.D. thesis, University of Maryland, Collage Park, 116pp, 2011.
- Sugiura, N., Awaji, T., Masuda, S., Mochizuki, T., Toyoda, T., Miyama, T., Igarashi, H., Ishikawa, Y.: Development of a four-  
30 dimensional variation coupled data assimilation system for enhanced analysis and prediction of seasonal to interannual climate variations. *J. Geophys. Res.*, 113(C10), C10017, doi: 10.1029/2008JC004741, 2008.
- Whitaker, J. S. and Hamill, T. M.: Ensemble data assimilation without perturbed observations. *Mon. Wea. Rev.*, 130, 1913-1924, doi:10.1175/1520-0493(2002)130<1913:EDAWPO>2.0.CO;2, 2002.

- Yang, X. and Coauthors: A predictable AMO-like pattern in GFDL's fully coupled ensemble initialization and decadal forecasting system. *J. Climate*, 26, 650-661, doi:10.1175/JCLI-D-12-00231.1, 2013.
- Zhang, S. and Anderson, J. L.: Impact of spatially and temporally varying estimates of error covariance on assimilation in a simple atmospheric model. *Tellus A*, 55(2), 126-147, doi: 10.1034/j.1600-0870.2003.00010.x, 2003.
- 5 Zhang, S., Harrison, M. J., Rosati, A. and Wittenberg, A.: System design and evaluation of coupled ensemble data assimilation for global oceanic climate studies. *Mon. Wea. Rev.*, 135, 3541-3564, doi: 10.1175/MRW3466.1, 2007.
- Zhang, S.: Impact of observation-optimized model parameters on decadal predictions: simulation with a simple pycnocline prediction model. *Geophys. Res. Lett.*, 38(L02702), 1-5, doi: 10.1029/2010GL046133, 2011a.
- Zhang, S.: A study of impacts of coupled model initial shocks and state-parameter optimization on climate predictions using  
10 a simple pycnocline prediction model. *J. Climate*, 24(23), doi:10.1175/JCLI-D-10-05003.1, 2011b.
- Zhang, S., Liu, Z., Rosati, A. and Delworth, T.: A study of enhance parameter correction with coupled data assimilation for climate estimation and prediction using a simple coupled model. *Tellus A*, 64, 1-20, doi:10.3402/tellusa.v64i0.10963, 2012.
- Zhang, S., Winton, M., Rosati, A., Delworth, T., and Huang, B.: Impact of enthalpy-based ensemble filtering sea ice data assimilation on decadal predictions: simulation with a conceptual pycnocline prediction model. *J. Climate*, 26(7), 2368-2378,  
15 doi:10.1175/JCLI-D-11-00714.1, 2013.
- Zhang, S., Chang, Y. -S., Yang, X., and Rosati, A.: Balanced and coherent climate estimation by combining data with a biased coupled model. *J. Climate*, 27(3), 1302-1314, doi:10.1175/JCLI-D-13-00260.1, 2014.




**Figure 1:** The schematic for the assimilation interval, the length of observational time window (OTW) as well as observational interval in terms of the model integration time step. Here  $L$  represents the time steps at one side of OTW. For example, OCN-OTW ( $L$ ) in the content stands for an ocean observational time window with total observations of  $2L+1$ .

5




**Figure 2:** Time series of the absolute errors of the slab ocean variable ( $\omega$ ) in 3 assimilation experiments based on the model states between 9100 TUs and 9110 TUs assimilation results in the perfect model experiment framework with the uni-variate adjustment scheme. Green – CDA control with the standard update intervals of 0.05 TU for  $\mathcal{X}_{1,2,3}$  and 0.2 TU for  $\omega$ ; Red – CDA with an ocean observational time window (OCN-OTW) of 5 time steps [OCN-OTW (5)]; Blue – CDA with OCN-OTW (100).

10



**Figure 3:** The power spectrum (green) of a)  $\mathcal{X}_2$  b)  $\omega$ , c)  $\eta$  based on the model states between 5000TUs and 9800TUs integrations after the spin-up which integrates for 10000TUs from the initial condition (0,1,0,0) with respect to the frequency, with 95% statistics significance (red).


5



**Figure 4:** Variations of root mean square errors (RMSEs) of a) “atmospheric” states  $\mathcal{X}_{1,2,3}$  (namely the average of  $\mathcal{X}_1$ ,  $\mathcal{X}_2$  and  $\mathcal{X}_3$  RMSEs) in the space of ATM-OTW length when the “oceanic” state ( $\omega$ ) only uses a single observation at the assimilation time; b) “upper ocean” state ( $\omega$ ) in the space of OCN-OTW length when the ATM-OTW is fixed 1 as shown in panel a (1 for the ATM-OTW, i.e. 3 observations in each window, see the caption of Fig. 1) but the OCN-OTW (for  $\omega$ ) is varying and c) “deep ocean” state ( $\eta$ ) in the space of  $\eta$ -OTW length when the “deep ocean” observations are assumed to be valid and the ATM-OTW and OCN-OTW are fixed as 1 and 10, respectively. The experiments are conducted in a perfect model setting with a simple uni-variate adjustment scheme. The red lines are the 20-case mean, each using different initial conditions taken from different periods in the control integration (see description in section 2.2), and the blue lines represent the upper/lower bounds (mean  $\pm$  standard deviations) of the RMSEs. An OTW with the length of 0 represents only assimilating the observation at the assimilation time (i.e. with no OTW, dashed-black lines). The RMSE values of the control case (no observational constraint, called CTL) are marked in the parenthesis.


10

15




**Figure 5:** Same as the panels a and b in Fig. 4 but for the variation of ensemble spreads of the model states. In panel b the optimal ATM-OTW is also set as 1. The area between the lower and upper bounds (blue) represents the range evaluated from the 20 cases. **And the blue shadow below the ensemble spreads represents the range of the uncertainty of state estimation in each assimilation experiment.**

5




**Figure 6:** The auto-correlation coefficient of a)  $\mathcal{X}_2$  b)  $\omega$ , c)  $\eta$  in the space of lag times are marked by corresponding time correlation coefficients at the time scale ( $L$ ) of optimal OTWs as detected by Fig. 4 for different media (The black dashed lines). What are shown is the mean of 20 cases. In each case, an independent section (each has 10000 data of the state – 100 TUs with the interval of 0.01 TU) is used to evaluate the lag correlation coefficient. The 20 independent sections are taken from the model states apart each 200TUs between 5000TUs and 9000TUs integrations after the spin-up of 10000TUs from the initial condition (0, 1, 0, 0, 0).


10



**Figure 7:** Same as Fig. 4 but using multi-variate adjustment scheme. In panels b the optimal ATM-OTW is also set as 1.




**5 Figure 8:** Same as Fig. 4 but using the biased model setting. In panel b) the optimal ATM-OTW is set as 3. And in panel c) the optimal ATM-OTW and OCN-OTW are kept as 3 and 20, when the “deep ocean” observations are assumed to be valid.



**Figure 9:** The power spectrum of a)  $\mathcal{X}_2$  and b)  $\omega$  based on the model states between 5000 TUs and 9800 TUs integrations after the spin-up which integrates for 10000TUs from the initial condition (0,1,0,0,0) with different coupling strength ( $\mathcal{C}_2$  is set as 1.5, 1.25, 1.0, 0.8, 0.5 and 0.1 while  $\mathcal{C}_1$  keeps as 0.1.). Panel c) shows the time series of the model state  $\omega$  between 5000 TUs and 5100 TUs integrations corresponding to the six cases.

5



**Figure 10:** Panel a) is same as Panel b) in Fig. 9 but for using six different coupling strength cases (with  $\mathcal{C}_2$  values as 1.5, 1.25, 1.0, 0.8, 0.5, 0.1 while  $\mathcal{C}_1$  keeps as 0.1). Panel b) is the variation of the length of the optimal OCN-OTW with respect to the values of  $\mathcal{C}_2$ .

10