

# ***Interactive comment on “Dynamical properties and extremes of Northern Hemisphere climate fields over the past 60 years” by Davide Faranda et al.***

**Davide Faranda et al.**

davide.faranda@cea.fr

Received and published: 31 October 2017

QUERY: The authors analyze dynamical properties of daily surface pressure, surface temperature and precipitation rate using two parameters from a universal distribution derived in previous research from extreme value theory of Poincaré recurrences. The two parameters, local dimension of the phase space attractor and the inverse of the average residence time, are physically intuitive, and have the potential to characterize predictability of the analyzed variables. The manuscript is of interest for both the Dynamical System and Applied Climatology scientific communities, illustrating an approach that could be used in other regions of the world and for other variables of inter-

Printer-friendly version

Discussion paper



est. Nonetheless, the analysis raises three main issues that I think need to be clarified before the paper can be considered for publication in NLPG. In addition, a number of minor changes/clarifications are also suggested to the authors.

ANSWER: We thank the referee for the detailed comments, which we have addressed as detailed below.

QUERY: The manuscript identifies seasonality as a key regulator of the behavior of local dimension and persistence. For example, for surface pressure, two maxima and minima per year are reported, in seasons considerably different (e.g., the typical associated teleconnections are not the same in the two max seasons, with potential implications for the phase-space trajectories and their characteristics). Nonetheless, my understanding is that the composite of anomalies shown in Figure 4 for the maxima and minima are computed from the "extreme" regions in the dimension/persistence diagram (Figure 3, in the example considered here), without discriminating between the two seasons. As a result, I wonder if the present composite analysis is not averaging fields that correspond to different situations. Please clarify this point and the possible differences that could be obtained if analyzing separately the two different maxima and minima.

ANSWER: The referee is right about the seasonal dependence of the results. We have now added a new Figure 3 showing the distribution of dynamical extremes for every season. We note that in most cases the bulk of the maxima or the minima of the dimension/persistence corresponds to a single season. This is however not true for SLP, which shows a greater spread between seasons. For all the cases where two or more seasons have > 25% of the extremes, we have therefore produced separate composites for each season. These are shown in the new Fig 3. We have further added a sign test to all geographical composites to display the regions where more than 2/3 of the composite members agree on the sign of the anomalies. We have extensively modified the manuscript in Section 3 to account for the updated analysis and the new figures.

[Printer-friendly version](#)[Discussion paper](#)

QUERY: My second major concern is somehow related to the first one. Although something similar might be happening for precipitation rate (there is only one maximum, but it is so long that different climate agents might be involved, and thus perhaps it makes sense to try to sub-sample that season), I wonder if less noisier results could be obtained if a different precipitation variable is used. As the authors are probably aware, precipitation rate is in general too noisy and far less predictable than accumulated rainfall or rainfall frequency, the latter being the most predictable of all three – at least in most parts of the world. How much would Figure 10 change if frequency of precipitation is used?

ANSWER: We thank the referee for the very good suggestion of using precipitation frequency. In the new version of the manuscript we have substituted the precipitation rate with the daily precipitation frequency. This variable is constructed as follows: for each grid point, if it rains during a specific day, we set the value to 1, and to 0 if it doesn't rain. With this choice, we obtain results that we believe correspond to a clearer narrative. As a result, we have radically altered the section of the manuscript dealing with the rainfall analysis.

QUERY: I also think that it is adequate that the authors include at least one comment on the fact that considering all grid boxes in the Northern Hemisphere might be "masking" the predictability and/or the dynamical properties of the variables in study, compared to performing the analysis in regions that are known to have a more homogeneous climatic response to different sources of predictability. In other words: how different are the results if instead of considering the entire Northern Hemisphere, regions of homogeneous predictability are considered? Seasonality is different for different regions, so is it possible that mixing the seasonal cycle of so many places degrades the characterization of the dynamical properties of the system?

ANSWER: In a recent paper (Faranda et al. 2017, Sci. Rep.), we analyzed a specific region of the northern hemisphere, i.e. the North Atlantic, which is more homogeneous in terms of predictability. Indeed the results obtained at the regional scale do not always

[Printer-friendly version](#)[Discussion paper](#)

match those obtained for the full Northern Hemisphere. However, rather than ascribing this discrepancy to a “masking” we rather interpret it as meaning that, despite the large regional differences, our methodology still allows us to draw interesting conclusions about the full hemispheric dynamics. Of course, other studies on restricted regions may answer more specific questions about the predictability and the persistence of certain patterns. Our study remains mostly theoretical at this stage and focuses on the general properties of the atmospheric attractor. We have clarified this in the conclusions of the manuscript.

QUERY: P2L12: excise "the" before "attractor"

ANSWER: Corrected

QUERY: P2L33: how many are "enough"?

ANSWER: We have now specified this.

QUERY: P2L34-P3L1: something is wrong. There is a "." misplaced, and I think the trajectory is around  $\xi$ , not around  $\theta(\xi)$ , which has not been defined yet.

ANSWER: We have corrected these typos.

QUERY: P3L7: I suggest to change "air" by something else. Several of the readers of the paper are going to be fast-readers, and in several other places "air" might be misleading. Perhaps use "t2m", which is one of the standards.

ANSWER: We have replaced “air” by “t2m” as suggested.

QUERY: P3L8: what do the authors mean by "peculiar subspace"? Clarify.

ANSWER: We have now explained this terminology.

QUERY: P5L2: further explain what the “length” of a cluster is.

ANSWER: This is now explained.

QUERY: P5L15-16: excise the subjective phrasing "a very".

ANSWER: Done.

QUERY: P5L22: maybe the semi-annual cycle is inherited from some semi-annual variability phenomena too? e.g., North Atlantic Sub-tropical High?

ANSWER: Good suggestion, we have emphasized this in the discussion of the slp results.

QUERY: P5L28: why is it more restrictive?

ANSWER: Because if one considers the typical partition into 4 weather regimes, the probability of being in one of them is roughly  $1/4=0.25$ , whereas the probability of being close to zeta is set by the quantile: for  $q=0.98$ , the probability of being close to zeta is 0.02. This is now better explained in the text.

QUERY: P6L16-17: it seems to me that the weakened Aleutian low is actually a high in that figure. Can you comment on that? Also, the magnitudes of the anomalies are really low (amplitude of 10 hPa). This might be associated, again, with the fact that all days are considered together, without seasonal discrimination. The Aleutian low is typical of winter; can the authors comment on why it should be expected to appear during the maxima of d

ANSWER: The plots show anomalies relative to a climatological seasonal cycle, so that a positive anomaly over a climatological low would correspond to a weakened low (provided that the anomaly is not so strong as to turn a low into a high). However, following the Reviewer's first major comment we have radically altered the presentation and discussion of the geographical composites, so that the figure mentioned here is not present in the revised version of the manuscript.

QUERY: P6L26: please explain why that means intra-seasonal correlation.

ANSWER: We have removed this sentence and now discuss the cross-correlation only in reference to Figure 4b.

Printer-friendly version

Discussion paper



QUERY: P7L11: any hypothesis for why is it absent?

ANSWER: This is linked to the slight offset in the minima of the seasonal cycle of  $d$  and  $\theta$ , which result in a positive ACF for  $d$  between winter and summer and a negative one for  $\theta$ . We illustrate this in Fig. R1 below, which shows the  $d$  and  $\theta$  curves for  $t_2m$  shifted by 6 months. We now briefly mention this in the text. Fig. R1: Local dimension (continuous lines) and  $\theta$  (dashed lines) for  $t_2m$ . The blue lines are lagged by 6 months relative to the red lines.

QUERY: P7L22: Explain what you mean by "high-dimensional situation".

ANSWER: Here we meant a situation with a high  $d$  value, and hence a high number of active degrees of freedom, which makes the evolution of the atmospheric state inherently hard to forecast. We have now rephrased this to clarify our point.

QUERY: P7L30: it is a bit strange that extremes cannot be matched to large-scale patterns, as they tend to be controlled by synoptic configurations. Maybe what the authors mean is that there is no global pattern associated with the extremes? I do not expect that case either. Right now, the sentence is confusing.

ANSWER: We have now repeated the analysis with a different variable (precipitation frequency) and this sentence has been removed as this is not the case anymore.

QUERY: P8L12: "Precipitation rate"

ANSWER: We have now corrected this to "Precipitation Frequency" to reflect our updated analysis. QUERY: P11L9-10: use "twenty" and "fifty", or "20" and "50".

ANSWER: Format of numbers is now consistent.

QUERY: P11L16: along the paper, the authors use 1000 hPa. Did they interpolate? Or do they just mean "surface". If the latter, please change to "surface".

ANSWER: 1000 hPa has been changed to surface.

Printer-friendly version

Discussion paper



QUERY: P11L26-27: I actually expect to see a correlation between pressure gradients and monsoonal precipitation. Maybe I'm not understanding something here.

ANSWER: We have now included a revised explanation in this section discussing the new precipitation frequency results.


---

Interactive comment on Nonlin. Processes Geophys. Discuss., <https://doi.org/10.5194/npg-2017-36>, 2017.

Printer-friendly version

Discussion paper





**Fig. 1.** Local dimension (continuous lines) and theta (dashed lines) for t2m. The blue lines are lagged by 6 months relative to the red lines.

Printer-friendly version

Discussion paper

