Response to Reviewer#1's comments

on paper "Tipping point analysis of ocean acoustic noise"

by V. N. Livina, A. Brouwer, P. Harris, L. Wang, K. Sotirakopoulos, S. Robinson

We would like to thank the Reviewer for the helpful comments on the paper. We have prepared a marked manuscript with added text as described below.

- **Comment 1.** Fig. 4: The record is only 14-yr long, and there is definitely a lot of internal variability in the climate system that can contribute to decadal trends on top of any possible anthropogenic effects.
 - **Response.** We agree with the reviewer that there may be various causes for the trend, and antropogenic influence may be only a part of contribution on the background of the various climatic factors this is why, in particular, we investigate the connection with El Niño events.

Acoustic noise at the depth of one kilometre can be influenced by multiple factors of natural and anthropogenic origin, and we discuss some of the possible components that could represent the dynamics of the acoustic system. While there may be various equivalent models reproducing the observed time series, we choose the simple stochastic dynamic system which generates simulated time series with very close statistical properties.

It has already been reported that antropogenic acoustic signals affect vocalizing baleen whales, with distance of impact up to 200 km [Risch et al 2012]: such disturbances cause behavioural changes in large animals, and consequently reduce their acoustic presence in the area of observation. This may be one of the reasons of declining trend in the ocean acoustic noise in particular frequency ranges.

We have added this paragraph into the manuscript.

- **Comment 2.** Fig. 5: The text claims a connection with ENSO, but its not immediately obvious from comparing the top and bottom panels of the figure. Could the authors produce a bit more quantitative measure of this association (for example, plotting a 1-D time series of the color plot and correlating it with ENSO indices)?
 - **Response.** We have attempted to generate a time series from the coloured potential plot at a chosen time scale (0.5 yr). Note that the values of such time series can only be integer numbers, as the variable is the number of the potential wells in the dynamical system. On the other hand, the El Niño indices are continuous variables, which denote an event when they are varying below or above certain level. To make them comparable with the extracted potential metrics described above, we produce a binary variable for the ONI index time series: such variable

is 1 when ONI is bigger than 0.2 (different sources suggest different threshold, sometimes 0.5, sometime 0 as indication of El Niño events). The varying number of wells in the potential plot was converted into a binary index, which equals 1 when there is "non-regular" (not triple-well-potential) state of the dynamic system. Accordingly, we have prepared a new version of Fig.5, with three panels, where this is illustrated.

For several known El-Nino events (2003, 2004, 2006, 2009, 2015-2016), there is agreement between the two binary indices. The potential index has several additional oscillations which might have been caused by different factors among many that affect this complex dynamic system (yet another possible cause may be La Niña events, when ONI index is negative — see the new version of Figure 5).

We have added this text in the manuscript.

- Comment 3. Fig. 6: Such skewed distributions can be generated by the stochastic models with nonlinear deterministic part driven by the additive noise, but also with linear models driven by multiplicative noise. See Compo et al. (2015) and references therein (http://journals.ametsoc.org/doi/abs/10.1175/JCLI-D-15-0020.1). So the choice of the appropriate stochastic model may not be that obvious.
 - **Response.** As we understand it, the reviewer mentions the paper by Sardeshmukh et al, JoC 2015. The paper considers extreme weather statistics and warns against seeking anthropogenic components in data with heavy-tailed non-Gaussian distributions. We agree with this statement, and we do not attempt to introduce an anthropogenic term in our model. However, the acoustic impact of shipping and other anthropogenic factors on the marine wildlife has been reported (as in the reference above and those cited in the paper), and we add this as a discussion topic rather than a modelling component.

We have added this reference and a few sentences on this into the manuscript.

- Comment 4. Section 5, eq. (11). I am confused about the model. Should it be z-dot (not zprime) on the left-hand side, as in (4)? But then are T(t) and P(t) added on top of the potential model driven by the red-noise (Phi)? In this case, they cannot be the part of Eq. (11)? Do the coefficients of the potential continuously depend on time (as implied by Fig. 5)? Or is the model trained on subsamples of data with a given potential structure (e.g., two-well, three-well)?
 - **Response.** The reviewer is obviously right about the typo in the left-hand-side of Eq.11, it is derivative in t and should be \dot{z} . We have corrected this.

The potential term and the stochastic noise in the model equation are used to reproduce the fluctuations that are obtained after subtraction of seasonal and long-term linear trend. The data that is modelled with the potential forecasting technique is shown in Figure 1, right panels. These fluctuations also include the red noise component. Its characteristics, from our point of view, do vary with time, as Figure 4 shows.

The original data (Figure 1, left panels) can be reproduced by combining all the components of model (11), which is a version of the nonautonomous Langevin equation. We follow this modelling approach as it was used in paleostudies, see for example [Ditlevsen et al, JoC 2005], although paleodata has different temporal resolution and patterns. This illustrates the flexibility of the stochastic modelling approach and its general applicability.

We have added this paragraph into the paper.

- **Comment 5.** Is it really a surprise that the model reproduces the observed statistics? This section needs to be clarified.
 - **Response.** In our opinion, only by taking into account all the components of the proposed model, i.e. the global trend, the seasonal trend, the asymmetric system potential structure, and the long-term correlated red noise, one can reproduce the considered acoustic data. Our analysis allows one to understand these main components and derive their specific parameters, which are then used to forecast data, and thus validate our understanding and the proposed stochastic model. Indeed, there may be other models that can produce similar structure of time series. The advantage of our model is its simplicity and adequate representation of the main geophysical processes of this dynamical system.

The hypothesis of the possible influence of El Niño appeared in the course of our research and was unexpected. Therefore, our modelling approach, in principle, is capable of discovering such interesting signatures in the data for further investigation. This demonstrates the capability of the proposed data analysis, on its part, to stimulate geophysical research.

We have added this text into the paper.

We hope that the revised manuscript is now suitable for publication in the Nonlinear Processes in Geophysics.

Yours sincerely,

V. N. Livina, A. Brouwer, P. Harris, L. Wang, K. Sotirakopoulos, and S. Robinson



Figure 1: Upper panel: potential analysis plot of the broadband SPL data, with varying window length (y-axis) from one day to one year. The colours denote the number of detected potential states: green two, cyan three. Specks of magenta denote very short periods of a higher number of states, which correspond to highly variable (possibly non-potential) subsets of data of small size. Middle panel: ENSO indices ONI and SOI, known to be anti-correlated, which indicate several ENSO events (El Niño and La Niña). These can also be seen in the potential analysis plot. ONI positive and negative values (roughly corresponding to El Niño and La Niña) are shaded by light red and light blue respectively, for better comparison with the indices in the lower panel. Lower panel: binary indices derived from the ONI index and potential plot (at the level 0.5 year of y-scale in the top panel) which have values 1 when there is an El Niño (in the case of ONI-based binary index) or anomalous potential state (in the case of the potential binary index).