

Please, find the revised version of the manuscript entitled “Parametric covariance dynamics for the nonlinear diffusive Burgers equation”. The comments of the both referees have been taken into account in the revised manuscript.

We have improve the description of the numerical experiments with some details on the implementation used: finite difference for the spacial discretization, a fourth order Runge-Kutta for the time scheme and we have specified the numerical setting (time step, numerical value of the diffusion coefficient). The ensemble size has been increased to 6400 in order to limit the sampling noise, and a single ensemble of normalized error has been generated then used with appropriate initial error magnitude – this reduces the sampling fluctuations when comparing the numerical results from a method to another.

In order to investigate the limitation of the tangent-linear covariance dynamics the manuscript incorporates new results (even if further research are still needed to investigation more completely the limitations of the parametric formulation as highlighted in the manuscript):

A study of the mean predicted by the parametric model and estimated from the ensemble has been introduced in order to illustrate the ability of the PKF to provide an estimation of the true mean state when small non-linearities are present: see Fig. 4 and the new section 4.3.1. A long term experiment has been introduced to determine if there is an exponential growth of the error that could be a side effect of the tangent-linear approximation: see Fig. 8 and the new section 4.3.3. The discussion of the results has been put in a new section 4.3.4.

Here are the answers to the referees:

Reviewer 1:

We would like to thanks S. E. Cohn for his comments and suggestions.

Major comments:

two errors:

1) *“First, there is an error passing from Eq. (27b) to Eq. (28b): a coefficient 2 has crept into the third-from-last term on the right side of Eq. (28b) which does not belong there, and it is repeated in the final Eq. (29c). If this is just a typo, it simply needs to be corrected. But if this error has also made its way into the computer code, then the numerical experiments will need to be re-run.”*

This has been corrected, thank you very much.

2) *“Second, the Gaussian initial covariance function, Eq. (30), is not appropriate for the geometry of the numerical experiments. Since the domain is periodic, the distance $|x-y|$ should be replaced by a distance function that reflects this periodicity, such as the great-circle distance or chordal distance. As it stands, the covariance function has a slight first-derivative discontinuity at $|x-y| = D/2$, and this introduces spurious odd-order terms in the series expansion of the correlation function that have been neglected. Although this might be a small effect initially, since the initial correlation length L was taken to be small, the numerical experiments showed that the correlation length grows by nearly an order of magnitude. The numerical experiments will need to be repeated with an appropriate initial covariance model.”*

For the experiment considered in the manuscript, only the initial covariance Eq.(30) (from the previous version) is needed. Since the length-scale considered is relatively small considering the length of the domain, the Gaussian correlation applies here. However, we agree with the referee that this is not strictly a valid correlation function, and a chordal distance has been introduced in

accordance with the geometry of the domain. This modification does not change the results but is theoretically better.

Minor comments and typos:

1. *It is mentioned a few times that “operator splitting” (p.2 l.25) or “time splitting” (p.6 l.26) is used in the derivation. Actually, the authors are simply carrying out the derivation term-by-term without any approximation introduced by doing so. The authors’ use of this terminology is not at all standard; usually it means that a time discretization error is introduced in a numerical method. I suggest removing the terminology altogether.*

We agree with the referee that the terminology “time-splitting” could introduce a confusion with the classical numerical time-splitting. In order to avoid this confusion, we have introduced the following lines in section 3.1, where the splitting method is mentioned:

“The splitting strategy is a theoretical method to deduced the so-called infinitesimal generator of an evolution equation, by taking advantage of the Lie-Trotter formula to separate each processes (or appropriate arrangements of the processes). This strategy should not be confused with the numerical time-splitting which introduces numerical errors (Sportisse, 2007).”

2. *P.7 l.13: I would change “Hilbert space” to the more general “function space” since no Hilbert space apparatus has been introduced in the paper.*

Yes the Hilbert structure is not important here and it has been replace by “function space” as suggested by the referee.

3. *In the title itself, the apostrophe after Burgers should be removed: the possessive is not correct here.*

The typos is now corrected.

4. *P.4 l.16: recipes → recipe*

The typos is now corrected.

5. *P.6 l.12: te → the*

The typos is now corrected.

6. *Eq. (25): One appearance of δx^2 in the second term, and one in the third term, should be removed.*

The Taylor expansion Eq.(25) has been corrected.

7. *P.10 l.8: express → expressed*

The typos is now corrected.

8. *Eq. (29): The subscript x on the symbol V should be used consistently.*

The subscript x has been removed from Eq.(29).

9. *P.13 l.33: 8.8 of → 8.8 times*

The typos is now corrected.

10. *P.14 l.9: 5.5 of → 5.5 times*

The typos is now corrected.

11. *P.14 l.12: 7.5 of → 7.5 times*

The typos is now corrected.

12. P.18 l.13: *variance field* → *normalized variance field*

The typos is now corrected.

13. P.18 l.21: *third term* → *third order term*

The typos is now corrected.

14. P.19 l.1: *fourth term* → *fourth order term*

The typos is now corrected.

15. P.19 l.8: *third order term* → *fourth order term*

The typos is now corrected.

Rev. 2:

We would like to thanks referee 2 for his/her comments.

1) page 1 Abstract: “Abstract: I am not sure if you can say that “this study extends the PKF to nonlinear dynamics”. Rather, It is a required preliminary step.”

Since the parametric model is designed to the nearly nonlinear dynamics we think this terminology is appropriate. This is supported by the introduction of the mean predicted by the ensemble versus the parametric model (Fig. 4) where the mean is different from the reference state at T.

2) page 2 line 20 “numerical tests”

The typos is now corrected

3) page 3 line 2 “Background on the”

The typos is now corrected

4) page 4 line 15 I am not really keen to use a semigroup in order to define a multivariate matrix. What stops you writing the matrix explicitly ?

The matrix results from numerical integration of the diffusion equation, it is not easy to write the matrix explicitly.

5) page 5 line 10 “meaningful”

The typos is now corrected.

6) page 5 line 13 The dimension of the space plays no role in here.

We agree with the referee comment but we think it is interesting to extrapolate the 1D situation to the 2D/3D case much important for further applications, where Eq. (11) offers a systematic derivation of the dynamics, whatever the dimension is.

7) page 6 line 5 Can I suggest that you avoid using the term “nonlinear Kalman filter”.

The Kalman filter is, by definition, linear. Yes, there are extensions of the Kalman filter methodology to nonlinear frameworks (the extended Kalman Filter, the ensemble Kalman filter, etc), but I believe that here you are referring to the framework which is either nonlinear or linear (Kalman).

The terminology has been employed by Cohn (1993) and corresponds to the next order of the extended Kalman filter where the magnitude of the fluctuation influences the mean state. The sentence has been rephrase as follows:

“ Note that the fluctuation-mean flow interaction leads to the Gaussian second-order filter \citep[sec. 9.3]{Jazwinski1970book},

and is important in nonlinear Kalman-like filters \cite{Cohn1993MWR}.”

8) page 6 line 12 “deduced from the difference”

The typos is now corrected.

9) page 7 line 21 As I understand it, $\overline{\partial x \varepsilon^2}$ is the expectation of $\partial x \varepsilon^2$, not the approximation obtained by the ensemble average.“

Yes this is correct: in this expression there is no approximation, and you are right this corresponds to the expectation operator of the derivative of ε^2 . At a theoretical level, the “ensemble average” considering an infinite ensemble size is equivalent to the expectation operator. And this infinite ensemble average should not be confused with the finite ensemble average as used with an EnKF: the theoretical derivation does not rely on any finite ensemble as encountered in the EnKF. The deterministic equations Eq.(29) of the parametric model does not need any ensemble.

10) page 9 line 13 There is a comma missing after the exponential.

The typos is now corrected.

11) page 10 line 8 “expressed” .

The typos is now corrected.

12) page 10 line 23 How large is the ensemble for the numerical experiment ? Why do you call it nonlinear ?

The ensemble size was mentioned in p11, line 12: 1600 members. The sentence has been rephrased as: “Then, the PKF is assessed using a large ensemble of nonlinear forecasts (6400 members)”.

13) page 11 Caption for Figure 1: the figure has 6 curves. I assume you plot the solution at 0.8T too (same for Figure 4 page 15, Figure 5 page 16) :

The time 0.8T has been added in the caption.

14) page 13 line 17 What does the 17.6% figure signify ?

This refers to the result at time $t=T$. The sentence has been rephrase to clarify: “with a low relative error $\frac{\|K - K_{GC}\|}{\|K\|}$ of 9.4% (respectively 17.6%) at time 0 (respectively T), ”

15) page 14 Figure 3 What is the maximum difference between the two graphs ?

We have precise the maximum difference of the two kurtosis normalized by K_G : 0.05

16) page 15 line 10 In the case of Burgers’ equation, can you provide an estimate of the length of the time interval throughout which the method presented in the paper gives meaningful results.

In order to improve this point, additional results have been incorporated in the manuscript:

-1- Prediction of the mean:

For the Burgers equation, in the parametric model Eq.(29), the dynamics of the mean flow is the true dynamics of the ensemble mean. We have insisted on the exactness of Eq.(14a) to predict the ensemble mean with the sentence:

“Moreover, as pointed out in Ménard (1994),

Eq. (14a) is the exact the ensemble mean for the Burgers dynamics, while Eq. (14b) is an approximated dynamics. As a consequence, if the variance field is the true one, then the mean predicted by Eq. (14a) is the true ensemble mean (Ménard, 1994, sec. 5.5.2).” (sec. 3.1)

The new section 4.3.1 illustrates the case of the mean (see also the new Fig. 4) .

-2- Introduction of a long term experiment:

The new Fig. 8 and new section 4.3.4 have been introduced to tackle (but only in part) the time where the parametric model is valid.

17) page 16 Figure 16. Again, it would help to know how large is the ensemble used in the numerical experiments.

In the new version of the manuscript, the ensemble size has been increased to 6400 with a single ensemble of normalized error to facilitate the comparison of the numerical experiments.

18) page 17 line 22 "The aim of this section"

The typos is now corrected.

19) page 18 line 2,4 Something went wrong with the display of the formulae (A3) and (A4)

This is due to the "manuscript version" configuration of the manuscript (single column while "article version" is two columns). The two column version of the manuscript does not present this wrong display.

Parametric covariance dynamics for the nonlinear diffusive Burgers ~~’~~-equation

Olivier Pannekoucke¹, Marc Bocquet², and Richard Ménard³

¹ INPT-ENM, CNRM UMR 3589, Météo-France/CNRS, CERFACS, Toulouse, France.

² CEREAs, joint laboratory École des Ponts ParisTech and EDF R&D, Université Paris-Est, Champs-sur-Marne, France

³ ARQI/Air Quality Research Division, Environment and Climate Change Canada, Dorval (Québec), Canada.

Correspondence to: O. Pannekoucke (olivier.pannekoucke@meteo.fr)

Abstract. The parametric Kalman filter (PKF) is a computationally efficient alternative method to the ensemble Kalman filter (EnKF). The PKF relies on an approximation of the error covariance matrix by a covariance model with a space-time evolving set of parameters. This study extends the PKF to nonlinear dynamics using the diffusive Burgers ~~’~~-equation as an application, focusing on the forecast step of the assimilation cycle. The covariance model considered is based on the diffusion equation, with the diffusion tensor and the error variance as evolving parameter. An analytical derivation of the parameter dynamics highlights a closure issue. Therefore, a closure model is proposed based on the ~~so-called~~-kurtosis of the local correlation functions. Numerical experiments compare the PKF forecast with the statistics obtained from ~~an~~-a large ensemble of nonlinear forecasts. These experiments strengthen the closure model and demonstrate the ability of the PKF to reproduce the tangent-linear covariance dynamics, at a low numerical cost.

10 *Copyright statement.* TEXT

1 Introduction

Covariance functions in geophysical flows are known to ~~be evolving~~-evolve in both time and space (e.g., Bouttier, 1993; Snyder et al., 2003). Yet, an accurate solution of the covariance dynamics is one of the major challenges in data assimilation and probabilistic forecasting. The Monte Carlo method, which is the most common approach, addresses nonlinear dynamics and is computationally efficient with parallel computers. However, it suffers from non-uniform sampling noise, which is a function of the true signal covariance.

Another route can be investigated that relies on analytical derivation of covariance tensor dynamics (Cohn, 1993), which has inspired application in chemical transport model (Ménard et al., 2000). Despite the theoretical interest resulting from the analytical derivation of covariance dynamics, it is still difficult to take advantage of this formulation in real applications. Moreover, the system presents closure problem for the diffusive error dynamics. An hybrid approach that mixes the Monte Carlo method based on an ensemble and an approximate propagation of the correlations by a surrogate model has also been proposed (Bocquet, 2016).

An intermediate formulation, between the approximation by an ensemble and the theoretical formulation by analytic derivation, has been recently introduced by Pannekoucke et al. (2016) who proposed to approximate the forecast error covariance matrix by a parametric covariance matrix, from which the dynamics of parameters ~~is a surrogate to stand for~~ the dynamics of the full covariance matrix. This formulation, ~~so-called~~ called the Parametric Kalman Filter (PKF), has been illustrated on a linear
5 advection-diffusion equation similarly to the equations encountered in ~~Chemical Transport Model~~ chemical transport model. As defined for general parametric covariance model, the PKF has been illustrated for the particular case where the covariance model is based on the diffusion equation (Weaver and Courtier, 2001). Hence, the error covariance matrix is reduced to the knowledge of its variance field and its local diffusion tensor field. The time evolution all along the forecast and the analysis steps of the data assimilation process is expressed in terms of variance and local diffusion tensor evolution.

10 As mentioned earlier, the PKF formulation has been tested so far on a linear dynamics. It is thus interesting for more general applications to consider extension to a ~~non-linear~~ nonlinear setting. The ~~motivation~~ goal of the present work is to formulate and illustrate the forecast step of the PKF for the ~~non-linear~~ nonlinear dynamics given by Burgers' equation. Burgers' equation is a nonlinear advection-diffusion model that usually involves one variable in a one-dimensional space – the u –wind. It is one of the simplest equations that display important features of geophysical interest, such as advection, frontogenesis and 1D
15 turbulence (Burgers (1974); Hopf (1950); el Malek and El-Mansi (2000) and references therein). Burgers' equation has been used in several data assimilation studies to examine the effect of nonlinearity error propagation and in Kalman filtering methods (Cohn, 1993; Ménard, 1994; Verlaan and Heemink, 2001), in maximum likelihood ensemble filtering (Zupanski et al., 2008), in adjoint methods (Apte et al., 2010), in model error estimation using 4D-Var (Lakshminarayanan et al., 2013), and in 4D-EnVar and ~~4D~~-localization (Desroziers et al., 2014, 2016).

20 However, preliminary numerical ~~test~~ tests have shown that the treatment of the physical diffusion as proposed in Pannekoucke et al. (2016), and deduced from analytical solution, was not able to reproduce the complexity of the Burgers dynamics. Hence, we need to develop a higher order representation of the PKF equation for the physical diffusion process.

In section 2 the parametric formulation based on the covariance modeling with the diffusion equation is first recalled, and we specify the methodology for developing the parametric dynamics under a systematic treatment. This method is then applied to
25 the Burgers' ~~equation~~, equation, in section 3, taking advantage of the operator splitting. In section 4, numerical simulations are conducted to illustrate the ability of the parametric dynamics to reproduce the main features of the true covariance dynamics emerging from a forecast Monte Carlo experiment. The conclusions are given in section 5.

2 Parametric formulation of covariance dynamics

2.1 ~~background~~ Background on the uncertainty propagation and covariance dynamics

Geophysical flow dynamics can be represented as a nonlinear ~~dynamical~~ system of the form

$$\begin{cases} \partial_t u = \mathcal{M}(u), \\ u(\mathbf{x}, 0) = u^0(\mathbf{x}), \end{cases} \quad (1)$$

5 that describes the time evolution of a state function u and where a unique solution is assumed to exist for any initial condition u^0 within an appropriate set.

Due to the lack of precise knowledge of the initial condition, u^0 is generally represented as a random state $u^0 = \bar{u}^0 + \varepsilon^0$ where ε^0 is a centered Gaussian random field characterized by its two-point covariance function $P^0(\mathbf{x}, \mathbf{y}) = \overline{\varepsilon^0(\mathbf{x})\varepsilon^0(\mathbf{y})}$, where $\overline{(\cdot)}$ stands for the expectation operator. The covariance function can be described by its variance field V^0 , where $V_{\mathbf{x}}^0 = P^0(\mathbf{x}, \mathbf{x})$ denotes the variance at position \mathbf{x} and by its error correlation function $\rho^0(\mathbf{x}, \mathbf{y}) = P^0(\mathbf{x}, \mathbf{y})/\sigma_{\mathbf{x}}^0\sigma_{\mathbf{y}}^0$ where $\sigma_{\mathbf{x}}^0 = \sqrt{V_{\mathbf{x}}^0}$ ~~(and~~ $\sigma_{\mathbf{y}}^0 = \sqrt{V_{\mathbf{y}}^0}$ ~~) is the standard deviation are the standard deviations~~ at point \mathbf{x} ~~(and~~ \mathbf{y}), respectively. When a first order Taylor expansion accurately approximates the error dynamics, then the tangent-linear dynamics of the initial error ε^0

$$\begin{cases} \partial_t \varepsilon = M\varepsilon, \\ \varepsilon(\mathbf{x}, 0) = \varepsilon^0(\mathbf{x}), \end{cases} \quad (2)$$

~~can accurately represent the evolution of~~ makes evolve the error about the mean state \bar{u} . $M = \partial_u \mathcal{M}|_{\bar{u}}$ is the tangent-linear dynamics along the nonlinear trajectory, \bar{u} , solution of Eq. (1) starting from the initial condition \bar{u}^0 .

The two-point covariance function $\overline{\varepsilon(\mathbf{x}, t)\varepsilon(\mathbf{y}, t)}$ of the error field ε at a given time t defines the covariance function $P(\mathbf{x}, \mathbf{y}, t)$. Thereafter, the covariance function is computed as a covariance matrix: when space is discretized, with the grid-point positions denoted by \mathbf{x}_i , the restriction of the covariance function to the grid-point positions is the matrix \mathbf{P} defined by $\mathbf{P}_{ij}(t) = P(\mathbf{x}_i, \mathbf{x}_j, t)$. With the discretized version of the tangent-linear model M being now the matrix \mathbf{M} , the dynamics of the covariance matrix is then given by the forecast error covariance equation

$$\begin{cases} \frac{d}{dt} \mathbf{P} = \mathbf{M}\mathbf{P} + \mathbf{P}\mathbf{M}^T, \\ \mathbf{P}(t=0) = \mathbf{P}^0, \end{cases} \quad (3)$$

where \mathbf{M}^T stands for the adjoint of the tangent-linear model \mathbf{M} . Thereafter, since the statistics depends from the time evolution, the explicit reference to the time t is dropped, excepted for the initial time $t = 0$ identified by superscript $(\cdot)^0$.

The numerical cost ~~to solve this covariance dynamics, of solving~~ Eq. (3) ~~, for high-dimensional dynamics~~ is beyond super-computer capacity. Different options have been considered in the literature to approximate the solution, among which one finds the ensemble Monte Carlo method employed in ensemble Kalman filter (Evensen, 1994).

The ensemble Kalman filter is a robust algorithm that applies to low order dynamical systems as well as to large dimension systems encountered in geophysical applications. The main differences for geophysical applications is that the covariance

matrix is closely related to the continuous covariance function, which may not be the case for all discrete low-order models. Thereafter, it is assumed that a discrete model results from the discretization of a continuous model, making a clear connection between the discrete and the continuous covariance representations. This offers simplifications in the following derivations. To that end, in what follows, the covariance function $P(\mathbf{x}, \mathbf{y})$ and its grid-point matrix representation \mathbf{P} are considered as equivalent, and denoted by the matrix notation.

We now give details about another approximation, that relies on the continuum, namely the parametric formulation.

2.2 Parametric formulation of the covariance forecast dynamics

Pannekoucke et al. (2016) have proposed to parameterize the covariance matrix by the covariance model, and they have illustrated this approach by considering the covariance model based on the pseudo-time diffusion equation (Weaver and Courtier, 2001).

The diffusion covariance model factorizes the covariance matrix as

$$\mathbf{P}^{\text{diff.}} = \mathbf{\Sigma} \mathbf{L} \mathbf{L}^T \mathbf{\Sigma}^T, \quad (4)$$

where $\mathbf{\Sigma}$ denotes the diagonal matrix of standard deviation $\sqrt{\varepsilon^2}$, and \mathbf{L} is the propagator of the diffusion equation

$$\partial_\tau \alpha = \nabla \cdot (\boldsymbol{\nu} \nabla \alpha), \quad (5)$$

integrated from the pseudo-time τ from $\tau = 0$ to $\tau = 1/2$, thus giving $\mathbf{L} = e^{\frac{1}{2} \nabla \cdot (\boldsymbol{\nu} \nabla)}$. The pseudo-time diffusion equation is a [recipe-recipe](#) to build Gaussian random field with Gaussian-like correlation functions. Note that, the pseudo-time τ has no link with the physical time t of Eq. (1). In this formulation the variance field V (deduced from $\mathbf{\Sigma}$) and the local diffusion tensor field ($\boldsymbol{\nu}$) are the only parameters to be determined.

Knowing the dynamics of the variance field V and the local diffusion tensor field $\boldsymbol{\nu}$ provides a [mean-means](#) to approximate the true covariance dynamics Eq. (3), where \mathbf{P} would be replaced by the covariance model $\mathbf{P}^{\text{diff.}}$ parameterized by using the diffusion equation, Eq. (4). This constitutes the parametric formulation of the dynamics. The challenge is now to determine the dynamics of the two parameter fields.

The dynamics of the variance field $V = \overline{\varepsilon^2}$ can straightforwardly be obtained from the trend $\partial_t \varepsilon$ following

$$\partial_t V = 2 \overline{\varepsilon \partial_t \varepsilon}. \quad (6)$$

However, the dynamics of the diffusion tensor is not as obvious to derive. A possible way to describe its dynamics is to consider some approximation that we will describe in the next section.

2.2.1 Approximate dynamics for the diffusion covariance model

The dynamical equations of the local diffusion can be obtained taking advantage of approximations used in data assimilation, for the estimation of the local diffusion tensor from ensemble data.

Following Pannekoucke and Massart (2008); Mirouze and Weaver (2010); Weaver and Mirouze (2013), the local diffusion tensor field can be deduced from the correlation functions, when assuming that the random error field is smooth. For a given position \mathbf{x} , the local Taylor expansion of the correlation function $\rho(\mathbf{x}, \mathbf{x} + \delta\mathbf{x})$ is related to the local correlation function in the form

$$5 \quad \rho(\mathbf{x}, \mathbf{x} + \delta\mathbf{x}) = 1 - \frac{1}{2} \|\delta\mathbf{x}\|_{\mathbf{g}_x}^2 + o(\|\delta\mathbf{x}\|^2), \quad (7)$$

where \mathbf{g}_x denotes the local metric tensor at point \mathbf{x} , with $\|\delta\mathbf{x}\|_{\mathbf{E}}^2 = \delta\mathbf{x}^T \mathbf{E} \delta\mathbf{x}$. In this expression little o means that for two functions, f_1 and f_2 , $f_1 = o(f_2)$ is equivalent to $\lim_{x \rightarrow a} \frac{|f_1(x)|}{|f_2(x)|} = 0$. From Eq. (7) we can define a diffusion tensor at \mathbf{x} by

$$\nu_x = \frac{1}{2} \mathbf{g}_x^{-1}. \quad (8)$$

The importance of the metric tensor comes from its direct connection with the error field. In dimension one, the metric is
 10 the scalar $g_x = \overline{(\partial_x \tilde{\varepsilon}_x)^2}$ where $\tilde{\varepsilon}$ denotes the normalized error field $\tilde{\varepsilon}_x = \frac{\varepsilon_x}{\sigma_x}$ (see appendix A). It is ~~meaningful~~ meaningful to relate the metric to a typical scale of correlation, the so-called error correlation length-scale (Daley, 1991; Pannekoucke et al., 2008)

$$L_x = 1/\sqrt{g_x}. \quad (9)$$

In dimension two (three), the metric is a 2×2 (3×3) matrix $\mathbf{g}_x = [g_{ij}(\mathbf{x})]$ given by

$$15 \quad g_{ij} = \overline{\partial_{x^i} \tilde{\varepsilon} \partial_{x^j} \tilde{\varepsilon}}. \quad (10)$$

Consequently, an approximation for the dynamics of the parametric formulation based on the diffusion equation is given by

$$\partial_t V = 2\overline{\varepsilon \partial_t \varepsilon}, \quad (11a)$$

$$\partial_t g_{ij} = \partial_t \left(\overline{\partial_{x^i} \tilde{\varepsilon} \partial_{x^j} \tilde{\varepsilon}} \right). \quad (11b)$$

Eq. Equation (11), has the advantage that we should be able to compute the time evolution of covariances for any error
 20 dynamics. This will be illustrated with Burgers ²-equation, that is a one dimensional dynamical model with nonlinear advection and diffusion processes similar to those of geophysical flows.

3 Dynamics of forecast error for Burgers ²-model

Here, we consider the dynamics associated with Burgers ²-equation

$$\partial_t u + u \partial_x u = \kappa \partial_x^2 u. \quad (12)$$

25 For any smooth function $u^0(x)$, there exists a unique solution $u(x, t)$ to Eq. (12) with the initial condition $u(x, 0) = u^0(x)$. A particular initial condition is now considered where $u^0(x)$ is a sample of a smooth random field of mean field

$\overline{u^0}(x)$. Hence, each sample u^0 is decomposed as $u^0(x) = \overline{u^0}(x) + \varepsilon^0(x)$, where ε^0 is a smooth random field. The dynamics of the mean $\overline{u}(x, t)$ and of higher order statistical moments are obtained from the Reynolds equations. Similarly to Cohn (1993), the ~~fluctuation-mean-flow~~ fluctuation-mean-flow dynamics deduced from the Reynolds equations is considered, in place of the more classical tangent-linear dynamics. Compared with the tangent-linear dynamics, ~~fluctuation-mean~~ fluctuation-mean-flow dynamics makes the mean flow depending on the fluctuation statistics evolution, which enlarge the tangent-linear setting. ~~The framework of the fluctuation-mean-flow dynamics is adapted to nonlinear Kalman filters similarly to the ensemble Kalman filter~~ Note that the fluctuation-mean-flow interaction leads to the Gaussian second-order filter (Jazwinski, 1970, sec. 9.3), and is important in nonlinear Kalman-like filters (Cohn, 1993). The next section presents the ~~fluctuation-mean-flow~~ fluctuation-mean-flow dynamics, and how it is used to describe the time evolution Eq. (11) of the two-points error covariance parameters.

3.1 Derivation of the ~~fluctuation-mean-flow~~ fluctuation-mean-flow dynamics for small error magnitude

The random field u can be decomposed into its ensemble-averaged and fluctuating parts $u = \overline{u} + \varepsilon$, where $\overline{u}(x, t) = \overline{u(x, t)}$ denotes the expectation of the random field u , and $\varepsilon = u - \overline{u}$ is a random field of zero mean. From this expansion, the mean flow dynamics is the ensemble ~~averaged-average~~ of the dynamics Eq. (12) and yields reads

$$\partial_t \overline{u} + \overline{u} \partial_x \overline{u} = \kappa \partial_x^2 \overline{u} - \overline{\varepsilon \partial_x \varepsilon}. \quad (13a)$$

The dynamics of the fluctuation ε is deduced from ~~te the~~ difference between the full dynamics Eq. (12) and the mean flow dynamics Eq. (13a), ~~leading to yielding~~ leading to yielding

$$\partial_t \varepsilon + \overline{u} \partial_x \varepsilon = -\varepsilon \partial_x \overline{u} + \overline{\varepsilon \partial_x \varepsilon} + \kappa \partial_x^2 \varepsilon - \varepsilon \partial_x \varepsilon. \quad (13b)$$

Hence, the dynamics of the mean flow and of the fluctuations are described by the coupled system Eq. (13).

Note that the term $-\overline{\varepsilon \partial_x \varepsilon}$ is the offset of the mean state due to the fluctuations. The offset term does not affect the statistical properties of the perturbations ε , while it is crucial to the dynamics of \overline{u} . From the commutativity of the ensemble mean with spatial derivative, $\overline{\partial_x(\cdot)} = \partial_x(\overline{\cdot})$, the offset term $\overline{\varepsilon \partial_x \varepsilon} = \frac{1}{2} \partial_x \overline{\varepsilon^2}$ can be written as a function of the variance $V = \overline{\varepsilon^2}$, $\overline{\varepsilon \partial_x \varepsilon} = \frac{1}{2} \partial_x V$.

If the magnitude of the perturbation ε is small, ~~Eqs Eq.~~ Eq. (13) can be simplified into the ~~fluctuation-mean-flow~~ fluctuation-mean-flow dynamics

$$\partial_t \overline{u} + \overline{u} \partial_x \overline{u} = \kappa \partial_x^2 \overline{u} - \frac{1}{2} \partial_x V, \quad (14a)$$

$$\partial_t \varepsilon + \overline{u} \partial_x \varepsilon = -\varepsilon \partial_x \overline{u} + \frac{1}{2} \partial_x V + \kappa \partial_x^2 \varepsilon, \quad (14b)$$

where the product $\varepsilon \partial_x \varepsilon$ has been discarded while keeping the ~~fluctuation-mean-flow~~ fluctuation-mean-flow interaction term $\frac{1}{2} \partial_x V = \overline{\varepsilon \partial_x \varepsilon}$. Note that ~~in the~~ tangent-linear dynamics, ~~the correspond to~~ Eq. (14) but where the offset term $\frac{1}{2} \partial_x V$ is discarded from Eq. Moreover, as pointed out in Ménard (1994), Eq. (14a) is the exact ensemble mean for the Burgers dynamics,

while Eq. (14b) is an approximate dynamics. As a consequence, if the variance field is the true one, then the mean predicted by Eq. (14a) is the true ensemble mean (Ménard, 1994, sec. 5.5.2).

The aim is now to determine the dynamics of the two-point error covariance function, $\overline{\varepsilon_x \varepsilon_y}$, that correspond, after spatial discretization, to the time evolution of the covariance matrix \mathbf{P} in data assimilation. Following the ~~time-splitting-splitting~~ strategy developed in Pannekoucke et al. (2016), the evolution of the perturbation ε is decomposed considering the effect of each process. The splitting strategy is a theoretical method to deduced the so-called infinitesimal generator of an evolution equation, by taking advantage of the Lie-Trotter formula to separate each processes (or appropriate arrangements of the processes). This strategy should not be confused with the numerical time-splitting which introduces numerical errors (Sportisse, 2007). Here, as seen in Eq. (14b), four processes influence the error statistics: (i) a production term due to the transport of the mean flow by the perturbation $-\varepsilon \partial_x \bar{u}$, (ii) the transport of the perturbation by the mean flow $\bar{u} \partial_x \varepsilon$, (iii) a diffusion term $\kappa \partial_x^2 \varepsilon$, and (iv) an offset term $\frac{1}{2} \partial_x V$ due to the averaged nonlinear self-interaction of the perturbation $\overline{\varepsilon \partial_x \varepsilon}$.

Since the offset (iv) modifies the mean but not the higher statistical moments of ε and without loss of generality, only the first three elementary processes are needed for the description of the covariance dynamics:

$$\partial_t \varepsilon = (-\partial_x \bar{u}) \varepsilon, \tag{15a}$$

$$15 \quad \partial_t \varepsilon = -\bar{u} \partial_x \varepsilon, \tag{15b}$$

$$\partial_t \varepsilon = \kappa \partial_x^2 \varepsilon. \tag{15c}$$

The effect of each process in Eq. (15) onto the dynamics Eq. (11) of the variance and the local diffusion tensor is now described.

3.2 Separate contribution of elementary processes

20 The contribution of the production term Eq. (15a) is first examined, then the transport Eq. (15b) and finally the diffusion Eq. (15c).

3.2.1 Contribution of the production term

The production term describes the amplification of the error due to the gradient of the mean field \bar{u} . This process can be viewed as a diagonal operator in the ~~Hilbert space where lies~~ function space where the random field ε lies. As a consequence, this error dynamics affects the variance but not the metric tensor. This leads to the following parameter dynamics

$$\partial_t V = -(2\partial_x \bar{u}) V, \tag{16a}$$

$$\partial_t \nu = 0. \tag{16b}$$

3.2.2 Contribution of the transport term

The time evolution of the variance and the diffusion fields due to the transport term Eq. (15b) is now tackled. Since the derivation is archetypal of how to proceed, the calculus are detailed.

The dynamics of error variance fields, deduced from Eq. (11a), yields

$$\partial_t V = 2\overline{\varepsilon(-\bar{u}\partial_x\varepsilon)} = -\bar{u}\partial_x\overline{\varepsilon^2}. \quad (17)$$

From the commutation of ensemble average and partial derivative, it simplifies to

$$\partial_t V = -\bar{u}\partial_x V. \quad (18)$$

5 Since $V = \sigma^2$, the dynamics of the standard deviation is given by

$$\partial_t \sigma = -\bar{u}\partial_x \sigma. \quad (19)$$

The dynamics of the metric tensor is deduced from Eq. (11b)

$$\begin{aligned} \partial_t g &= \partial_t \overline{(\partial_x \tilde{\varepsilon})^2}, \\ &= \overline{2\partial_{tx}^2 \tilde{\varepsilon} \partial_x \tilde{\varepsilon}} \\ 10 &= \overline{2\partial_x \left[\partial_t \left(\frac{\varepsilon}{\sigma} \right) \right] \partial_x \tilde{\varepsilon}} \\ &= \overline{2\partial_x \left[\frac{1}{\sigma} \partial_t \varepsilon - \frac{\varepsilon}{\sigma^2} \partial_t \sigma \right] \partial_x \tilde{\varepsilon}}. \end{aligned}$$

With the normalized error $\tilde{\varepsilon} = \frac{1}{\sigma}\varepsilon$ and the dynamics of the standard-deviation Eq. (19), the dynamics of the metric reads

$$\begin{aligned} \partial_t g &= \overline{2\partial_x \left[-\frac{\bar{u}}{\sigma} \partial_x (\sigma \tilde{\varepsilon}) + \tilde{\varepsilon} \bar{u} \partial_x \ln \sigma \right] \partial_x \tilde{\varepsilon}}, \\ &= -2\partial_x \bar{u} \overline{(\partial_x \tilde{\varepsilon})^2} - 2\bar{u} \partial_x \tilde{\varepsilon} \overline{\partial_x^2 \tilde{\varepsilon}}. \end{aligned}$$

15 From the identity $\partial_x(\partial_x \tilde{\varepsilon} \partial_x \tilde{\varepsilon}) = 2\partial_x \tilde{\varepsilon} \partial_x^2 \tilde{\varepsilon}$, and from $g = \overline{(\partial_x \tilde{\varepsilon})^2}$, ~~the last term of the right hand side is~~ we obtain

$$\partial_t g + \bar{u}\partial_x g = -2(\partial_x \bar{u})g. \quad (20)$$

Hence, the variance and the local diffusion $\nu = \frac{1}{2g}$ evolve following

$$\partial_t V + \bar{u}\partial_x V = 0, \quad (21a)$$

$$\partial_t \nu + \bar{u}\partial_x \nu = 2(\partial_x \bar{u})\nu. \quad (21b)$$

20 These equations represent the transport of the variance and of the diffusion by the mean flow. ~~Moreover:~~ the variance is conserved, while the diffusion tensor is warped by the mean flow.

3.2.3 Contribution of the diffusion term

Following the same procedure, the dynamics of V and g , Eq. (11), is given for the diffusion process Eq. (15c) by

$$\partial_t V = \kappa \left(\partial_x^2 V - \frac{1}{2} \frac{1}{V} \frac{1}{2V} (\partial_x V)^2 \right) - 2g\kappa V, \quad (22a)$$

$$\begin{aligned} 25 \quad \partial_t g &= 2\kappa g \frac{1}{V} \partial_x^2 V - 2g\kappa \frac{1}{V^2} (\partial_x V)^2 + \kappa \partial_x g \frac{1}{V} \partial_x V + \\ &\quad \kappa \partial_x^2 g + 2g^2 \kappa - 2\kappa \overline{(\partial_x \tilde{\varepsilon})^2}. \end{aligned} \quad (22b)$$

As it is expected while dealing with Reynold equations, a closure problem appears since the term $\overline{(\partial_x^2 \tilde{\varepsilon})^2}$ cannot be deduced from either V or g . Hence, a parameterization is needed to pursue.

To proceed further, we take advantage of the link between the unknown quantity $\overline{(\partial_x^2 \tilde{\varepsilon})^2}$ and the fourth order term K_x of the Taylor expansion of the error correlation function (see appendix A)

$$5 \quad \rho(x, x + \delta x) = 1 - \frac{1}{2} g_x \delta x^2 + S_x \delta x^3 + K_x \delta x^4 + o(\delta x^4), \quad (23)$$

where

$$g_x = \overline{\partial_x \tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x}, \quad (24a)$$

$$S_x = -\frac{1}{4} \partial_x g_x, \quad (24b)$$

$$K_x = \frac{1}{24} \overline{(\partial_x^2 \tilde{\varepsilon}_x)^2} - \frac{1}{12} \partial_x^2 g_x. \quad (24c)$$

10 The quantities S_x and K_x are later called the skewness and the kurtosis of the correlation function $\rho(x, \cdot)$. Note that, due to the symmetry of the two-points correlation functions, $\rho(x, y) = \rho(y, x)$, the skewness S_x is entirely determined by the metric field g . As a result, a choice of the kurtosis implies choosing the closure.

Two particular cases are interesting to discuss: when the random field is statistically homogeneous and, moreover, when the correlation function is a Gaussian function. In the case where the error random field is homogeneous, the error correlation function is homogeneous too: $\rho(x, y) = \rho(x + \delta, y + \delta), \forall \delta \in \mathbb{R}$. As a result, the fields of metrics, skewness and kurtosis are constant fields denoted by g^h, S^h and K^h . Due to the homogeneity of the metric field g^h , the skewness Eq. (24b) is zero, and the kurtosis Eq. (24c) is $K^h = \frac{1}{24} \overline{(\partial_x^2 \tilde{\varepsilon}_x)^2}$. In the case where the homogeneous correlation function is the Gaussian $\rho_G(x, x + \delta x) = e^{-\frac{1}{2} \frac{\delta x^2}{L_G^2}}$ where L_G stands for the homogeneous error correlation length-scale, the Taylor expansion reads

$$20 \quad \rho_{\underline{G}}(x, x + \delta x) = 1 - \frac{1}{2} \frac{\delta x^2}{L_G^2} + \frac{1}{2} \left(\frac{1}{2} \frac{\delta x^2}{L_G^2} \right)^2 \delta x^4 + o(\delta x^4), \quad (25)$$

and by identification with Eq. (23), $g_G = \frac{1}{L_G^2}, S_G = 0$ and $K_G = \frac{1}{8} g_G^2, g_G = \frac{1}{L_G^2}, S_G = 0$ and $K_G = \frac{1}{8} g_G^2$.

We propose to use these results to formulate a closure model: for a general smooth error random field of metric field g_x , the kurtosis K_x , Eq. (24c), is approximated by

$$K_x \underline{\sim GC} = \frac{1}{8} g_x^2 - \frac{1}{12} \partial_x^2 g_x, \quad (26)$$

25 where the first term of the right hand side is the kurtosis of the equivalent local homogeneous Gaussian correlation function. This closure is hereby called the *locally homogeneous Gaussian closure* or simply the *Gaussian closure*.

With this Gaussian closure Eq. (26), the dynamics of diffusion Eq. (22) is

$$\partial_t V = \kappa \left(\partial_x^2 V - \frac{1}{2} \frac{1}{V} (\partial_x V)^2 \right) - 2g\kappa V, \quad (27a)$$

$$\begin{aligned} \partial_t g = & 2\kappa g \frac{1}{V} \partial_x^2 V - 2\underline{g}\kappa\underline{g} \frac{1}{\underline{V}^2} (\partial_x V)^2 + \\ & \kappa \partial_x g \frac{1}{V} \partial_x V + \kappa \partial_x^2 g - 4g^2 \kappa. \end{aligned} \quad (27b)$$

- 5 In the one dimensional case, the dynamics of the local diffusion tensor are deduced from the dynamics of the metric from $\partial_t \nu = -2\nu^2 \partial_t g$ with $g = \frac{1}{2\nu}$. Thus, the variance and local diffusion tensor evolution are equivalently expressed as

$$\partial_t V = \kappa \left(\partial_x^2 V - \frac{1}{2} \frac{1}{V} \frac{1}{\underline{V}} (\partial_x V)^2 \right) - \frac{\kappa}{\nu} V, \quad (28a)$$

$$\begin{aligned} \partial_t \nu = & 2\kappa - 2 \frac{\kappa}{V} \frac{\kappa \nu}{\underline{V}} \partial_x^2 V \underline{\nu} + 2 \frac{\kappa}{V^2} \frac{\kappa \nu}{\underline{V}^2} (\partial_x V)^2 \underline{\nu} + \\ & \underline{2}\kappa \frac{1}{V} \partial_x V \partial_x \nu + \kappa \partial_x^2 \nu - 2\kappa \frac{1}{\nu} (\partial_x \nu)^2. \end{aligned} \quad (28b)$$

- 10 Contrary to the production Eq. (16) and the transport Eq. (21) processes, the effect of the diffusion process Eq. (28) creates a nonlinear coupling between the variance and the local diffusion field.

The parametric covariance dynamics for the Burgers equation is now ~~express~~expressed collecting all these results.

3.3 Parametric covariance dynamics of the Burgers equation

From Eq. (16), Eq. (21), and Eq. (28), the complete parametric covariance dynamics for the Burgers equation under the

- 15 Gaussian closure, is given by the coupled system

$$\partial_t \bar{u} + \bar{u} \partial_x \bar{u} = \kappa \partial_x^2 \bar{u} - \frac{1}{2} \partial_x V, \quad (29a)$$

$$\partial_t V + \bar{u} \partial_x V = -2(\partial_x \bar{u})V + \kappa \partial_x^2 V \underline{x} - \frac{\kappa}{2} \frac{1}{V_x} \frac{1}{\underline{V}} (\partial_x V \underline{x})^2 - \frac{\kappa}{V_x} \frac{\kappa}{\underline{V}} V \underline{x}, \quad (29b)$$

$$\begin{aligned} \partial_t \nu \underline{x} + \bar{u} \partial_x \nu \underline{x} = & 2(\partial_x \bar{u})\nu \underline{x} + 2\kappa - 2 \frac{\kappa}{V_x} \frac{\kappa \nu}{\underline{V}} \partial_x^2 V \underline{x} \nu \underline{x} + 2 \frac{\kappa}{V_x^2} \frac{\kappa \nu}{\underline{V}^2} (\partial_x V \underline{x})^2 \nu \underline{x} \\ & + \underline{2}\kappa \frac{1}{V_x} \frac{1}{\underline{V}} \partial_x V \underline{x} \partial_x \nu \underline{x} + \kappa \partial_x^2 \nu \underline{x} - 2\kappa \frac{1}{V_x} \frac{1}{\underline{V}} (\partial_x \nu \underline{x})^2. \end{aligned} \quad (29c)$$

- 20 Equation (29) exhibits a nonlinear coupling between the variance, Eq. (29b), and the local diffusion tensor, Eq. (29c), which illustrates the intricacy of the action of the diffusion process on the error dynamics. Moreover, Eq. (29) differs from its tangent-linear equivalent by the term $-\frac{1}{2} \partial_x V$ in Eq. (29a).

A numerical experiment is now proposed to illustrate and assess these theoretical results.




Figure 1. Nonlinear solution of the diffusive Burger' equation for the times $t = \{0, 0.2, 0.4, 0.6, 1\}T$ $t = \{0, 0.2, 0.4, 0.6, 0.8, 1\}T$.

4 Numerical experiment

A numerical experiment is proposed to illustrate the ability of the PKF forecast to reproduce the statistical evolution of the errors in the diffusive Burgers model. The numerical setting is first introduced, followed by an evaluation of the kurtosis closure. Then, the PKF is assessed using a large **nonlinear ensemble ensemble of nonlinear forecasts (6400 members)**. A sensitivity test on the different terms in the PKF concludes the section.

4.1 Numerical setting

For the numerical validation, a front-like situation is considered on a periodic domain of length $D = 1000$ km, discretized with $N = 241$ grid-points. The initial reference state, shown in Fig. 1, is the velocity field $\bar{u}^0(x) = U_{max} [1 + \cos(2\pi(x - D/4)/D)]/2$ with $U_{max} = 20$ km/h. From the nonlinear forecast of Eq. (12) starting from \bar{u}^0 , the maximum initially at 250 km, develops a front structure at 750 km after $T = 24$ hours of forecast. The diffusion coefficient κ is set so the front thickness $\delta \sim \frac{\kappa}{U_{max}}$, of order 50 km. The simulations considered here are integrated from the initial time $t = 0$ to the final time $t = T$. The Burgers equation Eq. (12) has been numerically integrating considering a classical finite difference spatial scheme and a fourth order Runge-Kutta time scheme with $\delta t = 0.002T$ and $\kappa = 0.0025D^2/T$.

The random perturbation at initial time, ε^0 , is set as a homogeneous random field of Gaussian distribution. The homogeneous covarianee Following Gaspari and Cohn (1999), the homogeneous correlation function is set as the Gaussian covarianee, in accordance with the geometry of the circle, as the chordal distance Gaussian correlation

$$P\rho^0(x, y) = (\sigma^0)^2 e^{-\frac{(x-y)^2}{2L_G^2} - \frac{d(x,y)^2}{2L_G^2}}, \quad (30)$$

where the homogeneous correlation length-scale is $L_G = 20$ km and where $d(x, y) = \frac{D}{\pi} |\sin \frac{\pi}{D}(x - y)|$ is the chordal distance between the two geographical positions x and y . Since the length-scale L_G is much smaller than the perimeter D , the Gaussian correlation Eq. (30) with the arc-length distance $d(x, y) = |x - y|$ is numerically very closed to the one with

the chordal distance (while at the theoretical level, the arc-length Gaussian is not strictly a correlation function on the circle, see Gaspari and Cohn (1999)), and leads to the same numerical results.

The covariance function is then defined as

$$P^0(x, y) = (\sigma^0)^2 \rho^0(x, y), \quad (31)$$

5 where σ^0 is the constant standard-deviation field. Thereafter, ~~two four~~ magnitudes of standard deviation σ^0 are considered: $\sigma_{1\%}^0 = U_{max}/100$ and $\sigma_{10\%}^0 = U_{max}/10$. Figure 2 illustrates the time evolution of two perturbed initial conditions whose perturbations are generated with the initial standard deviation $\sigma_{1\%}^0$ in panel (a) and $\sigma_{10\%}^0$ in panel (b) $\sigma_{1\%}^0 = 0.01 U_{max}$, $\sigma_{10\%}^0 = 0.1 U_{max}$, $\sigma_{20\%}^0 = 0.2 U_{max}$, and $\sigma_{50\%}^0 = 0.5 U_{max}$.

The time evolution of the true error covariance functions is computed considering a large ensemble of N_e nonlinear forecasts
 10 of Eq. (12). ~~Later the ensemble size is set to $N_e = 1600$ with $N_e = 6400$.~~ From the non-parametric convergence, the expected sampling error should thus represents about $1/\sqrt{N_e} = 0.25\%$ ~~$1/\sqrt{N_e} = 1.25\%$~~ of the real statistics. In order to limit the differences when comparing the results and due to the sampling noise, a single large ensemble of normalized error $(\tilde{\varepsilon}_k)_{k \in [1, N_e]}$ has been generated as $\tilde{\varepsilon}_k = \mathbf{C}^{1/2} \zeta_k$ where $\mathbf{C}^{1/2}$ is the square-root of the correlation matrix deduced from the correlations Eq. (30), and ζ_k is a sample of a random Gaussian noise of zero mean and covariance matrix the identity matrix. The ensemble
 15 of initial perturbation is then generated as $\varepsilon_k = \sigma^0 \tilde{\varepsilon}_k$ with $\sigma^0 \in \{\sigma_{1\%}^0, \sigma_{10\%}^0, \sigma_{20\%}^0, \sigma_{50\%}^0\}$.

Since the parametric covariance dynamics Eq. (29) has been theoretically derived for small perturbations, it has to be compared with the statistics from ~~an the~~ ensemble of small magnitude noise. Hence the validation is later conducted by considering ~~an the~~ ensemble generated from the initial standard deviation $\sigma_{1\%}^0$. Limits of predictability of the parametric covariance dynamics Eq. (29) are also addressed considering ~~an ensemble the ensembles~~ of larger initial uncertainty, ~~later set to from~~ $\sigma_{10\%}^0$
 20 ~~to $\sigma_{50\%}^0$.~~

Figure 2 illustrates the time evolution of the four perturbed initial conditions whose perturbations are generated from the normalized perturbation $\tilde{\varepsilon}_1$ scaled with the initial standard deviations from $\sigma_{1\%}^0$ in panel (a) to $\sigma_{50\%}^0$ in panel (d).

These ensembles are first used to tackle the closure of kurtosis, as discussed now.

4.2 Evaluation of the kurtosis closure

25 The aim of this section is to compare the kurtosis diagnosed from the true error covariance Eq. (24c), with the kurtosis resulting from the Gaussian closure Eq. (26). This validation is a crucial step since the quality of the closure will affect the skill of the parametric covariance dynamics Eq. (29). ~~If Even~~ though the closure is likely to be wrong for an arbitrary covariance matrix, it is expected to apply to most statistics encountered in applications. ~~Hence, the The~~ large ensemble, whose the initial perturbations are sampled by using $\sigma^0 = \sigma_{10\%}^0$, is considered for the validation. ~~The results are equivalent for the other error~~
 30 ~~magnitudes.~~ For this experiment, the error covariance matrices at times $t = 0$ and $t = T$ are representative of intermediate covariance matrices ; the computation of the true kurtosis and its closure is achieved by considering the ensemble at both times. The computation of the closure Eq. (26) relies on the local metric tensor, which has to be diagnosed from the ensemble.




Figure 2. Solutions from the initial perturbations of magnitude (a) $\sigma_{1\%}^0$ and (b) $\sigma_{10\%}^0$, (c) $\sigma_{20\%}^0$ and (d) $\sigma_{50\%}^0$, for the times $t = \{0, 0.2, 0.4, 0.6, 1\}T$ and $t = \{0, 0.2, 0.4, 0.6, 0.8, 1\}T$.

The local metric and kurtosis can be computed from the ensemble considering Eqs. (24a) and (24c). It is also possible to compute these quantities from the direct estimation of the local correlation function expansion Eq. (23), with the benefit of validating the theoretical derivations made in Eq. (24b) for the skewness and Eq. (24c) for the kurtosis. This motivates the estimation of these quantities from the computation of the local polynomial expansions, which are computed as follows.

- 5 For each position x , the fourth order polynomial approximation of the correlation function $\rho(x, \cdot)$ is estimated as the Lagrange interpolating polynomial $Q_x(X) = \sum_{k=-2}^2 \rho(x, x_k) \prod_{i \neq k} \frac{(X-x_i)}{(x_k-x_i)}$, computed from the five correlation values $(\rho(x, x_k))_{k \in [-2, 2]}$ where $x_k = x + k\delta x$. Expanding the Lagrange interpolating polynomial as $Q_x(X) = p_0 + p_1X + p_2X^2 + p_3X^3 + p_4X^4$, one obtains $g_x = -2p_2$ and $K_x = p_4$.

Figure 3 illustrates the results computed from the ensemble at $t = 0$ (panel a) and $t = T$ (panel b), showing in continuous line the length-scale $L_x = 1/\sqrt{g_x}$ (top panels) and the kurtosis (bottom panels), normalized by the initial homogeneous Gaussian




Figure 3. Diagnoses of the length-scale L_x normalized by $L_G L_G$ (top panels) and of the kurtosis (normalized by $K_G = \frac{1}{8}g_G^2$, $K_G = \frac{1}{8}g_G^2$) (bottom panels) deduced from the ensemble of forecast with initial uncertainty magnitude $\sigma_{10\%}^0$, at times $t = 0$ (a) and $t = T$ (b) (continuous line). The kurtosis is compared with the its closure $K_x^{GC} = \frac{1}{8}g_x^2 - \frac{1}{12}\partial_x^2 g_x$, $K_x^{GC} = \frac{1}{8}g_x^2 - \frac{1}{12}\partial_x^2 g_x$ (dashed line) computed from the metric field.

values L_G and $K_G = g_G^2/8$ with $g_G = 1/L_G^2 L_G$ and $K_G = g_G^2/8$ with $g_G = 1/L_G^2$. The kurtosis' closure Eq. (26) computed from the metric is in dashed line (bottom panels).

At $t = 0$, the length-scale (respectively the kurtosis) field is homogeneously equal to the initial values L_G ($K_G L_G$ (respectively K_G)). The small fluctuations visible at this time are due to the sampling noise. For $t = T$, the length-scale is larger than at the initial time, and presents an area of small values at in the vicinity of the front position $x = 0.75D$. The kurtosis is negligible, except at the vicinity of the front position, where the field is oscillating. For these two both times, it appears that the kurtosis' closure is able to reproduce the main behavior of the true kurtosis, with a low relative error $\|K - K^{GC}\|/\|K\|$ of 9.4% (17.6% $\|K - K^{GC}\|/\|K\|$ of 5.2% (respectively 21.5%) at time 0 (respectively T), where $\|\cdot\|$ stands for the L^2 norm. At time T , the maximum difference between the two normalized kurtosis, is 0.05.

Note that all the previous results are similar for the smaller initial uncertainty magnitude $\sigma_{1\%}^0$, with a relative error of 9.0% (14.3% 5.2% (respectively 14.5%) at time 0 (respectively T) (not shown here). For the larger error magnitude $\sigma_{20\%}^0$ and $\sigma_{50\%}^0$, the relative error at time T is respectively 40% and 63%. Hence, for this numerical simulation, the Gaussian closure proposed for the kurtosis appears relevant to approximate the real feature of the correlation shape. This is now used to explore the ability of the PKF to accurately predict the error statistics.

4.3 Theoretical Parametric versus ensemble length-scale and variance statistics

The theoretical-parametric setting is based on the online-time integration of the nonlinear coupled system Eq. Eq. (29) considering an equivalent numerical scheme than the one solving the Burgers equation Eq. (12), i.e. finite difference and RK4, with the same time step as detailed in Section 4.1. The numerical cost is of the order of a nonlinear time integration of the




Figure 4. Mean state at time $t = T$ computed from the ensemble of forecasts (solid line) and predicted from the parametric model Eq. (29) (dashed line), for the initial perturbations of magnitude (a) σ_1^0 , (b) σ_{10}^0 , (c) σ_{20}^0 and (d) σ_{50}^0 . The reference state is in grey solid line.

nonlinear Burgers equation. In this one-dimensional case, only two scalar fields are propagated: the variance V and the local diffusion field ν .

~~Error~~ The mean, the error variance and length-scale fields are reproduced in Fig. 4, Fig. 5 and Fig. 6, respectively, considering either a small or a large error at $t=0$ a range of initial errors. These figures compare the diagnosis from the ensemble with the analytical solution provided by of nonlinear forecasts of Eq. (12) with the statistics predicted by the parametric model Eq. (29). The mean diagnosed from the ensemble and predicted by the parametric model is first considered.

For the small

4.3.1 Comparison of the means

In order to appreciate the differences between the ensemble and the parametric means, the discussion is focused on the results at the final time T . When the initial error magnitude is small (Fig 4-(a)), corresponding to the tangent linear regime, the ensemble mean (continuous line) and the mean state predicted from the parametric model (dashed line) coincide with the reference state $u^0(T, x)$ (grey solid line, reproduced from Fig. 1). This is within the tangent linear validity regime where the small magnitude of the fluctuation has not impact on the ensemble mean which is then equal to the reference trajectory. For larger error magnitudes, the ensemble mean is expected to deviate from the reference trajectory due to the nonlinear interaction between the fluctuation and the mean. In the Burgers equation, the deviation is due to $-\frac{1}{2}\partial_x V$ in Eq. (29a) which implies here that the ensemble mean decreases as if the diffusion increased with the error magnitude (left panels-panels (b-d)). The mean predicted by the parametric model is very close to the ensemble mean (panel (b-c)) for the moderate error magnitudes of $\sigma_{10\%}$ and $\sigma_{20\%}$, but presents an anomalous distortion at the inflexion point for the larger error magnitudes (panel (d)). Hence, for the particular case of the Burgers dynamics, the parametric prediction of the mean is an accurate approximation of the ensemble mean, even for mild error magnitude.

4.3.2 Comparison of the variance and length-scale statistics

The variance (Fig. 5) and length-scale (Fig. 6) statistics are now discussed. For the small error magnitude, as seen in panels (a), the uncertainty is spreading at the initial time due to the physical diffusion, resulting in a strong dampening of the variance. This is accompanied by a global increase of the length-scale, except in the vicinity of the inflexion point located near $x = 0.5D$ (see Fig. 1). Then, as time goes on, the dynamical front generates a source of uncertainty where a beam of variance appears and increases with time, yielding a maximum of ~~8.8 of 10.0 times~~ the initial variance at $t = T$. The length-scale remains short close to the front position, except for a peak emerging from time $t = 0.6T$, evolving with the flow at the inflexion point. Compared-Comparing to the ensemble statistics, the PKF is able to capture all the details of the dynamics, ~~with a slightly smaller magnitude of variance at $t = T$~~ . This strongly support Eq. (29) as well as the underlying assumptions: the derivation of the tangent-linear dynamics for the error variance and length-scale fields, and the Gaussian closure for the kurtosis. In particular, the large length-scale values visible at the inflexion point of the front, is a real signal and not a numerical artifact of the diagnosis, since it is produced in both simulations.

The case of ~~large error magnitude is now considered~~the error magnitude of $\sigma_{10\%}$, where the tangent-linear approximation should no more be valid(~~right panels~~, is now considered (see panels (b)). Key features previously described are still present: emergence of a beam of uncertainty, increase of the length-scale except in the vicinity of the front. However, two differences appear compared to the ensemble statistics reference. Firstly, the magnitude of the uncertainty is lower than in the tangent-linear case; the maximum of variance beam at $t = T$ is now close to ~~5.5 of 6.0 times~~ the initial variance. Secondly, the local large length-scale value as depicted in the tangent-linear setting is nearly flat at the bottom of the small length-scale basin associated to the front. The main features of the PKF predictions are recovered: the variance beam has lower magnitude than in the tangent-linear case, and there is still a low length-scale area near the front.




Figure 5. Parametric (dashed line) versus ensemble estimated (continuous line) variance fields for initial perturbations of standard deviation magnitude $\sigma_1^0\%$ (a) (tangent-linear dynamics), $\sigma_{10}^0\%$ (b) (weakly nonlinear dynamics), $\sigma_{20}^0\%$ (c) and $\sigma_{50}^0\%$ (d). The fields are represented only for the times $t = \{0, 0.2, 0.4, 0.6, 0.8, 1\}T$.

Beyond the variance attenuation, a maximum at $t = T$ of ~~7.5~~ 7.8 times the initial variance, is much greater than the ensemble statistics result ~~showing an exponential growth, with a relative error of 29%~~. Moreover, the length-scale field displays a peak at the front, similar to the one described for the tangent-linear.

In order to assess the role of the nonlinear term, $\varepsilon \partial_x \varepsilon$, ~~on~~ in the error dynamics Eq. (13b), an evaluation with an ensemble has been performed. The results are displayed in Fig. 7, which shows, at $t = T$, the error variance and length-scale fields estimated with Eq. (29) (dashed line), compared to the fields diagnosed from a large ensemble (continuous line, and also shown in Fig. 5-(b) and 6-(b)). Then it shows the statistics computed from an ensemble of forecast of the tangent-linear dynamics Eq. (14) (small dashed line) (finite difference and RK4), and using the Reynolds equations Eq. (13) (dash-dotted line) (also finite difference and RK4). It appears that the statistics computed from the tangent-linear dynamics are equivalent to the error




Figure 6. Parametric (dashed line) versus ensemble estimated (continuous line) length-scale field for initial standard deviations $\sigma_{1\%}^0$ (a) (tangent-linear dynamics), $\sigma_{10\%}^0$ (b) (weakly nonlinear dynamics), $\sigma_{20\%}^0$ (c) and $\sigma_{50\%}^0$ (d). The fields are represented only for the times $t = \{0, 0.2, 0.4, 0.6, 0.8, 1\}T$.




Figure 7. Verification of the parametric variance (a) and correlation length-scale (b) at $t = T$ and for an initial perturbation standard deviation $\sigma_{10\%}^0$. The statistics from the ensemble of nonlinear forecasts of Eq. (12) are in black line, the statistics predicted by the parametric model are in blue dashed line, the statistics from the Reynolds equations Eq. (13) (respectively from the fluctuation-mean-flow dynamics Eq. (14)) with (respectively without) the second order term $\varepsilon \partial_x \varepsilon$ is in small red dashed-dotted line (respectively yellow dash-dotted line).

variance and length-scale fields predicted by the ~~theoretical-parametric~~ model, while the statistics from the Reynolds equations ~~are equivalent to the full nonlinear ensemble~~ equal those deduced from the ensemble of nonlinear forecasts of Eq. (12). Hence, the difference is well explained by the contribution of the nonlinear term $\varepsilon \partial_x \varepsilon$.

The case of the larger initial error magnitudes of $\sigma_{20\%}$ and $\sigma_{50\%}$ show similar results to the magnitude $\sigma_{10\%}$ case: the small length-scale area is captured by the PKF but with a spurious oscillation not present in the ensemble estimation (Fig. 6, panels c-d), the position of the beam of uncertainty is well predicted by the PKF but with a larger magnitude than the ensemble estimation (Fig. 5, panels c-d). Since the magnitude of the variance predicted by the PKF seems to increase faster than the ensemble estimation, it is interesting to investigate what happens for longer time window.

4.3.3 Long term behaviour

The increase of the PKF variance prediction might be a side effect due to the tangent linear like derivation of the PKF which could fails to predict the saturation of the error magnitude. In order to tackle the long term behaviour a comparison is conducted with a longer time window of $[0, 5T]$. Since the location of the uncertainty beam is well predicted by the PKF, the comparison focuses on the magnitude of the variance fields maximum. The time series of the variance maximum predicted by the PKF and estimated from the ensemble is shown in Fig. 8. The time evolution is equivalent for all the initial error magnitudes: after a short transition (where the variance decreases), two phases are seen, where the variance increases (phase 1), then saturates and decreases in long term (phase 2). The time where the maximum of variance is reached shifts with the magnitude of the initial error: it occurs after (respectively before) the time T for $\sigma_{1\%}$ (respectively $\sigma_{50\%}$). We associate the increase to the advection




Figure 8. Parametric (dashed line) versus numerical (continuous line) time series of the variance fields' maximum, from $t = 0$ to $t = 5T$, for initial perturbation of standard deviation magnitude $\sigma_{1\%}^0$ (a), $\sigma_{10\%}^0$ (b), $\sigma_{20\%}^0$ (c) and $\sigma_{50\%}^0$ (d).

contribution that involves the source term $-2(\partial_x \bar{u})V$ in the variance dynamics Eq. (29b), while the damping is related to the diffusion. Thanks to the competition between the advection and the diffusion, the increase of the variance first saturates then decreases. The PKF reproduces the two phases with a magnitude prediction closed but different from the ensemble estimation. For the small initial error magnitude $\sigma_{1\%}$ the PKF underestimates the variance for the long term behaviour, while the variance is overestimated for larger initial error magnitudes. From numerical investigations with smaller error magnitude at the initial time, the PKF prediction of the phase 2 appears more difficult than for the phase 1. This could be related to the choice of closure made for the kurtosis: while the locally homogeneous Gaussian closure is in accordance with the one diagnosed from the ensemble, an heterogeneous closure might improve the results. Beyond these deficiencies, it is interesting to retain that the theoretical derivation of the parametric model, which is partly based on the tangent linear assumption, is able to capture the main part of the uncertainty dynamics in the Burgers equation.

4.3.4 Discussion

From these results, we can conclude that the PKF forecast, ~~which implements as implemented by~~ Eq. (29), reproduces the tangent-linear evolution of the statistics as given by the covariance forecast equation Eq. (3). Put differently, the PKF forecast reproduces the tangent-linear covariance dynamics occurring in the extended Kalman filter. Since the PKF forecast model is deduced from ~~the tangent-linear dynamics a small error expansion~~, it is not meant to recover strongly nonlinear effects, which has been verified in the numerical experiment. Yet, even when the nonlinearity is stronger and the tangent-linear assumption is invalid, the solution of the PKF still shares some features with the empirical ensemble statistics. This may not be the case anymore for long-time integration of more complex geophysical dynamics. However, this suggests that some ~~part~~ of the statistics could be ~~accounted for predicted~~, at least for medium range forecast.

~~(a) (b) Analytical (dashed line) versus numerical (continuous line) variance fields for initial perturbation of standard deviation magnitude $\sigma_{1\%}^0$ (a) (tangent-linear dynamics) and $\sigma_{10\%}^0$ (b) (weakly nonlinear dynamics). The fields are represented only for the times $t = \{0, 0.2, 0.4, 0.6, 1\}T$.~~

~~(a) (b) Analytical (dashed line) versus numerical (continuous line) length-scale field for initial standard deviations $\sigma_{1\%}^0$ (a) — tangent-linear dynamics — and $\sigma_{10\%}^0$ (b) — weakly nonlinear dynamics. The fields are represented only for the times $t = \{0, 0.2, 0.4, 0.6, 1\}T$.~~

~~(a) (b) Verification of the theoretical variance (a) and correlation length-scale (b) at $t = T$ and for an initial perturbation standard deviation $\sigma_{10\%}^0$. The results for the nonlinear ensemble are in continuous line, the theoretical statistics are in dashed line, the ensemble without (with) second order term $\varepsilon \partial_x \varepsilon$ is in small dashed line (dash-dotted line), respectively.~~

5 Conclusions

This study focused on the forecast step of the parametric Kalman filter (PKF) applied to the nonlinear dynamics of the diffusive Burgers ²-equation. The parametric approach consists in approximating the error covariance matrix by a covariance model with evolving parameter fields. Here the covariance model considered is based on the diffusion equation, parameterized by the error variance and local diffusion fields. Hence, the forecast of the error covariance matrix, which is ~~practically impossible computationally very demanding~~ in real application with high-dimensional systems, amounts to the forecast of the error variance and local-diffusion fields, whose numerical cost is of the order of a single nonlinear forecast. In comparison, ensemble methods need dozens of members for the covariance forecast (which could be parallelized though), as well as localization to address the rank deficiency.

The derivation of the PKF dynamics has first been rigorously deduced from ~~the~~ dynamics of the perturbation under small magnitude assumption. However, a closure problem appears ~~,~~ due to the physical diffusion process. This closure issue has been related to the fourth term in the Taylor expansion of correlation function, the kurtosis, and a closure has been proposed based on an homogeneous Gaussian approximation for the kurtosis.

Numerical experiments where the true covariance evolution has been diagnosed from an ensemble ~~forecasting forecast~~ have been performed. A comparison with the PKF prediction has first shown the relevance of the closure, even for large error

magnitude. Moreover, [it has these experiments have](#) demonstrated the ability of the parametric formulation to reproduce the main features of the error dynamics when the tangent-linear approximation is valid. When tangent-linear dynamics is no more valid, the PKF could only reproduce part of the error statistics evolution, at least for mid-term forecast.

This contribution is a step toward the PKF formulation of more complex dynamics in geophysics. From the present study, we learned the difficulties of handling the higher order derivatives, since the coupling between the error variance and diffusion fields has been due to the physical diffusion. The Gaussian closure, similar to the one introduced in the kurtosis' treatment, will be useful to provide prognostic dynamics. But we expect that the main difficulties will be encountered in the forecast of multi-variate statistics that govern the balance between geophysical fields. Theoretically, the PKF formulation enables the forecast of covariance matrices in high dimension. Hence, it might offer new theoretical tools to approximate and to investigate important aspects of the dynamics of errors, such as the unstable subspace of chaotic dynamics. These points will be investigated in further developments.

Appendix A: Correlation fourth order derivative

The [aims aim](#) of this section is to show the following theorem (given here [in for](#) $d=1$).

Theorem: For smooth and centered error random field ε , with a smooth normalized error random field defined by $\tilde{\varepsilon}_x = \varepsilon_x / \sigma_x$, the error correlation function $\rho(x, y) = \overline{\tilde{\varepsilon}_x \tilde{\varepsilon}_y}$ locally expands as

$$\rho(x, x + \delta x) = 1 - \frac{1}{2}g_x \delta x^2 + S_x \delta x^3 + K_x \delta x^4 + o(\delta x^4), \quad (\text{A1})$$

with

$$g_x = \overline{\partial_x \tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x}, \quad (\text{A2a})$$

$$S_x = -\frac{1}{4} \partial_x g_x, \quad (\text{A2b})$$

$$K_x = \frac{1}{24} \overline{\partial_x^2 \tilde{\varepsilon}_x \partial_x^2 \tilde{\varepsilon}_x} - \frac{1}{12} \partial_x^2 g_x. \quad (\text{A2c})$$

Proof: For smooth normalized error field, the Taylor expansion, reads

$$\tilde{\varepsilon}_{x+\delta x} = \tilde{\varepsilon}_x + \partial_x \tilde{\varepsilon}_x \delta x + \frac{1}{2} \partial_x^2 \tilde{\varepsilon}_x \delta x^2 + \frac{1}{6} \partial_x^3 \tilde{\varepsilon}_x \delta x^3 + \frac{1}{24} \partial_x^4 \tilde{\varepsilon}_x \delta x^4 + o(\|\delta x\|^4). \quad (\text{A3})$$

Multiplication by $\tilde{\varepsilon}_x$ yields the correlation function

$$\rho(x, x + \delta x) = \overline{\tilde{\varepsilon}_x^2} + \overline{\tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x} \delta x + \frac{1}{2} \overline{\tilde{\varepsilon}_x \partial_x^2 \tilde{\varepsilon}_x} \delta x^2 + \frac{1}{6} \overline{\tilde{\varepsilon}_x \partial_x^3 \tilde{\varepsilon}_x} \delta x^3 + \frac{1}{24} \overline{\tilde{\varepsilon}_x \partial_x^4 \tilde{\varepsilon}_x} \delta x^4 + o(\delta x^4). \quad (\text{A4})$$

The aim is to reformulate terms of the form $\overline{\tilde{\varepsilon}_x \partial_x^k \tilde{\varepsilon}_x}$ in order to specify links between the coefficients.

The variance $\overline{\tilde{\varepsilon}_x^2}$ of the normalized error field is first studied. From linearity of the expectation operator it can be deduced that $\overline{\tilde{\varepsilon}_x^2} = \frac{1}{\sigma_x^2} \overline{\varepsilon_x^2} = 1$. Hence, the zero order term of the Taylor expansion Eq. (A4) is equal to 1.

The term $\overline{\tilde{\varepsilon} \partial_x \tilde{\varepsilon}}$ is now considered. Since $\partial_x \tilde{\varepsilon}_x^2 = 2\tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x$, the commutation rule $\partial_{x^i}(\cdot) = \overline{\partial_{x^i}(\cdot)}$ and the stationarity of the

$$\overline{\tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x} = 0, \quad (\text{A5})$$

so that the first order term of the Taylor expansion Eq. (A4) is zero.

For the term $\overline{\tilde{\varepsilon}_x \partial_x^2 \tilde{\varepsilon}_x}$: by using the identity $\partial_x(\tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x) = \partial_x \tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x + \tilde{\varepsilon}_x \partial_x^2 \tilde{\varepsilon}_x$, and the commutation rule, [it results we have](#) that $\overline{\tilde{\varepsilon}_x \partial_x^2 \tilde{\varepsilon}_x} = \partial_x(\overline{\tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x}) - \overline{\partial_x \tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x}$. With, Eq. (A5), this simplifies into

$$15 \quad \overline{\tilde{\varepsilon}_x \partial_x^2 \tilde{\varepsilon}_x} = -\overline{\partial_x \tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x}. \quad (\text{A6})$$

Identifying with Eq. (A1), the second order term of the Taylor expansion Eq. (A4) can be written as

$$g_x = \overline{\partial_x \tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x}. \quad (\text{A7})$$

The third [order](#) term $\overline{\tilde{\varepsilon}_x \partial_x^3 \tilde{\varepsilon}_x}$ is reformulated as follows: $\partial_x(\tilde{\varepsilon}_x \partial_x^2 \tilde{\varepsilon}_x) = \partial_x \tilde{\varepsilon}_x \partial_x^2 \tilde{\varepsilon}_x + \tilde{\varepsilon}_x \partial_x^3 \tilde{\varepsilon}_x$ implies $\overline{\tilde{\varepsilon}_x \partial_x^3 \tilde{\varepsilon}_x} = \partial_x(\overline{\tilde{\varepsilon}_x \partial_x^2 \tilde{\varepsilon}_x}) - \overline{\partial_x \tilde{\varepsilon}_x \partial_x^2 \tilde{\varepsilon}_x}$, which is transformed using Eq. (A6) into $\overline{\tilde{\varepsilon}_x \partial_x^3 \tilde{\varepsilon}_x} = -\partial_x(\overline{\partial_x \tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x}) - \overline{\partial_x \tilde{\varepsilon}_x \partial_x^2 \tilde{\varepsilon}_x}$. But with $\partial_x(\partial_x \tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x) =$

15 $2\partial_x \tilde{\varepsilon}_x \partial_x^2 \tilde{\varepsilon}_x$, it results that

$$\overline{\partial_x \tilde{\varepsilon}_x \partial_x^2 \tilde{\varepsilon}_x} = \frac{1}{2} \partial_x(\overline{\partial_x \tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x}), \quad (\text{A8})$$

and then

$$\overline{\tilde{\varepsilon}_x \partial_x^3 \tilde{\varepsilon}_x} = -\frac{3}{2} \partial_x(\overline{\partial_x \tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x}). \quad (\text{A9})$$

Identifying with Eq. (A1), the third order term of the Taylor expansion Eq. (A4) can be written as

$$20 \quad S_x = -\frac{1}{4} \partial_x g_x. \quad (\text{A10})$$

For the last fourth [order](#) term $\overline{\tilde{\varepsilon}_x \partial_x^4 \tilde{\varepsilon}_x}$, the computation of the derivative $\partial_x(\tilde{\varepsilon}_x \partial_x^3 \tilde{\varepsilon}_x) = \partial_x \tilde{\varepsilon}_x \partial_x^3 \tilde{\varepsilon}_x + \tilde{\varepsilon}_x \partial_x^4 \tilde{\varepsilon}_x$, leads to $\overline{\tilde{\varepsilon}_x \partial_x^4 \tilde{\varepsilon}_x} = \partial_x(\overline{\tilde{\varepsilon}_x \partial_x^3 \tilde{\varepsilon}_x}) - \overline{\partial_x \tilde{\varepsilon}_x \partial_x^3 \tilde{\varepsilon}_x}$. From Eq. (A9), this reads $\overline{\tilde{\varepsilon}_x \partial_x^4 \tilde{\varepsilon}_x} = -\frac{3}{2} \partial_x^2(\overline{\partial_x \tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x}) - \overline{\partial_x \tilde{\varepsilon}_x \partial_x^3 \tilde{\varepsilon}_x}$. The last term can be deduced from the use of the homogeneity of the normalized error variance $\partial_x^4 \overline{\tilde{\varepsilon}_x^2} = 0$. Since $\partial_x^4 \tilde{\varepsilon}_x^2 = \partial_x^3[2\tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x]$, the Leibniz rule states

$$25 \quad \partial_x^4 \tilde{\varepsilon}_x^2 = 2(\partial_x^3 \tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x + 3\partial_x^2 \tilde{\varepsilon}_x \partial_x^2 \tilde{\varepsilon}_x + 3\partial_x \tilde{\varepsilon}_x \partial_x^3 \tilde{\varepsilon}_x + \tilde{\varepsilon}_x \partial_x^4 \tilde{\varepsilon}_x). \quad (\text{A11})$$

With the ensemble average, it follows that $\overline{\partial_x \tilde{\varepsilon}_x \partial_x^3 \tilde{\varepsilon}_x} = -\frac{1}{4} \overline{\tilde{\varepsilon}_x \partial_x^4 \tilde{\varepsilon}_x} - \frac{3}{4} \overline{\partial_x^2 \tilde{\varepsilon}_x \partial_x^2 \tilde{\varepsilon}_x}$. Hence,

$$\overline{\tilde{\varepsilon}_x \partial_x^4 \tilde{\varepsilon}_x} = -2\partial_x^2(\overline{\partial_x \tilde{\varepsilon}_x \partial_x \tilde{\varepsilon}_x}) + \overline{\partial_x^2 \tilde{\varepsilon}_x \partial_x^2 \tilde{\varepsilon}_x}. \quad (\text{A12})$$

Identifying with Eq. (A1), the [third-fourth](#) order term of the Taylor expansion Eq. (A4) can be written as

$$K_x = \frac{1}{24} \overline{\partial_x^2 \tilde{\varepsilon}_x \partial_x^2 \tilde{\varepsilon}_x} - \frac{1}{12} \partial_x^2 g_x. \quad (\text{A13})$$

Author contributions. TEXT

Competing interests. TEXT

Disclaimer. TEXT

Acknowledgements. This work was supported by the french national program LEFE/INSU ([Étude du filtre de Kalman Paramétrique, KAPA](#)). The authors are thankful to [S. E. Cohn](#) and an anonymous reviewer for their useful comments, suggestions and corrections. CERA is a member of Institut Pierre-Simon Laplace (IPSL).



References

- Apte, M., Auroux, D., and Ramaswamy, M.: Variational data assimilation for discrete Burgers equation, *Electronic Journal of Differential Equations Conf.*, 19, 15–30, 2010.
- Bocquet, M.: Localization and the iterative ensemble Kalman smoother, *Quarterly Journal of the Royal Meteorological Society*, 142, 1075–1089, doi:10.1002/qj.2711, 2016.
- Bouttier, F.: The dynamics of error covariances in a barotropic model, *Tellus A*, 45, 408–423, 1993.
- Burgers, J.: *The nonlinear diffusion equation: asymptotic solutions and statistical problems*, D. Reidel Publishing Company, 1974.
- Cohn, S.: Dynamics of Short-Term Univariate Forecast Error Covariances, *Monthly Weather Review*, 121, 3123–3149, 1993.
- Daley, R.: *Atmospheric Data Analysis*, Cambridge University Press, 1991.
- 10 Desroziers, G., Camino, J.-T., and Berre, L.: 4DEnVar: link with 4D state formulation of variational assimilation and different possible implementations, *Quarterly Journal of the Royal Meteorological Society*, doi:10.1002/qj.2325, 2014.
- Desroziers, G., Arbogast, E., and Berre, L.: Improving spatial localization in 4DEnVar, *Quarterly Journal of the Royal Meteorological Society*, 142, 3171–3185, doi:10.1002/qj.2898, 2016.
- el Malek, M. B. A. and El-Mansi, S. M.: Group theoretic methods applied to Burgers' equation, *Journal of Computational and Applied Mathematics*, 115, 1–12, doi:10.1016/s0377-0427(99)00170-3, 2000.
- 15 Evensen, G.: Sequential data assimilation with a nonlinear quasi-geostrophic model using Monte Carlo methods to forecast error., *J. Geophys. Res.*, 99, 10 143–10 162, 1994.
- Gaspari, G. and Cohn, S. E.: Construction of correlation functions in two and three dimensions, *Quarterly Journal of the Royal Meteorological Society*, 125, 723–757, 1999.
- 20 Hopf, E.: The partial differential equation $u_t + u u_x = u u_{xx}$, *Communications on Pure and Applied Mathematics*, 3, 201–230, doi:10.1002/cpa.3160030302, 1950.
- Jazwinski, A.: *Stochastic Processes and Filtering Theory*, Academic Press, 1970.
- Lakshminarayanan, S., Lewis, J. M., and Phan, D.: Data Assimilation as a Problem in Optimal Tracking: Application of Pontryagin's Minimum Principle to Atmospheric Science, *Journal of the Atmospheric Sciences*, 70, 1257–1277, doi:10.1175/jas-d-12-0217.1, 2013.
- 25 Ménard, R.: Kalman filtering of Burgers' equation and its application to atmospheric data assimilation., Ph.D. thesis, Ph.D. Thesis. McGill University, 1994.
- Ménard, R., Cohn, S., Chang, L.-P., and Lyster, P. M.: Assimilation of Stratospheric Chemical Tracer Observations Using a Kalman Filter. Part I: Formulation, *Monthly Weather Review*, 128, 2654–2671, 2000.
- Mirouze, I. and Weaver, A. T.: Representation of correlation functions in variational assimilation using an implicit diffusion operator, *Quarterly Journal of the Royal Meteorological Society*, 136, 1421–1443, 2010.
- 30 Pannekoucke, O. and Massart, S.: Estimation of the local diffusion tensor and normalization for heterogeneous correlation modelling using a diffusion equation., *Q. J. R. Meteorol. Soc.*, 134, 1425–1438, 2008.
- Pannekoucke, O., Berre, L., and Desroziers, G.: Background error correlation length-scale estimates and their sampling statistics, *Quarterly Journal of the Royal Meteorological Society*, 134, 497–508, 2008.
- 35 Pannekoucke, O., Ricci, S., Barthelemy, S., Ménard, R., and Thual, O.: Parametric Kalman Filter for chemical transport model, *Tellus*, 68, 31 547, 2016.

- Snyder, C., Hamill, T. M., and Trier, S. B.: Linear Evolution of Error Covariances in a Quasigeostrophic Model, *Monthly Weather Review*, 131, 189–205, 2003.
- Sportisse, B.: A review of current issues in air pollution modeling and simulation, *J. Comp. Geosci.*, 11, 159–181, 2007.
- Verlaan, M. and Heemink, A. W.: Nonlinearity in Data Assimilation Applications: A Practical Method for Analysis, *Monthly Weather Review*, 129, 1578–1589, doi:10.1175/1520-0493(2001)129<1578:nidaaa>2.0.co;2, 2001.
- 5 Weaver, A. and Courtier, P.: Correlation modelling on the sphere using a generalized diffusion equation (Tech. Memo. ECMWF, num. 306), *Quart. J. Roy. Meteor. Soc.*, 127, 1815–1846, 2001.
- Weaver, A. T. and Mirouze, I.: On the diffusion equation and its application to isotropic and anisotropic correlation modelling in variational assimilation, *Quarterly Journal of the Royal Meteorological Society*, 139, 242–260, 2013.
- 10 Zupanski, M., Navon, I. M., and Zupanski, D.: The Maximum Likelihood Ensemble Filter as a non-differentiable minimization algorithm, *Quarterly Journal of the Royal Meteorological Society*, 134, 1039–1050, doi:10.1002/qj.251, 2008.