

1 Introduction

Further to the reviews of this submission, we have considered all of the comments made by both reviewers and Editor and taken them fully into account in revising our manuscript. We have accepted most of them and respond to each comment as follows. In addition, we include a version of the revised manuscript with significant changes highlighted in blue.

2 Reviewer 1

We are grateful to the referee for these comments and suggestions. We have implemented most of them in the text and figures, as indicated below, with changes to the text noted in blue in the new version of the manuscript.

- (1) *Fig 7. from Young (2014) is incorrect for FZ: 2.08 to be replaced by 1.69*
Now done

- (2) *Fig.8. Please define the Peclet number Pe ($Nu-1$?)*
Péclet number is indeed defined as $Nu-1$ (advective/conductive transport). This is now defined in the text.

- (3) *3.1 and Fig.9. Saturn and Jupiter do have super-rotation in the equator (and multiple jets, Note that Saturn "S" is missing in Fig.9) in contradiction with Fig.9. This part seems to suggest that only large Rossby numbers yield super-rotation: this is not what is observed. It might be the case that the authors mean "global (?)" super-rotation and arguing that one is within the PUMA realm but a comment on this would be very welcome.*

Global super-rotation was intended in this discussion and a comment has been added to clarify this (with a reference to the review by Read & Lebonnois 2018). We also note that Jupiter and Saturn have locally super-rotating equatorial jets, although the global super-rotation is not known for either planet. The reference (S) to Saturn in Fig. 9 has been added.

- (4) *Fig 12. The arrow thicknesses must be consistent with their numerical values. The convention of using negative values is also misleading. Why not use positive values provided the arrows are in the right direction ?*

Noted. We have now changed the convention to use positive numbers everywhere and have adjusted the thickness of arrows to be clearer.

- (5) *12(a) NCEP data are used here: plz correct -0.48 to 0.48 since the arrow direction is in the opposite direction. Note that one should look at Boer and Lambert instead of Fig. 1 of Tabataba-Vakili et al. (pls correct text as well).*

Number changed to positive in Fig. 12(a) and references are now entirely to Boer & Lambert for these numbers.

- (6) 12(b) pls refer explicitly to Fig.12 (b) in 4.1.2

Now added.

- (7) 12(c) correct to positive values like in Lee and Richardson.

Done.

3 Reviewer 2

This is an interesting and generally well-written review of barotropic and baroclinic instabilities, and their equilibration, in planetary atmospheres. I have mostly fairly minor comments and clarifications for the authors to consider.

Thanks for this. We respond to these points below and have modified the text of the paper as indicated.

- (1) *I think it was Margules who introduced the idea of a potential energy that is available for conversion to kinetic energy. Granted it was Lorenz who (as the authors describe) came up with the eponymous energy cycle, but in a review article such as this it might be appropriate to reference Margules.*

Noted. We have added a reference to Margules (1903) as suggested.

- (2) *Regarding baroclinic equilibration and baroclinic adjustment. It is fairly apparent that Earth's ocean is in a supercritical state, not one of marginal criticality. The ocean may be beyond the scope of this article, but that observation prompted a somewhat more general discussion of why supercriticality does occur that may be found in Jansen and Ferrari in 2012/2013. Going back a little further, precursors may be found in Salmon (GAFD 1980) and Vallis (QJ 1988) who both found supercritical regimes. The authors may wish to consider a brief discussion of all this.*

This is a good point. Although the ocean is probably beyond the intended scope of the article, similar super-critical regimes may occur in gas giant atmospheres. We have added some extra discussion of this in Section 5, including mention of some of the scaling arguments of Jansen and Ferrari. A notable addition to this from the present work is to emphasise the non-negligible role of the zonally-symmetric meridional overturning in addition to eddy transports in maintaining a criticality close to 1 under some conditions. This is now clarified in our discussion.

- (3) *Regarding Venus, it seems to me rather an 'ill-posed planet' as regards quasigeostrophy, or at least its troposphere is. Both N and f are close to zero (compared to values on Earth) so the deformation radius, NH/f (or equivalently the square root of the Burger number) is poorly defined. At cloud level N is finite but f is still small, so the conventional deformation radius is very large, probably larger than the planet itself. The discussion toward the end of section 4.2 needs a little more clarification, since as*

defined by equation (13) the Burger number does not vary with the reference frame, unless f is being considered as changing, which is perhaps what the authors mean. But in any case this needs clarifying and a bit more discussion.

Noted. We have clarified in our discussion in Section 4.2 that the modified Burger number in the cloud layers assumes a change in f to a value representative of the mean angular velocity of the cloud layer itself - with a rotation period of around 4 Earth days. This is what has been assumed in other published work, such as that by Young et al. (1984) and Sugimoto et al. (2014).

- (4) *It is not just realistically forced models that struggle to get the key features of Venus. Different models with Held-Suarez forcing can give very different results, as I think some recent model intercomparisons have shown (although I don't have a reference to hand). I suspect this is at least in part because of the ability of models to conserve angular momentum, and the fact that models are sensitive to the ratio of N/f , which might vary among models.*

Agreed. We have included some additional discussion of idealised models of the Venus and Titan atmospheres and now reference the recent inter-comparisons of such models by Lebonnois and Lee & Richardson.

- (5) *The authors say, early in in the conclusions, that 'the distinction between baroclinic and barotropic instabilities is to a large extent an artificial one.' Is that really the case? Granted the CSP criterion covers both cases, but as the authors themselves say the difference is associated with whether the basic state has a store of potential energy that is converted to eddy kinetic energy, and this doesn't seem like a minor difference. And the Lorenz cycles for barotropic and baroclinic instabilities are quite different. If the authors are trying to make a profound statement that, in spite of these differences, the instabilities are really of the same type then I think they need to justify this more. If not, I'd suggest they moderate that statement.*

Point taken. The similarities are from a vorticity point of view only, but the distinction remains energetically as noted here. We have clarified this in Section 5.

- (6) *At the beginning of the section on Jupiter the wording suggests that there are a number of gas giants in the Solar System, whereas in fact there are only two, if ice giants are regarded as separate. This is made clearer later on, but a slight rephrasing to clarify might help.*

OK though the dynamical distinction between gas and ice giants is perhaps rather a fine and subtle one. Section 4.3 does make a distinction between gas and ice giants in a few places, but we have added emphasis to the distinction in our discussion of Jupiter.

- (7) *Is it really the case that (line 581) 'the release of latent heat energy from condensation of water vapour plays an important role ...in the weather layer'? Do we know that or is that really just a conjecture? After all, the moist and dry adiabatic lapse rates are almost the same on Jupiter because of the hydrogen atmosphere, suggesting that moisture may have a limited effect. Granted there are arguments the other way (lightning is seen, there is a strong virtual temperature effect, some numerical simulations), but the importance otherwise of moisture seems to me an open question. If I am mistaken and it is a settled issue then the authors need to point to some definitive evidence and give references.*

Precisely what the role of latent heat release in moist convection is in the weather layers of Jupiter and Saturn is still an open one. But evidence for the occurrence of moist convection I think is fairly unequivocal (notably through observations of lightning and convective water clouds). Also, estimates of upward energy flux by moist convection (e.g. by Gierasch et al. (2000)) suggest that an $O(1)$ fraction of the interior heat on Jupiter ($> 50\%$) may be carried by moist convective processes. This led Ingersoll et al. (2000) to suggest that the dominant eddies driving the zonal jets might also be from moist convection, although the recent work of Young and Read (2017) suggests that baroclinic instabilities (on scales similar to the Rossby deformation radius) may be most important for this. We have added some discussion in Section 4.3 to clarify this.

4 Editor

Thank you for these comments and your detailed and careful reading of this manuscript. We will take these into account in producing a revised version of the paper, which we hope will then be acceptable. Our detailed responses follow.

- *The line numbers below are those of the version I have had access to, once printed. For some reason, the numbers seem to be shifted by one or two units from the ones on the version I visualize on my screen.*

I have noticed this too! But I think most of your references are clear.

- (1) *1. Eq. (1). An integral sign (from 0 to H I presume) is missing for the integral with respect to z on the left-hand side of the equation. And it might be useful to specify at this stage that y is the latitudinal coordinate and z the vertical coordinate.*

Well spotted! Yes, the integral sign has been added. We also specify the coordinates here as suggested.

- (2) *There seems to be an inconsistency as concerns the values of the thermal Rossby number Ro_T . The text (starting l. 388) says that the eddy meridional heat transport peaks at $Ro_T \sim 0.07$, while Fig. 11(a) shows a peak at $Ro_T \sim 0.3$ (see also Table 1, and ll. 724–730 and 748–751).*

A straightforward error – now corrected to $Ro_T \sim 0.3$ for the peak.

- (3) *Fig. 3(c). There are two curves on the figure. What is the difference between them ?*

This is just a single curve that is double valued, and comes from plotting u vs dQ/dy point wise across the domain represented in the radial profiles in 3(a) and (b). u takes different values at different radii, despite have the same value of dQ/dy . This is presumably because the flow is not precisely at the inviscid marginally stable state, though we don't think it is necessary to labour this point in the text.

- (4) *Ll. 444-445, ... leads to a value of \mathcal{T}_R which is much larger than $O(1)$ but not hugely so Well, the value given in Table 2 is 1.3×10^5 .*

OK point taken. We have omitted the “but not hugely so” phrase.

- (5) *Ll. 710-711, ... to infer the existence of a unique reference frame on each planet ... A unique reference frame with which properties ?*

This is the unique state at which the gravest Doppler-shifted Rossby wave trains are just able to propagate at the same phase speed and hence couple together and interact to grow via over-reflection. We have added text in Section 5.2 to clarify this.

- (6) *Ll. 461-462, ... the observed near suppression of baroclinic instability in Martian summers It would be better to give appropriate reference(s).*

OK. A reference to the paper by Lewis et al. (2016) has been inserted here.

- (7) *Fig. 6. What is the precise connection between the vertical coordinate (Stability parameter) and the thermal Rossby number ?*

OK - they are the same in this diagram (a consequence of using a figure from another source). We have clarified this in the caption.

- (8) *Fig. 8. Inset. It would be preferable to say explicitly that that Pe refers to heat transport by the axisymmetric flow, Pe_s to transport by the eddies, and m to the number of longitudinal waves.*

This has been clarified in the caption.

- (9) *L. 100, ... for comparison, in Section 4, with the known properties ...*

Well spotted! Reference to Section 4 has been added.

- (10) *Eq. (15) Inconsistency of notation. θ or θ with overbar ?*

OK, overbar is intended here and has been added.

- (11) *Ll. 418-419, ... the values of Bu , Ro_T and \mathcal{T}_R [...], based on Eqs (12-14)*

Noted and corrected.

- (12) *Ll. 780-781. Contrary to what the text implies, Table 2 does not mention values for Saturn.*

Reference to Saturn has been deleted.

- (13) *Fig. 10, caption, and l. 382. What is PUMA-S with respect to PUMA, introduced earlier ?*

This was used in Wang et al. (2018) to mean the Held-Suarez simplified version of the PUMA model. But this distinction is unnecessary in the present context so the “-S” suffix has been omitted throughout the paper.

- (14) *It would be preferable to define the Burger number when it is first introduced (l. 241) rather than later on (Eq. 13).*

OK, the definition of Bu has been added here (as well as the modified definition in Eq (14) in the revised version (line 350).

- (15) *L. 125. Say that u^* and v^* are perturbations with respect to zonal mean, and that the overbar denotes a longitudinal mean.*

These definitions have been added.

- (16) *Table 1 does not seem to be referenced in the text. It could be on l. 363, after mention of the range of variation of Ω^* .*

Noted - thanks for the suggestion. Now added.

- (17) *L. 350, $\Delta\theta_{EP} \rightarrow \Delta\theta_E P$*

Noted - now corrected.

- (18) *Caption of Table 2. Expand PDS (Planetary Data System)*

Now added.

- (19) *L. 64, ... quasi-geostrophic potential vorticity (QGPV), ...*

Now added.

- (20) *L. 468, expand LMD (Laboratoire de Météorologie Dynamique)*

Now added.

- (21) *L. 451, ... tilt with altitude.*

We have clarified this by “...pronounced latitudinal tilt with altitude”.

- (22) *L. 141, ... yet it is observed ...*

Now changed.

- (23) *L. 114, ... stability criterion (i) (parentheses, a similar correction is to be made in other places, please check).*

Well spotted. Now corrected in this and a few other places.

BAROCLINIC AND BAROTROPIC INSTABILITIES IN PLANETARY ATMOSPHERES - ENERGETICS, EQUILIBRATION AND ADJUSTMENT

Peter Read¹, Neil Lewis¹, Daniel Kennedy¹, H       Scolan^{2,1}, Fachreddin Tabataba-Vakili^{3,1}, Yixiong Wang¹, Susie Wright¹, and Roland Young^{4,1}

¹Department of Physics, University of Oxford, Clarendon Laboratory, Parks Road, Oxford, OX1 3PU, UK

²Laboratoire de Mécanique des Fluides et d'Acoustique, Université Lyon, France

³Jet Propulsion Laboratory, Pasadena, California, USA

⁴Department of Physics & National Space Science and Technology Center, UAE University, Al Ain, United Arab Emirates

Correspondence: Peter Read (peter.read@physics.ox.ac.uk)

Abstract. Baroclinic and barotropic instabilities, are well known as the mechanisms responsible for the production of the dominant energy-containing eddies in the atmospheres of the Earth and several other planets, as well as the Earth's oceans. Here we consider insights provided by both linear and nonlinear instability theories into the conditions under which such instabilities may occur, with reference to forced and dissipative flows obtainable in the laboratory, in simplified numerical atmospheric circulation models and in the planets of our Solar System. The equilibration of such instabilities is also of great importance in understanding the structure and energetics of the observable circulation of atmospheres and oceans. Various ideas have been proposed concerning the ways in which baroclinic and barotropic instabilities grow to large amplitude and saturate, whilst also modifying their background flow and environment. This remains an area that continues to challenge theoreticians and observers, though some progress has been made. The notion that such instabilities may act under some conditions to adjust the background flow towards a critical state is explored here in the context of both laboratory systems and planetary atmospheres. Evidence for such adjustment processes is found relating to baroclinic instabilities under a range of conditions where the efficiency of eddy and zonal mean heat transport may mutually compensate to maintain a nearly invariant thermal structure in the zonal mean. In other systems, barotropic instabilities may efficiently mix potential vorticity to result in a flow configuration that is found to approach a marginally unstable state with respect to Arnol'd's second stability theorem. We discuss the implications of these findings and identify some outstanding open questions.

1 Introduction

One of the great achievements of the past 100 years in fluid dynamics has been the development of a theory of dynamical instability, both linear and nonlinear. In a geophysical context, this has led to a quantitative understanding of a variety of phenomena, including the processes that lead to the development of large-scale energetic eddies in rotating, stratified atmospheres and oceans. These instabilities, known as baroclinic and barotropic instabilities, are well known as the mechanisms responsible for the production of the dominant energy-containing eddies in the atmospheres of the Earth and several other planets, as well

as the Earth's oceans. In general they occur in flows for which background rotation plays an important role, while baroclinic instabilities also require statically stable stratification. They are typically distinguished energetically by the respective dominance of exchanges of either kinetic or potential energy with the background (usually zonal) flow. Barotropic instability is associated with direct exchanges of kinetic energy between background flow and eddies, while baroclinic instability entails the growth of eddy kinetic and potential energy at the primary expense of the potential energy of the background flow.




Figure 1. (a) Schematic representation of energy storage and interconversions in the form suggested by Lorenz (1967), with each box representing the energy reservoirs for eddy and zonal mean kinetic (KE,KZ) and available potential energies (AE,AZ) and arrows representing conversion rates between them (CZ,CK,CE,CA). Sources or sinks of potential or kinetic energy are also shown as external arrows (GZ,GE,FZ,FE). (b)-(d) are idealised examples representing the cycles expected for isolated processes (e.g. see James, 1994): (b) thermally-direct Hadley cell, (c) barotropic instabilities and (d) baroclinic instabilities.

These exchanges of energy between zonally symmetric flows and eddies, and between potential and kinetic energies, are commonly quantified in the manner originally suggested by Lorenz (1955, 1967), [building on the much earlier work of Margules \(1903\)](#), in which the flow is partitioned between zonally symmetric (zonally averaged) components and departures therefrom (“eddies”), with four main energy reservoirs (AE, AZ, KE, KZ) representing respectively the eddy and zonal available

potential energy, and the corresponding kinetic energies. From considerations of the energy conservation equations, expressions can be defined to represent the rates of conversion between the four energy reservoirs (e.g. see Lorenz, 1967; Peixoto and Oort, 1974; James, 1994), typically shown schematically in Figure 1(a).

As will be discussed further below, baroclinic or barotropic instabilities and axisymmetric overturning circulations can be distinguished energetically by differing routes of energy conversion between potential and kinetic energies. But schematically, barotropic instabilities are characterised by the dominance of the CK conversion term from zonal mean kinetic energy KZ to eddy KE (see Figure 1(c)), while energy conversions in baroclinic instabilities are dominated by the CA and CE terms, converting zonal mean potential energy AZ to eddy potential and kinetic energy, AE and KE (see 1(d)). The direct kinetic energy conversion between KE and KZ may also take place in the latter case in either direction and so is indicated by the dashed arrows in Fig. 1(d). Such circulations may also be accompanied by direct, axisymmetric overturning circulations that primarily involve the CZ conversion between AZ and KZ (e.g. see Fig. 1(b)).

In addition to straightforward energy considerations, however, the vorticity configuration (either absolute or potential vorticity) also plays a key role in determining whether an instability develops and the character of the instability when it occurs. The subsequent nonlinear development of the instability then depends also on both energetic and vorticity constraints, posing significant challenges to our understanding of how such instabilities will equilibrate.

Predicting and quantifying such equilibration processes are crucial tasks if we are to gain a full understanding of the factors that govern the observable structure and evolution of atmospheric circulation systems under a variety of different conditions. Recent exploration of our Solar System has already provided much information on the structure and properties of large- and medium-scale circulation systems across a wide range of parameter space, from very slowly rotating terrestrial planets such as Venus and Titan, through rapidly rotating terrestrial planets such as Earth and Mars to the fast rotating gas and ice giants. Baroclinic and/or barotropic instabilities likely play important roles in most of these atmospheric systems, but precisely how the occurrence and nonlinear development of such instabilities governs the structure and energetics of their circulation is still very much the subject of ongoing research, using both observations and numerical circulation models. Many planetary atmospheric circulation systems, for example, manifest complex, often anisotropic and inhomogeneous macroturbulent cascades of energy and enstrophy (squared vorticity), but how are such cascades energised and what processes determine the typical equilibrated distribution of energy? How are such states maintained by the input of thermal energy from the Sun or from the deep planetary interior?

Determining the occurrence or otherwise of either barotropic or baroclinic instabilities has traditionally appealed to the results of linearised theories, often (though not exclusively) derived from simplified forms of the equations of motion, continuity and energy conservation. The most well known approach derives from the Charney-Stern-Pedlosky (CSP) criterion for the existence of a non-zero growth rate of an infinitesimal perturbation to a background zonal flow (e.g. Vallis, 2017). This essentially reduces to the requirement that

$$-c_i \int_{y_1}^{y_2} \left\{ \int_0^H \frac{\partial Q / \partial y}{|U - c_r|^2} |\tilde{\psi}|^2 dz + \left[\frac{F \partial U / \partial z}{|U - c_r|^2} |\tilde{\psi}|^2 \right]_0^H \right\} dy = 0, \quad (1)$$

where $U(y, z)$ is the background zonal flow and $Q(y, z)$ the corresponding quasi-geostrophic potential vorticity (QGPV), y is the lateral and z the vertical coordinate, $c = c_r + ic_i$ is the complex phase speed of the perturbation, $\tilde{\psi}(y, z)$ the amplitude of the perturbation streamfunction and $F = f_0^2/N^2$. $f_0 = 2\Omega \sin \phi_0$ is the mean Coriolis parameter, centred at latitude ϕ_0 with Ω the planetary rotation rate, and N is the Brunt-Väisälä frequency. The integrals are carried out over a domain in (y, z) bounded by rigid walls or other suitable boundary conditions at $y = y_1, y_2$ and $z = 0, H$.

For $c_i \neq 0$, therefore, we require one or more of the following criteria to be satisfied.

- (i) $\partial Q/\partial y$ changes sign in the interior domain;
- (ii) The interior $\partial Q/\partial y$ take the opposite sign to $\partial U/\partial z$ at the upper boundary at $z = H$;
- (iii) The interior $\partial Q/\partial y$ take the same sign as $\partial U/\partial z$ at the lower boundary at $z = 0$; or
- (iv) $\partial Q/\partial y = 0$ in the interior and $\partial U/\partial y$ takes the same sign at both $z = 0$ and $z = H$.

Thus, the Earth's atmosphere typically satisfies criterion (iii), being dominated by the planetary vorticity gradient $\beta = df/dy$ away from the lower boundary and $\partial U/\partial z > 0$ at the ground, consistent with geostrophic balance and an equatorward temperature gradient. In the oceans, in contrast, either criteria (i) or (ii) may be satisfied to result in mesoscale eddies generated by baroclinic instability. But whether or how these instability criteria may be satisfied on other planets, where conditions may be very different from Earth, is often unclear, leaving open questions as to the respective roles of baroclinic or barotropic instability in the global circulation.

Even where it can be established that one or other CSP criterion can be satisfied, the fully-developed nonlinear form of the instability may also be hard to predict without the use of complex and expensive numerical models. One possible hypothesis, however, that has proved insightful and interesting in some circumstances is the suggestion that the instability grows vigorously and feeds back onto the background flow, tending to restore it to a state of marginal instability or some other well defined state. For baroclinic instabilities, this process is often referred to as *baroclinic adjustment* (e.g. Stone, 1978; Zurita-Gotor and Lindzen, 2007; Schneider, 2007), by analogy with the concept of rapid convective adjustment towards a state of neutral static stability in a convectively unstable flow. Such a process might be expected to occur where the background unstable flow is only weakly forced or maintained, on timescales that are long compared with the advective timescale of the developing baroclinic instability. The applicability of this concept to the Earth's atmosphere was first suggested by Stone (1978), based on an application of the Phillips two-layer model of baroclinic instability. However, the validity of this application is controversial because the Earth's atmosphere is always baroclinically unstable, even though the dominant perturbations may become much shallower than the height of the troposphere. Thus, a state of marginal instability is not well defined, leading to alternative hypotheses for the adjusted state.

In this paper we consider the application of the CSP and other stability criteria to rotating, stratified flows that include laboratory experiments and both idealised and observed planetary atmospheres. We also explore some of the properties of the fully developed circulations with a view to quantifying the energetics of the equilibrated state and identifying the action and consequences of either baroclinic or barotropic adjustment. Section 2 reviews the results of laboratory experiments on

fully developed baroclinic and barotropic instabilities across a broad range of parameter space, examining the usefulness or otherwise of the CSP criteria and any evidence for signatures of adjustment or other forms of self-organized criticality. Section 3 reviews the corresponding properties of idealised general circulation model (GCM) simulations of an Earth-like planetary atmosphere circulation over a broad range of rotation rates, for comparison in Section 4 with the known properties of the atmospheric circulations of the main planetary bodies of the Solar System. We draw some conclusions and an outlook for future research in Section 5.

2 Baroclinic and barotropic instabilities in the laboratory

Laboratory experiments on rotating, stratified flows provide a powerful vehicle for exploring and understanding the fully developed forms of processes such as baroclinic and barotropic instability. The imposed boundary conditions and experimental parameters can often be closely controlled so that experiments can be repeated and checked, while the geometry can be kept relatively simple so that the respective influence of different factors can be tested over a wide range of conditions. Measurement techniques have also advanced considerably in recent years, allowing optical and other non-intrusive flow measurement as well as the use of in situ probes. Both baroclinic and barotropic flow configurations have been explored in past work over the past 50 or more years, with insightful results on bifurcations between steady, periodic and chaotic flow states and heat and momentum transfer properties.

2.1 Barotropic instabilities of shear layers and jets

2.1.1 CSP stability criteria and energy exchanges

For a purely barotropic flow, the CSP analysis reduces $\partial Q/\partial y$ to $\beta - \partial^2 U/\partial y^2$, leading to stability criterion (i) taking the familiar Rayleigh-Kuo form for the stability of a two-dimensional flow. In effect this represents the need for an inflection point (or related feature) in a zonal shear flow as a necessary but insufficient criterion for its instability. Such a criterion is necessary to enable the possibility of supporting a pair of zonally-propagating Rossby-like waves that can phase-lock across the inflection point and lead to the growth of an instability e.g. via over-reflection (e.g. see Lindzen, 1988). The combination of local vorticity gradients and Doppler shifting of Rossby-like wave trains located in different parts of the flow allow for the possibility of pairs of such wave trains to propagate at the same velocity relative to their common frame of reference. Provided their lateral extent is sufficient for them to interact, these pairs of wave trains can remain mutually coherent and stationary in a common frame of reference moving with the waves, allowing them to interact strongly and draw kinetic energy from the background flow if their phase relationship is favourable.

This generally entails a phase tilt across the channel in such a way that the waves “lean into” the sheared zonal flow so that the momentum flux $\overline{u^*v^*}$ acts to reduce the shear of the background zonal flow. This correlation between the eddy momentum flux and zonal shear is quantified by the CK term in the Lorenz energy budget:

$$CK = \left\langle \frac{d\overline{u}}{dy} \overline{[u^*v^*]} \right\rangle, \quad (2)$$

where u^* and v^* are perturbation velocities with respect to the zonal mean, indicated by the overbar, angle brackets denote a vertically integrated areal average and square brackets denote a time average. This leads naturally to $CK > 0$ such that kinetic
130 energy is transferred from KZ to KE.

2.1.2 Barotropically unstable flows in the laboratory

It turns out that generating a flow in the laboratory with such an inflection point favourable for supporting phase-locked Rossby wave pairs is reasonably straightforward, because such a flow structure occurs spontaneously in a viscous Stewartson shear layer. A detached shear layer may be produced in a rotating, homogeneous fluid by use of differentially-rotating horizontal
135 boundaries, e.g. by the use of a differentially rotating disk or ring at the centre of a cylindrical tank. This was originally investigated by Hide and Titman (1967) and later explored in experiments by Fröh and Read (1999a), Fröh and Read (1999b) and Aguiar et al. (2010). A schematic diagram of the apparatus is shown in Figure 2(a-b) and a typical apparatus illustrated in Figure 2(c).




Figure 2. Schematic cross-sections of two versions of apparatus for producing and studying barotropic instabilities in a cylindrical tank of fluid on a rotating table at angular velocity Ω . In (a) a differentially-rotating, axisymmetric ring is used to drive a barotropic jet at relative angular velocity ω at radius R , while in (b) a pair of differentially rotating disks, centred on the rotation axis, generate a detached Stewartson shear layer, also at radius R . In these cases, a topographic β -effect is also included by use of a conically sloping bottom with slope angle θ . (c) Photograph of the detached shear layer apparatus as used by Fröh and Read (1999a) for investigating fully developed barotropic instabilities.

The resulting directly forced zonal flow has a clear inflection point in its radial velocity profile in such a way that the radial
140 vorticity gradient clearly changes sign. A typical example is illustrated in Figure 3 for a jet-like flow, driven by a differentially-rotating ring. The azimuthal velocity is shown in Fig. 3a while the corresponding vorticity gradient is shown in Fig. 3b.

This would appear to suggest that the flow is unstable, yet it is observed in this state in a time average under conditions in which a regular azimuthal wavenumber $m = 6$ flow is found. The CSP condition, however, is only a necessary condition




Figure 3. A typical azimuthal and time mean jet flow, driven by a differentially rotating ring as used by Aguiar et al. (2010), derived from particle image velocimetry measurements. (a) shows the radial profile of azimuthal velocity while (b) shows the corresponding radial gradient of shallow water potential vorticity, clearly confirming a change of sign of $\partial Q/\partial y = -\partial Q/\partial r$ at $r \simeq 12$ and 17 cm, on either side of the barotropic jet ($\partial Q/\partial y$ due solely to the gradient of $2\Omega/h$ is shown as a dashed line). (c) shows a scatter plot of u against $\partial Q/\partial y$, illustrating the nearly linear relationship between the two quantities within the barotropically unstable jet for $\bar{u} > 0.2 \text{ cm s}^{-1}$.

for instability, not a sufficient one. An alternative condition for instability was considered by Arnol'd (1966) and is often
 145 referenced as “Arnol'd’s second stability theorem” (e.g. Dowling, 1995). This approach considers the instability that could
 arise from the interaction of two (or more) parallel wave trains (commonly, as herein, taken to be Rossby waves, although
 this may be generalised to other interacting wave modes, e.g. see Ripa (1983); Hayashi and Young (1997); Sakai (1989); Gula
 et al. (2009); Flór et al. (2011)), each centred on a local maximum or minimum of potential vorticity gradient. The existence or
 otherwise of a steadily moving reference frame in which both wave trains, Doppler shifted by the zonal flows in their vicinity,
 150 can be at rest relative to each other, provides a criterion for their sustained mutual interaction and hence exchange of momentum
 to gain energy at the expense of the background zonal flow. The resulting condition for perturbations to be stable is of the form:

$$\left(\frac{\bar{u} - \alpha}{\partial \bar{Q}/\partial y} \right) \geq L_0^2, \quad (3)$$

where L_0 is a constant length scale which, depending on the model, may represent the Rossby radius of deformation (L_D), and
 155 α is a constant representing the speed of the waves in a shifted inertial reference frame. The marginal stability condition is set
 by replacing the inequality in Eq (3) by an equality, such that \bar{u} is linearly related to $\partial \bar{Q}/\partial y$ with an offset α at the intercept
 where $\partial \bar{Q}/\partial y = 0$ (e.g. Dowling, 1995).

In practice, however, marginally stable real flows with both forcing and dissipation will not obey this condition exactly, but may approximate to it if the forcing and dissipation are relatively weak compared to the advective transport of Q . For the laboratory flow presented in Fig. 3(a) and (b), this is illustrated in Fig. 3(c), which shows a plot of \bar{u} against $-\partial\bar{Q}/\partial r$ ($= \partial Q/\partial y$ with y directed towards the rotation axis), where Q is now the shallow water potential vorticity

$$Q = \frac{\zeta + 2\Omega}{h(r)}, \quad (4)$$

ζ is the relative vorticity and $h(r)$ is the radius-dependent depth of the fluid layer. Fig. 3(c) clearly shows a strong and nearly linear correlation between $\partial Q/\partial y$ and \bar{u} where $\bar{u} > 0.2 \text{ cm s}^{-1}$, corresponding to the core of the zonal jet in Fig. 3(a). Thus, even though the equilibrated time-averaged flow continues to satisfy the CSP condition for instability, it has effectively stabilised the flow to a configuration that is close to marginal stability with regard to Arnol'd's second stability condition (hereafter *Arnol'd II*), as a result of a process which can be regarded as a form of *barotropic adjustment*.

2.1.3 Circulation regimes and wavenumber selection

The resulting flows thus equilibrate to a marginally stable state in which the original jet meanders in the form of a train of azimuthally travelling waves, on either side of which are patterns of closed vortices whose vorticity matches in sign with that of the flanks of the zonal jet. The flow is typically found to select just one dominant zonal wavenumber in many of the published experiments (e.g. see Figure 4), the wavelength of which depends mainly upon a combination of the Rossby Ro and Ekman E numbers, commonly defined (e.g. Fröh and Read, 1999a, b) as

$$Ro = \frac{R\omega}{2\bar{\Omega}d}, \quad (5)$$

$$E = \frac{\nu}{\bar{\Omega}d^2}, \quad (6)$$

where R is the radius and ω the relative angular velocity of the shear layer or jet, ν the kinematic viscosity of the fluid and d the depth of the fluid layer. $\bar{\Omega}$ is the mean angular velocity of the fluid, representing the average of the turntable rotation rate Ω and the differential rotation rate ω .

As shown by Niino and Misawa (1984) and Fröh and Read (1999a), although the inviscid CSP criterion (*i*) or (more realistically) Arnol'd II criterion needs to be satisfied for the driven jet to be unstable, the observed stability boundary in a real, viscous fluid is also consistent with the existence of a critical Reynolds number Re_c , defined with respect to a length scale of the same order as that of the $E^{1/4}$ Stewartson layer, such that

$$Re_c = \frac{UL_S}{\nu} \simeq Ro.E^{-3/4} \simeq 27, \quad (7)$$

where

$$L_S = (E/4)^{1/4}d \quad (8)$$

is the width of the viscous Stewartson layer. This expression provides a reasonably accurate description of the observed stability boundary (e.g. see Fig. 4) for both the jet and detached shear layer flows, such that for $Re < Re_c$ the flow observed is essentially axisymmetric with no discernible wave perturbation.




Figure 4. Regime diagram for the barotropic jet flows as a function of Rossby and Ekman number, as obtained by Aguiar et al. (2010), showing snapshots of typical equilibrated flows as visualised using dye and side illumination. The trend towards higher wavenumber as the stability boundary at $Re = Re_c$ is approached is clearly seen.

The corresponding boundaries between different flow regimes dominated by different wavenumbers are observed empirically to lie almost parallel to the main stability boundary, suggesting that the wavenumber selected is largely determined by the effective Reynolds number, and hence the effective criticality of the barotropic instability ($Re - Re_c$). In fully developed flows this may be consistent with other work on scale selection in barotropic instability, which associates the most unstable wavelength with a scale comparable to the width of the background jet or shear layer (e.g. Sommeria et al., 1991; Vallis, 2017). In these barotropic instability experiments, however, the equilibrated amplitude of unstable waves grows with criticality, effectively broadening the width of the original jet beyond its initial Stewartson layer width. At its equilibrated finite amplitude, therefore, strongly supercritical flows will tend to favour longer wavelength disturbances, qualitatively consistent with the observed ordering of different wavenumber regimes. This can be clearly seen in Fig. 4, in which the lower wavenumber states are seen to fill the entire tank with large meanders and vortices, whereas high wavenumber flows tend to be more tightly

confined in radius to the vicinity of the unstable jet beneath the rotating ring. At large supercritical Reynolds numbers, however, the equilibrated flow may no longer be steady but may become chaotic, as observed at the lowest Ekman numbers (see Fig. 4), although this has not been explored in much detail as yet.

This method of forcing and maintaining a barotropically unstable flow is relatively strong and fast, so even at large equilibrated amplitudes the zonally averaged flow maintains a reversal in the lateral gradient of potential vorticity. The forcing therefore would seem to be too strong to allow the zonally symmetric component of the flow to maintain itself close to a marginally unstable state. The resulting equilibrated flow is therefore strongly nonlinear, especially at relatively large effective Reynolds numbers.

2.2 Baroclinic instabilities

Baroclinic instabilities require the maintenance of a zonally symmetric distribution of density or temperature that is statically stable in the vertical direction together with a horizontal temperature gradient. This configuration has been studied intensively for more than 60 years in the laboratory using the thermally-driven, rotating annulus (see Figure 5). A fluid is contained between two upright, coaxial metal cylinders, which can be maintained at two different temperatures. Most typically, the inner cylinder is maintained at a cooler temperature than the outer, schematically representing the thermal contrast maintained between the equator and poles of an Earth-like planetary atmosphere. The annular tank is mounted on a rotating table which rotates at angular velocity Ω , with the axis of rotation aligned with the axis of symmetry of the cylinders, again by analogy with a planetary atmosphere. The thermal contrast, ΔT , between the cylinders drives an axisymmetric overturning circulation in the annular channel, mainly confined to boundary layers, which enables a stable stratification to develop and equilibrate, while the effect of background rotation is to induce a vertically sheared azimuthal flow as fluid moves between inner and outer cylinders while roughly conserving its angular momentum.

At a fast enough rotation rate, radial flow becomes largely confined to shallow Ekman layers close to the horizontal (thermally insulating) boundaries, while the isotherms in the interior of the annular channel develop an inclined slope with respect to the horizontal, thus effectively storing potential energy. The flow patterns observed in such a system are then found to depend strongly upon both the rotation rate and the imposed thermal contrast between the cylindrical boundaries (e.g. see Hide and Mason, 1975; Read et al., 2015, for reviews). The approximate sequence of bifurcations observed between different circulation regimes is shown schematically in Figure 6 as a function of the principal dimensionless thermal Rossby number, \mathcal{Ro}_T , and Taylor number, \mathcal{T} , defined respectively as

$$\mathcal{Ro}_T = \frac{g\alpha\Delta T d}{\Omega^2 L^2}, \quad (9)$$

$$\mathcal{T} = \frac{4\Omega^2 L^5}{\nu^2 d}, \quad (10)$$

where g is the acceleration due to gravity, α the volumetric expansion coefficient of the fluid, ν is the kinematic viscosity, L is the horizontal width of the annular channel and d its depth.




Figure 5. Schematic layout of the thermally-driven, rotating annulus experiment (a) as a vehicle for the study of rotating, stratified flows, with (b) a photograph of a typical apparatus in the laboratory, clearly showing the metallic inner and outer cylinders and connections to heat exchangers for maintaining the temperature of the outer wall and a ring of thermocouple junctions to measure temperatures in the interior.

230 As is well known, for a given imposed thermal contrast ΔT , the flow is observed to be stable to baroclinic instabilities for rotation rates slower than a critical value (corresponding to $\mathcal{Ro}_T \gtrsim 2$), resulting in an axisymmetric overturning circulation with prograde flow at upper levels. For higher rotation rates ($\mathcal{Ro}_T \lesssim 2$), the flow becomes unstable to baroclinic disturbances that cause the originally axisymmetric azimuthal flow to meander in the form of a train of azimuthally propagating waves. These baroclinic waves are the fully developed manifestation of baroclinic instability, sometimes referred to as *sloping convection*.
 235 (e.g. Hide, 1969; Hide and Mason, 1975), and may take the form of either regular, near-monochromatic wave trains or more chaotic or even turbulent flows at the highest rotation rates.

2.2.1 CSP instability criteria

Although from an energetic viewpoint it is reasonably clear why the basic state maintained by the differential heating in the rotating annulus experiment can lead to the release of potential energy to energise the growth of wave-like perturbations, it
 240 is less clear how the flow might satisfy the CSP criteria for instability. Early work by Hide (1969) compared the conditions under which active sloping convection was observed in the laboratory with various extensions of the linear model of baroclinic instability by Eady (1949). Provided account was taken of the dissipative effects of viscous Ekman layers adjacent to the upper




Figure 6. Regime diagram for the baroclinically stable and unstable flows in a thermally-driven, rotating annulus as a function of thermal Rossby number (shown here as “Stability parameter”) and Taylor number (e.g. Read et al., 2015), showing snapshots of typical equilibrated flows as visualised using neutrally buoyant tracer particles and side illumination to obtain streak images.

and lower boundaries, these comparisons found generally good quantitative agreement for the onset of the instability around a Burger number, defined as

$$Bu = \frac{N^2 d^2}{4\Omega^2 L^2}, \quad (11)$$

of order $Ro_T/4 \simeq 0.5$. Such an analogy, however, presumes that potential vorticity gradients in the interior are negligible compared with the thermal gradients at the horizontal boundaries, satisfying the CSP stability condition through criterion (iv).

In practice, however, the strong radial flow within the Ekman layers in typical annulus experiments rapidly transfers hot or cold fluid across the domain, maintaining relatively weak horizontal thermal gradients at the top of each Ekman layer. This

would suggest that the potential vorticity dynamics is controlled much less by the structure of the flow close to the horizontal boundaries and is more strongly influenced by a change of sign of $\partial Q/\partial y$ in the interior flow. This would indicate that the basic zonal flow maintained by the differential heating at the side boundaries is closer in character to an internal jet (e.g. Charney and Stern, 1962; Bell and White, 1988). Baroclinic or barotropic instability would therefore occur through satisfying the CSP conditions directly through criterion (i). The consequences of this in the context of rotating annulus experiments were explored by Bell and White (1988), who noted that idealised, inviscid, baroclinic internal jets without any lateral shear would become unstable at around the same value of Burger or thermal Rossby number as (or slightly larger than) the classical Eady problem. This was broadly in agreement with experiments, which did show a tendency for the onset of instability to occur at slightly larger values of Ro_T than the Eady problem would suggest. However, the critical value of Bu or Ro_T was found to be quite strongly sensitive to the addition of lateral shear to a baroclinic internal jet (Bell and White, 1988).

Numerical analyses, using more realistic linearised models based on the full Boussinesq Navier-Stokes equations with molecular viscosity and thermal diffusion (e.g. Lewis and Nagata, 2004; Lewis et al., 2015, and references therein), however, show close agreement with experimental measurements and also reveal sensitivity of the instability to other factors such as the Prandtl number, especially at low values of both Ro_T and \mathcal{T} along the boundary of the so-called “lower symmetric” regime. At high Prandtl number, the boundary roughly follows a line of constant ΔT , indicative of a critical Rayleigh (\mathcal{Ra}_c) or Grashof number (\mathcal{Gr}_c) in a similar way to the criterion for Rayleigh-Bénard convection. The product $Ro_T \cdot \mathcal{T}$ actually has the form of a Grashof number, such that $Ro_T \cdot \mathcal{T} = \mathcal{Gr}_c$ is consistent with the shape of the lower left instability boundary shown in Fig. 6.

2.2.2 Lorenz energy cycle

As instabilities develop and equilibrate, azimuthally travelling waves emerge with characteristic three-dimensional structures and azimuthal phase tilts in radius and height. These result in eddy fluxes of heat and momentum that lead to energy conversions between potential and kinetic energies in the zonally averaged and perturbation fields. A typical example of the Lorenz energy cycle for an equilibrated baroclinic wave developed from a baroclinically unstable initial state is shown in Figure 7. This was computed by Young (2014) from a numerical simulation of flow in a rotating annulus experiment from solutions of the Boussinesq Navier-Stokes equations in cylindrical annular geometry. The boundary conditions represent non-slip, rigid sidewalls and horizontal endwalls, thermally insulating endwalls and isothermal sidewalls at temperatures T_a and T_b at inner and outer radii r_a and r_b , with $\Delta T = T_b - T_a = 4$ K, $r_a = 2.5$ cm, $r_b = 8.0$ cm, $d = 14$ cm and rotation rate $\Omega = 3.0$ rad s⁻¹, corresponding to $Ro_T = 0.056$.

The values shown in Fig. 7 clearly indicate that the conversion terms implicated in energising baroclinic instability (CA and CE; see Fig. 1(b)) are dominant in the exchanges between eddies and the zonal mean flow. CK is at least a factor of 10 smaller than CA though in the same sense for the generation of KE, indicating that the dominant instability has a somewhat mixed baroclinic-barotropic character. The direct zonal mean exchanges (CZ) are also of a similar magnitude to CA and CE, and are consistent with a thermally direct overturning circulation in parallel with the conversion of available potential energy to KE.




Figure 7. An example Lorenz energy cycle, computed by Young (2014) from a numerical simulation of an equilibrated baroclinic wave flow in a differentially-heated, rotating annulus. Energy reservoir and conversion and source/sink terms are in units of $10^{-7} \text{ J kg}^{-1}$ and W kg^{-1} respectively.

2.2.3 Signatures of baroclinic adjustment in heat transport?

285 As mentioned above, baroclinic adjustment is commonly associated with a tendency for baroclinic instability to equilibrate at finite amplitude in such a way as to modify the initially unstable basic state towards a less unstable configuration. But it is less clear how this tendency might manifest itself in practice. However, one possible way this might be observable could be in the way heat transport by baroclinic disturbances varies with external parameters as the instability threshold is exceeded.

Figure 8 shows a series of calorimetric measurements of total heat transport in a differentially heating rotating annulus experiment for fixed ΔT as Ω is varied (see Read, 2003). In this case, heat transport is non-dimensionalised with respect to thermal conduction, either by the Nusselt number, Nu, defined by

290

$$\text{Nu} = \frac{\mathcal{H} \ln(r_b/r_a)}{2\pi\kappa d\rho c_H \Delta T}, \quad (12)$$

where \mathcal{H} is the total heat transport in W, κ is the thermal diffusivity of the working fluid and c_H its specific heat capacity, or by the Péclet number $\text{Pe} = \text{Nu} - 1$. Thus the solid symbols in Fig. 8 show little variation with Ω or \mathcal{R}_{OT} until the highest rotation rates. This corresponds to where the regular wave regime in the laboratory experiments begins to break down towards small-scale irregular flows and geostrophic turbulence. The solid line in Fig. 8, on the other hand, shows the variation of Nu for a purely axisymmetric flow, obtained from numerical simulations in which non-axisymmetric eddies were suppressed (see Read, 2003, for details). This shows that the axisymmetric flow under similar conditions decays much more quickly with Ω than when baroclinic waves are allowed to develop. The corresponding contribution to the dimensionless total heat transport




Figure 8. Variations of radial heat transport by baroclinic eddies and axisymmetric flows in a differentially heated rotating annulus, as measured in the laboratory (Read, 2003) as a function of rotation rate and thermal Rossby number \mathcal{Ro}_T . The total advective heat transport is indicated by the dimensionless Péclet number ($Pe = Nu - 1$, where Nu is Nusselt number; see Eq (12)). Solid symbols represent total heat transport as measured by calorimetry, while the continuous line represents the total heat transport by the axisymmetric flow alone. Open symbols indicate the difference, Pxs , in Pe between measured total advective heat transport and the axisymmetric component, representing the heat transport due to eddies alone.

300 by baroclinic waves is indicated by the open symbols, which are obtained from the difference between total heat transport and that of the pure axisymmetric flow. This clearly shows that the eddy contribution increases systematically from zero as the rotation rate is increased beyond the critical value for the onset of instability, until it begins to saturate and even begin to turn over for $\mathcal{Ro}_T < 0.1$.

The observation that total heat transport may be almost independent of rotation rate over a wide range of Ω , from the axisymmetric regime into the fully developed baroclinic wave regime, has been noted since the early work of Bowden (1961),
305 who was the first to carry out calorimetric measurements of heat transport in rotating annulus experiments. This tendency for the heat transport to remain close to its value at $\Omega = 0$, even though the (mainly axisymmetric) boundary layer contribution should reduce substantially over the same range of parameters, is unlikely to be coincidental, but more likely reflects a systematic result of equilibrated baroclinic instability in modifying the overall flow to transport almost as much heat (and therefore release

almost as much potential energy) as the non-rotating flow. The non-rotating flow represents a fully relaxed state, in which isotherms and isopycnals are as near horizontal as possible (except in conductive boundary layers), and is therefore a state that would be energetically stable to further baroclinic instabilities. In this sense, these results demonstrate that rotating annulus flows are capable of exhibiting a relatively strong form of baroclinic adjustment throughout the regular baroclinic wave regime. It is only when the regular regime begins itself to become unstable at values of $\text{Ro}_T \ll 0.1$, where the dominant energetic length scale of the baroclinic waves becomes significantly smaller than the width of the imposed baroclinic zone, that the eddies are no longer able to sustain the level of heat transfer required to maintain the fully relaxed flow state and the baroclinically adjusted state breaks down to a more strongly supercritical flow.

Such a tendency would seem to be quite general, and so it is of interest to see whether similar trends might be observable in a planetary atmosphere, in which differential heating is maintained by (radiative) relaxation towards a baroclinically unstable radiative equilibrium state.

3 Baroclinic and barotropic instabilities in idealised planetary atmospheric circulations

The Earth and other Solar System planets present us with just a few samples of atmospheric circulation systems in different parts of a broader parameter space (e.g. see Read, 2011). A better way of exploring trends and scaling of various properties of planetary atmosphere circulations is to use a simplified numerical model, in which the key forcing processes are represented schematically but in a physically consistent manner as a function of the principal planetary parameters, so that the model can be used to explore the relevant dynamical response. This approach has a long history, dating back to the early work of Hunt (1979), Williams and Holloway (1982) and Geisler et al. (1983), but has become more common in recent years, inspired in part by the diversity of newly discovered extrasolar planets (e.g. Merlis and Schneider, 2010; Mitchell and Vallis, 2010; Kaspi and Showman, 2015; Wang et al., 2018). Here we focus on diagnostics of energetics, instabilities and heat transfer in a typical set of idealised GCM simulations, based on the work of Wang et al. (2018).

3.1 Circulation regimes

As demonstrated by various authors (e.g. Kaspi and Showman, 2015; Wang et al., 2018), many generic features of the large-scale circulation of almost any planetary atmosphere can be captured in a general circulation numerical model that solves the hydrostatic primitive equations of meteorology, together with simple parameterizations of diabatic heating and small-scale turbulent mixing and friction. In recent work, Wang et al. (2018) has used the University of Hamburg PUMA model (e.g. Frisius et al., 1998; Fraedrich et al., 2005) to explore the dependence of the simulated fully three-dimensional, time-dependent circulation of an Earth-like planetary atmosphere on dimensionless parameters such as the thermal Rossby number, Ro_T , as quantities such as the planetary rotation rate, $\Omega^* = \Omega/\Omega_E$ (where Ω_E is the rotation rate of the Earth), are varied. This model represents fields in the horizontal as projections onto sets of spherical harmonic functions but in finite difference form in the vertical. Diabatic heating and cooling is represented by a linear relaxation towards a prescribed zonally-symmetric temperature field, intended to represent the diurnally and seasonally averaged radiative-convective equilibrium of an Earth-like

Ω^*	\mathcal{R}_{oT}	\mathcal{T}_F	\mathcal{T}_R
1/16	20.5	238	1.71×10^5
1/8	5.14	3.80×10^3	2.73×10^6
1/4	1.28	6.09×10^4	4.67×10^7
1/2	0.32	9.74×10^5	6.98×10^8
1	0.08	1.56×10^7	1.12×10^{10}
2	0.02	2.49×10^8	1.79×10^{11}
4	0.005	3.99×10^9	2.86×10^{12}
8	0.0013	6.38×10^{10}	4.58×10^{13}

Table 1. Key dimensionless parameters for the baseline set of numerical simulations with $\Delta\theta_{EP} = 60\text{K}$, $\tau_{ft} = 5$ Earth days and $\tau_R = 25.9$ Earth days, as defined by Equations (13) and (15).

planet, with a prescribed relaxation timescale τ_R . Surface friction is represented by a height-dependent, linear Rayleigh friction parameterization with local timescale τ_F and total (Ekman) spin-down timescale of τ_S .

The model was run to equilibrium over 10-20 Earth years before computing various diagnostics of the statistically equilibrated state. The circulation regime could then be characterised as a function of various dimensionless parameters, such as the thermal Rossby number,

$$\mathcal{R}_{oT} = \frac{R\Delta\theta_{EP}}{\Omega^2 a^2}, \quad (13)$$

where R is the specific gas constant, $\Delta\theta_{EP}$ is the horizontal temperature contrast between equator and poles and a is the planetary radius, the Burger number,

$$\mathcal{B}u = \frac{N^2 H^2}{4\Omega^2 a^2} \simeq \frac{R\Delta\theta_z}{4\Omega^2 a^2}, \quad (14)$$

where $\Delta\theta_z$ is the vertical contrast in potential temperature, and frictional or radiative Taylor numbers,

$$\mathcal{T}_{(F,R)} = 4(\Omega(\tau_F, \tau_R))^4. \quad (15)$$

The results of a scan through parameter space as Ω^* is varied may be summarised in a regime diagram with respect to \mathcal{R}_{oT} and \mathcal{T}_F , which identifies and classifies different circulation regimes according to their location in parameter space.

Figure 9 presents the regime diagram for the PUMA simulations of Wang et al. (2018), together with some sample snapshots of typical flows at different planetary rotation rates which present visualisations of the zonal wind speed at the upper tropospheric level of 200 hPa. As discussed more fully by Wang et al. (2018), the sequence of circulation patterns follows a clear




Figure 9. Regime diagram showing the various circulation regimes obtained in a simplified atmospheric general circulation model (Wang et al., 2018) with respect to characteristic dimensionless parameters (\mathcal{R}_T and \mathcal{T}_F). Stars refer to experiments in which wavy flows are observed, whereas open diamonds indicate experiments in which axisymmetric flows were found. The approximate location in parameter space of some Solar System planets (Earth, Mars, Titan, Jupiter, Saturn, Uranus and Neptune) are labeled by their initial letters. The solid line delineates the boundary between axisymmetric circulations and circulations with wavy/turbulent flows. The dashed lines indicate the boundaries between different circulation regimes within the wavy/turbulent region. Some example snapshots of typical circulation patterns, visualised by shading of the zonal wind speed at the 200 hPa level, are indicated by the arrows.

trend with planetary rotation rate. Keeping $\Delta\theta_{EP}$ constant, the locus of varying rotation rate follows a diagonal path from upper left to bottom right in $(\log \mathcal{R}_T, \log \mathcal{T}_F)$. At the lowest values of Ω^* the circulation is dominated by unsteady circumpolar vortices, a large-scale meridional overturning and strongly super-rotating zonal winds, both globally and locally (cf Read and Lebonnois, 2018). These evolve into a pattern of more Earth-like, meandering mid-latitude jet streams at intermediate values of Ω^* . This then breaks down into a more irregular pattern of parallel zonal jets, reminiscent of Jupiter and Saturn (although the latter both exhibit locally super-rotating equatorial jets, unlike the present model simulations), the number of which increases with increasing Ω^* .

From an examination of the temperature and PV structure of the flow, it is evident that the basic circulation in these models allow for either barotropic or baroclinic instabilities, through satisfying either CSP criteria (i) or (iii). In the latter case, the method of diabatic forcing maintains a strong equatorward thermal gradient at the lower boundary which, for quasi-geostrophic




Figure 10. Terms in the Lorenz energy budgets for the series of PUMA simulations as a function of Ω^* and thermal Rossby number: (a) globally averaged energies (in 10^5 J m^{-2}); (b) the main energy conversion rates CZ, CA, CE and CK. Conversion rates are in unit of W m^{-2} .

conditions, enables CSP criterion (iii) to be satisfied. Where a strong mid-latitude or circumpolar jet is formed, however, then criterion (i) may be satisfied through lateral variation in the vorticity of the jet. Precisely which will dominate in particular cases, however, is not immediately clear without a more detailed analysis.

3.2 Energetics

Figure 10 presents the results of an analysis of the time-averaged Lorenz energy cycles for each of a set of PUMA simulations from the work of Read et al. (2018) with fixed thermal forcing, covering a range of Ω^* from 1/16 to 8 (representing a range in Ro_T from $\sim 20 - 10^{-3}$; see Table 1). The energy content of the main reservoirs (Fig. 10(a)) is seen to change from a mainly KZ dominated regime at low Ω^* to a strongly AZ dominated regime at fast Ω^* ($Ro_T \ll 1$). Both the eddy components, AE and KE, remain relatively small but rise to a peak for rotation rates corresponding to $Ro_T \sim 0.1 - 1$. All terms except AZ then decay strongly with increasing Ω^* at the highest rotation rates.

Energy conversion rates exhibit even more complex variation with Ω^* (see Fig. 10(b)). For $Ro_T \gg 1$ the CZ term is strongly dominant, indicating a circulation that is energetically dominated by a thermally direct, zonal mean meridional overturning. Within this range, eddies gain energy from the zonal mean components through both the barotropic CK and baroclinic CE terms, but CK is dominant over CE (and CA) for $Ro_T > 10$. At intermediate Ω^* , the baroclinic conversions, CE and CA, rise to a peak around $Ro_T \sim 0.3$, while CK and CZ decrease and actually change sign for $Ro_T < 1$. This indicates that the energetics of the circulation are dominated by barotropically unstable eddies at the lowest rotation rates, although the baroclinic

conversion term CE is also positive in this range, indicating a mixed barotropic/baroclinic instability as the origin of these
 385 eddies. For $\text{Ro}_T \lesssim 1$, however, CK becomes negative while CE rises to a positive maximum around $\text{Ro}_T \sim 0.3$, indicating
 that eddies are predominantly baroclinic in character and energetics at relatively fast rotation rates. At the highest rotation rates
 all conversion terms are seen to decrease in magnitude as Ro_T decreases, leading to relatively weak and increasingly inefficient
 circulation.

3.3 Heat transfer and baroclinic adjustment

390 The positive contributions to both CA and CE at all rotation rates imply that the simulated circulation involves significant
 transfers of sensible heat across the planet. Figure 11(a) shows how the peak values of vertically integrated meridional heat
 transport vary with Ω^* and Ro_T in the PUMA simulations of Wang et al. (2018). The zonal mean and eddy heat transports are
 shown separately by open triangles and crosses respectively, with the total heat transport (zonal mean plus eddy) indicated by
 solid squares. This is similar in form to the results shown in Fig. 8(a) of Kaspi and Showman (2015) for their simulations using
 395 a GCM driven by a gray radiation scheme with moisture transport, apart from showing the combined total heat transport.

These results clearly show that, as in the simulations of Kaspi and Showman (2015), the zonal mean contribution to heat
 transport dominates at slow rotation rates (for $\text{Ro}_T > 1$) but decreases monotonically with increasing Ω^* . The eddy contribu-
 tion to heat transport, on the other hand, steadily increases (almost linearly) with Ω^* until it peaks around $\text{Ro}_T \sim 0.3$, beyond
 which it also decreases rapidly with increasing Ω^* . The sum of the two contributions, however, remains nearly independent
 400 of Ω^* and Ro_T until $\text{Ro}_T \sim 0.07$, indicating that for values of Ro_T between ~ 1 and 0.07 the eddy heat transport is able to
 compensate for the decrease in zonal mean transport to maintain the total heat transport close to its slowly rotating value.
 At faster rotation rates, however, for which $\text{Ro}_T \lesssim 0.07$, eddies can no longer maintain this strength of heat transport so the
 overall heat transport begins to drop.

Fig. 11(b) indicates that the global temperature structure also reflects these variations in total heat transport as Ω^* is varied.
 405 This shows η , a measure of the mean isentropic slope in the mid-troposphere $[\partial z / \partial y]_\theta$, normalised by the mean slope of
 the isentropes in the radiative equilibrium state to which the thermal state of the flow is relaxed, and the mean equator-pole
 temperature contrast, ΔT_{EP} , also in mid-troposphere. These clearly show the isentropic slope and ΔT_{EP} increasing with Ω^* ,
 but with a plateau in this variation for Ro_T between ~ 1 and $\sim 0.1 - 0.05$. For $\text{Ro}_T \lesssim 0.05$, however, the reduction in heat
 transport as the baroclinically adjusted state breaks down leads to a resumption of the increase in isentropic slope with rotation
 410 rate as the thermal structure approaches the radiative-convective equilibrium configuration.

These results show a close similarity with the heat transport results discussed in Section 2.2 for thermally-driven rotating
 annulus experiments, and clearly indicate a regime in which eddy heat fluxes exert a strong influence on the thermal structure
 of the global circulation. This is despite an absence of a clear instability threshold for baroclinic instability in the GCM
 simulations, unlike what is found in the annulus experiments. However, the onset of this compensation effect of eddy heat
 415 fluxes against the decreasing zonal mean transport (effectively acting as an “eddy thermostat”) around a value of $\text{Ro}_T \simeq 1$
 does correspond to where the vertical scale of Charney-type baroclinic instabilities first becomes of the same order as the
 pressure scale height and altitude of the tropopause (e.g. Branscombe, 1983) as Ω^* increases from smaller values. This would




Figure 11. (a) Variations of peak values of meridional heat transport (in W) within the PUMA GCM simulations of Wang et al. (2018) and at slower values of Ω , shown as total poleward transport (blue squares), zonal mean transport ($\int [c_p \bar{v} \bar{T}] 2\pi a \cos(\phi_{max}) dp/g$) and eddy transport ($\int [c_p \bar{v}^* T^*] 2\pi a \cos(\phi_{max}) dp/g$); (b) ratio η of the slope obtained in fully active circulations in the latitude-height plane for experiments with different Ω^* to the globally mass-weighted average isentropic slope in radiative-convective equilibrium (towards which the model atmosphere relaxes, shown as filled triangles connected by solid lines), plotted alongside the variation of mean equator-pole thermal contrast in the mid-troposphere, ΔT_{EP} (shown as open squares connected by a dotted line). The (red) dashed line in (a) indicates a gradient of +1 with respect to Ω suggesting linear proportionality.

suggest that the baroclinic component of the eddies need to span a large fraction of the active troposphere in order to be able to exert a strong influence on its thermal structure. But it does, therefore, indicate that when this condition is satisfied, it can exhibit a similar form of baroclinic adjustment to that found in the laboratory.

It would clearly be of interest to explore this trait in more detail in other, more realistic, GCM studies that also include the effects of radiative transfer and latent heat transport.

4 Baroclinic and barotropic instabilities in the Solar System

As discussed above in Section 1, the Earth's atmosphere satisfies the CSP necessary conditions for instability mainly through criterion (iii), associated with persistent equatorward temperature gradients at the surface and the planetary vorticity gradient in the free atmosphere. The flanks of the upper level mid-latitude jet might also lead to local changes of sign of $\partial Q/\partial y$ in the interior, while horizontal thermal gradients at the tropopause might also allow the CSP conditions to be satisfied at times through criterion (iv) (e.g. see Vallis, 2017, section 9.9). These factors suggest multiple ways in which large-scale instabilities might occur in the Earth's atmosphere, so which mechanism (baroclinic or barotropic) is likely to dominate?

Sections 2.2 and 3 both highlight the importance of the dimensionless parameters, $\mathcal{B}u$ and $\mathcal{R}o_T$, in favouring which mechanism is likely to be most important, although the distribution of $\partial Q/\partial y$ and its sign changes in the horizontal or vertical also plays a role. Table 2 presents some rough estimates of the values of $\mathcal{B}u$ and $\mathcal{R}o_T$ for the Earth, Mars, Venus and Jupiter, based on Eqs (13–15). These parameters for Earth are typically much less than $O(1)$, that would tend to suggest that baroclinic instabilities dominate, as is well known (e.g. James, 1994; Vallis, 2017).

Table 2. Estimates of the main dimensionless parameters for the Earth, Mars, Venus and Jupiter. Radiative time constant estimates are based on values from NASA’s [Planetary Data System \(PDS - See https://pds-atmospheres.nmsu.edu/education_and_outreach/encyclopedia/radiative_time_constant.htm\)](https://pds-atmospheres.nmsu.edu/education_and_outreach/encyclopedia/radiative_time_constant.htm).

Planet	$\mathcal{B}u$	$\mathcal{R}o_T$	τ_R (s)	\mathcal{T}_R
Earth	0.02	0.06	4.1×10^6	3×10^{10}
Mars	0.04	0.20	1.9×10^5	1.3×10^5
Venus (planet)	140	370	$2.2 \times 10^5 - 4 \times 10^9$	$7.5 \times 10^{-5} - 8 \times 10^{12}$
Venus (cloud tops)	0.04	0.10	2.2×10^5	10^3
Jupiter (cloud tops)	8×10^{-4}	10^{-3}	1.4×10^8	1.5×10^{18}

This is confirmed in typical calculations of the Lorenz energy budget for the Earth’s atmosphere. Figure 12(a) shows the result of a typical Lorenz energy cycle, as computed by Boer and Lambert (2008) [from US National Centers for Environmental Prediction \(NCEP\) data](#), with energy reservoirs shown in units of J m^{-2} and conversion rates in W m^{-2} . This clearly shows that much of the convertible energy in the atmosphere resides in the AZ reservoir, which hosts more than twice as much as the rest of the reservoirs put together. The strongest internal conversions are the CA and CE terms, which indicate a net transfer of energy from zonal potential energy AZ to KE via AE, which is the classical route representative of baroclinic instability (cf Fig. 1(d)). The kinetic energy conversion, CK, is relatively small but robustly negative, indicating a transfer from KE to KZ consistent with the driving of a zonal mean eddy-driven jet stream at mid-latitudes (e.g. James, 1994; Vallis, 2017). The AZ reservoir is maintained through the GZ term, representing the effects of differential radiative heating and cooling, while energy is ultimately removed from the system via the dissipative flux terms FZ and FE.

This pattern of energy reservoirs and fluxes is well known for the Earth, and is broadly consistent with the corresponding Earth-like case discussed in Section 3. But what about other planets in the Solar System with substantial atmospheres which may be in very different circulation regimes from Earth? Where might they fit in comparison to the scenarios indicated by the regimes found with simplified GCMs, such as in Section 3?

4.1 Mars

Mars is arguably the most Earth-like planet elsewhere in the Solar System, at least so far as its atmosphere and climate are concerned. It is roughly half the linear size of Earth (radius $a \simeq 3400$ km) with a rotation period of just over 24 hours. It lies at




Figure 12. Representations of energy storage and interconversions in the form suggested by Lorenz (1955) and Lorenz (1967) for a selection of the main planets of the Solar System: (a) Earth (calculations from Boer and Lambert, 2008); (b) Mars (calculations from Tabataba-Vakili et al., 2015); (c) Venus (calculations from Lee and Richardson, 2010) and (d) Jupiter’s weather layer (from preliminary calculations using a higher-resolution version of the model described in Young et al., 2019a). As before, energy reservoirs are shown in units of 10^5 J m^{-2} and conversion rates and source/sink terms in units of W m^{-2} .

a distance of around 1.3-1.5 Astronomical Units (AU) with an orbital period of around 687 days and, with a rotation axis tilted by approximately 25° from the perpendicular to its orbit, experiences a strong seasonal cycle much like the Earth. It possesses an atmosphere mainly consisting of CO_2 with a surface pressure of around 6 hPa which, although much less than on Earth, is sufficient to interact strongly with the rocky surface. Both water and CO_2 can condense as ices to form clouds and surface deposits.

Its temperature structure and rapid planetary rotation lead to relatively small values of \mathcal{R}_{OT} and $\mathcal{B}u$ (see Table 2), though not as small as for the Earth. Its relatively short radiative time constant leads to a value of \mathcal{T}_R which is much larger than $\mathcal{O}(1)$, indicating relatively weak radiative damping compared with Coriolis forces.




Figure 13. Zonal mean sections of (a) zonal wind (m s^{-1}) and (b) $\partial Q/\partial \phi$ ($\text{s}^{-1} \text{ rad}^{-1}$) typical of Martian northern hemisphere winter. Data were taken from the Mars Climate Database (e.g. Lewis et al., 1999, see also <http://www-mars.lmd.jussieu.fr/mars/access.html>) for $L_s = 270^\circ$ using the standard Mars climatology. Contour intervals are (a) 15 m s^{-1} and (b) $0.0002 \text{ s}^{-1} \text{ rad}^{-1}$ and negative contours are shown dashed.

460 4.1.1 PV structure

Its similarities with Earth are also reflected in its potential vorticity structure, as can be seen e.g. in Figure 13, which shows the zonal mean zonal wind and QGPV gradient for a typical atmospheric state during northern hemisphere winter on Mars, from data obtained from the Mars Climate Database (e.g. Lewis et al., 1999, ; see also <http://www-mars.lmd.jussieu.fr>). As can be seen in Fig. 13(a), zonal winds are typically concentrated into deep, mid-latitude jet streams during winter and the adjacent
465 seasons, with strong vertical and horizontal shears and a pronounced **latitudinal tilt with altitude**. The corresponding structure in QGPV (see Fig. 13(b)) shows $\partial Q/\partial y$ ($\partial Q/\partial \phi$) positive in the jet itself (cf Fig. 13(a)) but with weaker reversals in sign on either side. Together with the positive equatorward temperature gradient at the surface in the north, this would indicate that the CSP conditions can be satisfied in this season either by criterion (i) or (iii), much as on Earth. The atmosphere might then be expected to exhibit either baroclinic or barotropic instabilities, or a mixture of both, although the linear instability analysis of
470 Barnes (1984) suggests that the baroclinic instability mechanism dominates for Martian conditions while the barotropic shear mechanism acts to damp the baroclinic instability.

It is interesting to note that, at this time during southern summer, the corresponding zonal jet stream becomes much weaker and even reverses direction in places. The QGPV gradient does exhibit a weak reversal in sign, though the equatorward thermal gradient near the surface also reverses in sign during summer, so that conditions are much less favourable for baroclinic
475 instability via CSP criterion (iii). This is consistent with the observed near suppression of baroclinic instability in Martian

summers at mid-latitudes (e.g. Lewis et al., 2016), except for occasional very shallow disturbances that are sometimes observed on small horizontal scales close to the polar cap edge (e.g. Gierasch et al., 1979b).

4.1.2 Energy budget

The corresponding Lorenz energy budget is needed to establish which is the dominant mechanism to energise the Martian eddy
480 circulation. This has been computed recently by Tabataba-Vakili et al. (2015), based on an assimilated analysis of orbiting spacecraft measurements of Mars over a period of at least 3 Mars years (Montabone et al., 2014). The data were assimilated into the UK version of the [Laboratoire de Météorologie Dynamique \(LMD\)](#) Mars GCM to produce a daily record of Martian meteorology over the entire period, from which the energy budget could be computed (see Figure 12(b)).

The results, illustrated e.g. in Figure 12(b), show a qualitatively similar pattern to that of the Earth, with the majority of
485 dynamically active energy in the atmosphere stored as zonal potential energy AZ. The dominant conversion terms are the CA and CZ conversions from AZ to AE and KZ respectively (at 50 mW m^{-2}), and especially the CE term (at 110 mW m^{-2}). This indicates a strong energetic role for the thermally direct axisymmetric (Hadley cell) overturning circulation as well as the principal baroclinic conversions CA and CE maintaining synoptic eddies. This confirms the role for baroclinic instability as suggested from the configuration of potential vorticity gradients in Fig. 13(b). The barotropic eddy-zonal conversion CK,
490 however, is positive and around half the amplitude of CA, suggesting a role for exchanges consistent with a mixed baroclinic-barotropic instability. The GE source term for AE is also quite substantial and of the same order as CA and CZ. This is because the day-night contrast in solar heating on Mars generates a strong thermal tide response which contributes to the AE and KE reservoirs, although this varies significantly with season and the amount of dust loading in the atmosphere. As discussed by Tabataba-Vakili et al. (2015), this budget does exhibit fairly significant seasonal variations, also between the northern and
495 southern hemispheres, but the overall structure of the energy budget remains largely unchanged.

4.2 Venus

Venus is the other main terrestrial planet in the inner Solar System that possesses a substantial atmosphere. It is also Earth-like in some respects, but very different in others. Around the same size as Earth, Venus is somewhat closer to the Sun than Earth at a mean distance of 0.72 AU and orbits the Sun in around 224.7 days. The planetary rotation period, however, is very long
500 compared to Earth, with a sidereal rotation period of 243.7 days in a retrograde sense and a very small obliquity angle (177.3° , equivalent to 2.6° if it rotated in the same sense as its orbit).

Its atmosphere is deep and massive, consisting mainly of CO_2 with a surface pressure of more than 90 bars and typical surface temperature of 740 K. The entire planet, however, is almost completely shrouded in thick clouds, thought to consist mainly of aqueous sulphuric acid droplets that scatter sunlight very efficiently (with a Bond albedo of 0.77) and located between
505 40 and 60 km above the surface. Dynamically, Venus is in a very different regime to the Earth. Its winds are dominated by a very rapid super-rotation with maximum zonal winds of more than 100 m s^{-1} , located around or slightly above the main cloud decks at altitudes of 50-70 km. This means that the atmosphere at the cloud level rotates around 60 times faster than the underlying surface. The processes driving such a remarkable circulation are still imperfectly understood, though likely entail

the role of waves and eddies on various scales (e.g. see Sánchez-Lavega et al., 2017; Read and Lebonnois, 2018, for recent reviews). Despite such strong winds and dynamical activity, horizontal temperature gradients are relatively weak compared with those in the vertical, with typical equator to pole thermal contrasts of no more than 10-20K.

These factors lead to somewhat different estimates of some of the key dynamical parameters, depending upon whether the atmosphere is viewed in the frame of the underlying planet or in corotation with the main cloud deck. In the frame of the planet, values of $\mathcal{B}u$ and \mathcal{R}_{OT} are likely to be much larger than $O(1)$ (although N is also close to zero in the deep troposphere), indicating that the flow is unlikely to be geostrophic (its cloud level zonal winds are predominantly in cyclostrophic balance with centrifugal accelerations dominating over Coriolis accelerations; e.g. see Sánchez-Lavega et al. (2017)) or baroclinically unstable, although it could be consistent with various forms of barotropic shear instability. If the circulation is viewed in the average frame of the cloud level winds (e.g. see Young et al., 1984), however, which rotate around the planet on average in around 4 days (corresponding to $f = 2\Omega \simeq 3.6 \times 10^{-5} \text{ s}^{-1}$), a different perspective emerges with typical values of $\mathcal{B}u$ and \mathcal{R}_{OT} of around 0.04-0.1. In this frame, therefore, the cloud level circulation may appear to be in quasi-geostrophic balance with the possibility of baroclinic instabilities that would likely be localised in the vertical around the levels of the main cloud decks, provided the potential vorticity configuration would allow the CSP necessary conditions for instability to be satisfied.

4.2.1 PV structure

A key difficulty in determining the PV structure of Venus’s atmosphere arises from a lack of detailed observations of winds and temperature within and beneath its main cloud decks (e.g. see Sánchez-Lavega et al., 2017). It is necessary, therefore, to fall back on numerical models of Venus’s atmospheric circulation for the kind of information needed to compute quantities such as $\partial Q/\partial y$. But the numerical simulation of Venus’s atmospheric circulation is still at a relatively immature state, with realistically forced models appearing only recently (e.g. Lebonnois et al., 2010, 2016; Mendonça and Read, 2016). Simpler models have been studied more extensively but have struggled to reproduce key features of the circulation and have exhibited significant divergence among different model formulations (e.g. Lee and Richardson, 2010; Lebonnois et al., 2013; Lewis et al., 2013). This may reflect important differences e.g. in the computation of static stability (and hence the ratio f/N) and in how well different dynamical cores are able to conserve and transport angular momentum.

For the present purpose, therefore, we follow Young et al. (1984) and Sugimoto et al. (2014) in examining a somewhat idealised form of the zonal mean state of a Venus-like atmosphere. This is illustrated in Figure 14, taken from the work of Sugimoto et al. (2014), and shows a zonal mean section of the zonal wind (Fig. 14(a)) and corresponding field of $\partial Q/\partial \phi$ (in the average frame of the zonal wind at an altitude of 56 km; Fig. 14(b)). The zonal wind section (in the planetary reference frame) captures the main features of the observed zonal winds on Venus, with a monotonic growth of zonal wind strength with altitude at all latitudes towards maximum values of 100-120 m s^{-1} at around 70 km altitude, but with the formation of jet-like features at mid-latitudes in both hemispheres. This idealisation then neglects any vertical shear above the cloud tops, although this may not be particularly realistic. However, the flow structure around the cloud top altitudes are of most interest here. The winds are constructed to be in gradient wind balance (between horizontal pressure gradients and a combination of Coriolis and centrifugal “forces”) with the corresponding pressure and temperature fields.




Figure 14. Zonal mean sections of (a) zonal wind (m s^{-1}) and (b) $a\partial Q/\partial\phi$ (m s^{-1}) typical of Venus’s global circulation, as presented by Sugimoto et al. (2014) from their numerical model simulations.

The distribution of $\partial Q/\partial\phi$ in Fig. 14(b) indicates a concentration of PV gradient at high latitudes, especially within the cloudy layers between altitudes of 55 and 65 km, (although the clouds are not represented explicitly in this model). Moreover, this gradient clearly changes sign in the vertical while remaining positive over most of the domain at lower levels. This would immediately suggest the possibility of a baroclinic instability localised within the cloud deck levels by satisfying the CSP conditions for instability via criterion (i). A change of sign in the horizontal is only discernible in the layer around 56 km, however, suggesting that barotropic instabilities might also be possible, though again centred mainly within the cloud decks. Nearer the surface, $\partial Q/\partial\phi > 0$ but surface temperature gradients are very weak, suggesting that Venus may be able to satisfy the CSP condition via criterion (iv) only marginally. Besides, the flow is far from being in geostrophic balance at these levels, although that would not necessarily rule out the possibility of a non-geostrophic form of baroclinic instability.

4.2.2 Energy budget

The energy budget for Venus is similarly not available from observations and so we must rely on model simulations to compute the likely energies and exchanges. To the authors’ knowledge, this has not so far been done for any realistically forced, comprehensive GCM of Venus that faithfully reproduces the observed winds and circulation. However, Lee and Richardson (2010) did compute Lorenz energy budgets for a set of somewhat simplified GCM simulations of Venus-like circulations and so this

is what is presented here. However, these model simulations did not fully represent the radiative forcing of the atmosphere, including its diurnal cycle, nor did they fully capture the observed distribution of angular momentum in the simulations. So the budget shown here is likely to be inaccurate in some degree and missing some key processes. Nevertheless, the results are
560 worthy of attention as a first step towards a more accurate calculation.

Figure 12(c) presents the energy budget calculations for the spectral core case of Lee and Richardson (2010) with full damping, as being reasonably representative of this class of Venus model (without an explicit diurnal cycle). The energy reservoirs show a very different distribution of energy from either Earth or Mars, with most of the dynamically active energy in the atmosphere residing in the zonal and eddy kinetic energies, KZ and KE. The potential energy reservoirs, however, are
565 certainly not negligible and suggest a significant concentration of available potential energy in the AE reservoir. AZ is relatively small, though, as would be expected for an atmosphere with relatively weak meridional temperature gradients. Among the conversion terms, CZ is the largest, indicative of a strong, thermally-direct Hadley overturning circulation to maintain the very large KZ reservoir. The next largest conversion is CK, which evidently acts in the positive direction to convert from KZ to KE. This would suggest a dominance of barotropic processes, much as found in the idealised Earth-like simulations of Section 3 at
570 low values of Ω^* .

CA, on the other hand, also acts in a positive direction (from AZ to AE), indicating some alignment with baroclinically active processes and perhaps suggesting a mixed baroclinic/barotropic character to the dominant eddy processes. CE, however, acts in a negative direction, from KE to AE, which would suggest that mechanical exchanges between eddy potential and kinetic energies dominate over buoyancy driven instabilities. The one term that may be anomalous when compared with Venus
575 is the potential energy source/sink GE. This appears to be acting to remove energy from AE at a fairly strong rate. But this model takes no account of the diurnal cycle, and hence the input of thermal energy to the thermal tides. More realistic model simulations suggest that the thermal tides in and above the main cloud decks play an important role in maintaining the observed very strong atmospheric super-rotation (e.g. Sánchez-Lavega et al., 2017). It seems more likely, therefore, that a more complete simulation of the Venus circulation would include a large positive contribution to GE, much as indicated for Mars in Fig. 12(b).
580 This is a question that should be followed up urgently through analysis of more realistic model simulations.

4.3 Jupiter

Jupiter is the largest of a quite different class of planet within the Solar System, known as the gas giants, which also includes Saturn but also shares a number of dynamical aspects in common with Uranus and Neptune, known as the ice giants because of differences in composition and internal structure from Jupiter and Saturn (e.g. Guillot, 2005). As these names would suggest,
585 these planets are much larger than the Earth (Jupiter's mass is around 316 times larger than Earth's), with the gas giants largely composed of hydrogen and helium in proportions similar to the Sun. Both gas and ice giants therefore do not support a solid surface, unlike the terrestrial planets, but remain fluid to great depths until reaching a core of heavier elements close to the planetary centre. At a certain depth, however, the highly compressed fluid becomes electrically conducting and at even greater depths the hydrogen undergoes a phase change from a supercritical molecular fluid to a liquid metal (at least for Jupiter and
590 Saturn, though not for the so-called ice giants, Uranus and Neptune; e.g. see Guillot (2005)). All four gas and ice giant planets

orbit far from the Sun, with Jupiter (the innermost of these planets) in an orbit of mean radius 5.2 AU and orbital period of 11.86 years.

All four gas and ice giant planets are fully cloud covered, with clouds composed mainly of chemically reduced compounds such as ammonia ice, NH_4SH , H_2S and water (and CH_4 on Uranus and Neptune). Motions of these clouds reveal a very different circulation at the tops of their tropospheres from what is seen on Earth or the other terrestrial planets. Winds are predominantly zonal and arranged in systems of alternating eastward and westward jet streams, on scales smaller than the planetary radius. Cloud bands partly align with these zonal flows but are perturbed by waves and oval vortical eddies across a range of scales. Near the main cloud decks, incoming sunlight contributes significantly to the radiative energy budget. However, both Jupiter and Saturn (and Neptune) are significant net emitters of excess thermal radiation from a heat source in the deep interior, most likely representing residual thermal energy still escaping to space from the time of the original formation of these planets (e.g. Guillot, 2005). The action of incoming sunlight, upwelling heat from the deep interior and even the release of latent heat energy from the condensation of trace amounts of water in these planets have been suggested to play an important role in an atmospheric layer around 200-300 km thick, commonly known as the “weather layer” (e.g. Gierasch et al., 2000; Ingersoll et al., 2000). In the latter case, deep convective clouds and lightning are observed on Jupiter and Saturn which some estimates (Gierasch et al., 2000) suggest might carry as much as half of the upward thermal energy flux from the deep interior into the weather layer. The extent to which such convective storms energise large scale motions on Jupiter or Saturn remains controversial, however, since the apparent horizontal scale of such moist convective clusters is likely significantly smaller than the ~ 3000 km scale inferred by Young and Read (2017) for the excitation of Jupiter’s inverse kinetic energy cascade. The zonally dominated circulation in this weather layer is not thought to penetrate very deeply below the bottom of this layer, though recent measurements from the Juno mission suggest that the zonal wind pattern could penetrate as much as 3000 km below the visible clouds (around 4% of Jupiter’s radius; Kaspi et al. (2018)) though this may be still somewhat uncertain (e.g. see Kong et al. (2018)).

Despite their massive sizes, both Jupiter and Saturn are very rapid rotators with rotation periods around 10 hours. This would imply that differential motions within their fluid envelopes are highly likely to be strongly geostrophic. The observable regions of their atmospheres, above, within and somewhat below the main NH_3 and H_2O cloud decks, are found to be stably stratified, indicating the possibility of instability processes that have dynamical properties in common with those found in the Earth’s atmosphere and oceans. Their rapid rotation leads to values of the main Rossby radius of deformation, L_D , to be much smaller than the planetary radius a . This is reflected in the very small values of $\mathcal{B}u$ and \mathcal{R}_{OT} indicated for Jupiter’s cloud tops in Table 2. Together with the very large value of \mathcal{T}_R , this would seem to suggest that geostrophic forms of barotropic, and possibly baroclinic, instabilities are likely to occur in the weather layers of these planets.

4.3.1 PV structures: observations

The occurrence of such instabilities, however, depends crucially on the distribution of PV within the weather layers of these planets. As with the other planets under discussion here, Jupiter’s atmosphere needs to satisfy the CSP necessary condition for instability somehow. Without a solid surface, however, CSP criterion (iii) cannot be satisfied, unlike on Earth or Mars. Such

625 an argument was used by Gierasch et al. (1979a) to suggest only a minor role at most for baroclinic instabilities on Jupiter. But as subsequently noted by Conrath et al. (1981), the possibilities of satisfying the CSP condition through criteria (i) or (ii) remain, especially if Jupiter or Saturn possess a strong tropopause which can act as an internal interface that supports a horizontal thermal gradient. The analysis of Conrath et al. (1981) noted that, in the absence of significant horizontal shear and with a lower boundary that was essentially isentropic and weakly stratified, westward jets could satisfy the CSP condition
630 through criterion (i) (or (ii) if the tropopause is considered as an upper boundary or interface). Their calculations indicated that such an "inverted Charney" form of baroclinic instability could have relatively fast growth rates and even lead to up-gradient momentum fluxes for instabilities on a small enough horizontal scale.

The vorticity structure of the zonal mean flow at the cloud tops has been measured since the time of the Voyager spacecraft encounters with Jupiter in 1979. By tracking features in the ubiquitous ammonia cloud decks in successive images taken by the
635 spacecraft, the pattern of eastward and westward jet streams could be derived as a profile in latitude from which the vorticity and its northward gradient could be obtained (e.g. Ingersoll et al., 1981; Salyk et al., 2006; Galperin et al., 2014). These profiles indicate that the northward gradient of *absolute* vorticity robustly changes sign with latitude in a number of places, typically around the peaks of westward jets, leading to some suggestions that these jets may be prone to barotropic instabilities since they would appear to satisfy criterion (i) for the CSP instability condition (Ingersoll et al., 1981). The stretching term associated
640 with the thermal structure of the flow may also be important, leading to some suggestions (e.g. Ingersoll and Cuzzi, 1969; Scott and Dunkerton, 2017) that vigorous lateral mixing of PV could result in a monotonic, staircase-like variation of Q with latitude which would be unconditionally stable to shear instabilities. Such a process that might hold the atmosphere close to a state of marginal instability would effectively constitute a form of *barotropic adjustment* by analogy with Stone's baroclinic adjustment hypothesis (Stone, 1978).

645 Observational studies that have combined wind measurements from cloud motions with remotely sensed temperature retrievals (Read et al., 2006, 2009a; Antuñano et al., 2019) and a geostrophic balance assumption for both Jupiter and Saturn, however, also indicate significant reversals of $\partial Q/\partial y$. Figure 15(a) shows an example of a section of $\partial Q/\partial y$ in latitude and height in the southern hemisphere of Jupiter, based on data from the Cassini spacecraft (Read et al., 2006). This clearly shows patches where $\partial Q/\partial y < 0$ overlying westward jets but persisting over significant ranges in altitude. $\partial Q/\partial y$ even appears to
650 change sign in the vertical direction in some locations in Jupiter's stratosphere, although these observations should be treated with caution since the reconstruction of the flow at high altitudes requires the integration of the geostrophic thermal wind shear relation which can amplify the effects of measurement noise and uncertainties, especially in quantities that require spatial differentiation (e.g. see Read et al., 2009a). Being based on nadir observations, the retrieved thermal measurements also have rather limited vertical resolution. So it cannot be ruled out that structures associated with sharp thermal gradients along the
655 lines suggested by Scott and Dunkerton (2017) might exist that would eliminate such reversals in sign of $\partial Q/\partial y$.

The distribution of Q beneath Jupiter's cloud tops, however, is virtually unknown because of a dearth of detailed measurements of winds, temperatures or composition. But there seems no reason *a priori* to conclude that internal changes of sign of $\partial Q/\partial y$ in the deep troposphere, or of $\partial T/\partial y$ at the tropopause, cannot exist, and that therefore the CSP necessary condition for baroclinic or barotropic instability could be satisfied through criteria (i) or (ii).




Figure 15. (a) Zonal mean section of $\partial Q / \partial y$ ($\text{s}^{-1} \text{m}^{-1}$) and (b) zonal mean zonal wind (m s^{-1}) in the southern hemisphere of Jupiter, derived from Cassini observations by Read et al. (2006).

660 4.3.2 PV structures: GCMs

A number of studies have appeared recently in which fully three-dimensional, time-dependent GCMs have been developed of Jupiter's or Saturn's weather layers which seek to capture the thermal and vorticity structures of the flow around and beneath the cloud tops (e.g. Liu and Schneider, 2010; Lian and Showman, 2008, 2010; Palotai et al., 2014; Young et al., 2019a, b; Spiga et al., 2020). These models have included the effects of solar heating in the stratosphere and upper troposphere and upwelling
665 internal heating from the deep interior, and have been capable of capturing a number of realistic features such as the equatorial prograde jets and multiple eddy-driven zonal jets at mid-latitudes.

Figure 16 shows a cross-section of $\partial Q / \partial y$ and \bar{u} from a GCM simulation of Jupiter's weather layer by Young et al. (2019a) (their run B, with a horizontal resolution of 512×256 points in longitude and latitude). As with the observations, this shows a generally clear correlation between \bar{u} and $\partial Q / \partial y$ in the upper troposphere but with some sign reversals of $\partial Q / \partial y$ in both
670 the horizontal and vertical in relation to both westward and eastward jets in several cases. This would seem to confirm that the simulated flow sustains the necessary conditions for baroclinic instability through CSP criterion (i) as discussed above.




Figure 16. Zonal mean latitude-height section of $\partial Q/\partial y$ ($\text{s}^{-1} \text{m}^{-1}$) and zonal mean zonal wind (m s^{-1}) in a numerical simulation of Jupiter’s weather layer, from Young et al. (2019a). Data correspond to run B of Young et al. (2019a) and were averaged over that last 100 days of the equilibrated run. Shading represents $\partial Q/\partial y$ and the line contours represent zonal mean zonal velocity \bar{u} , shown for -10, -5, +5 and +10 m s^{-1} in grey and 0 m s^{-1} in black.

4.3.3 Energy Budget

A preliminary Lorenz energy budget, computed using an updated, higher-resolution version of the Jupiter weather layer model described by Young et al. (2019a), is shown in Fig. 12(d). This represents the model’s attempt to capture the energetics of a Jupiter-like weather layer between the pressure levels of 18 bars and 10 hPa. The energy reservoirs reflect a distribution of potential and kinetic energy typical of a rapidly rotating, stratified atmosphere, with most of the stored energy residing in the zonal mean potential energy AZ. Smaller amounts are found in the KZ and AE reservoirs with KE the weakest, though still containing much more energy per unit area than the corresponding reservoirs for Earth.

Energy conversions, however, are remarkably Earth-like in their relative sizes and directions, with CA and CE as the largest conversion rates in the sense expected for baroclinic instabilities. This confirms the indication from the PV structure that anticipated that baroclinic instabilities could occur and play a significant role in the circulation. CK acts to maintain the zonal mean flow through transfers from KE, at a vertically integrated rate (0.61 W m^{-2}) that is not unreasonable with regard to the observed transfer rate near the cloud tops (e.g. Ingersoll et al., 1981; Salyk et al., 2006; Galperin et al., 2014). CZ acts in the thermally direct sense but is relatively weak, reflecting the comparatively minor role played by zonally symmetric overturning

685 circulations in the global circulation. The largest terms overall are CA, GZ and GE, indicating some very strong forcing of
the zonal mean potential energy field and conversion to eddy potential energy, much of which is then extracted by diabatic
processes. A complication is that this model simulation includes both latent heat fluxes and exchanges associated with a moist
convection parameterization, which may not be fully accounted for in the conventional Lorenz energy budget approach. But
the overall structure of this cycle strongly suggests an important role for baroclinic instability, at least in this model simulation
690 of Jupiter's weather layer.

5 Conclusions

Throughout this review it has been clear that baroclinic and/or barotropic instabilities play a key role in generating some of the
most energetic eddies and waves in most rotating, stratified systems, including systems that are very different from the Earth's
atmosphere and oceans. The distinction between baroclinic and barotropic instabilities is, *in certain respects, an artificial one,*
695 *at least concerning the constraints deriving from the vorticity structure of the flow. The principal distinction is a real one,*
however, based largely upon whether the dominant conversion of energy from an initial basic state is from its stored potential
or kinetic energy, *as revealed in their Lorenz energy budgets.* But the CSP necessary condition for instability *applies to* both
mechanisms, which are governed by similar vorticity constraints. In practice, eddy formation in realistic systems with both
vertical and horizontal shear could entail contributions from both potential and kinetic energy stored in the basic state, so that it
700 is natural to consider either mechanism as different facets of a common process. This suggests the value of a unifying approach,
although a number of outstanding issues remain.

5.1 Stability theory

The body of stability theory continues to provide important and useful guidance to help predict the onset of instabilities in
various practical geophysical and astrophysical contexts, and in particular to establish conditions for when instabilities will
705 not occur. But there are still many situations that are not fully worked out in complete generality and most published stability
criteria have shortcomings that fail to take certain factors into account. After more than 50 years, the CSP instability condition
is still widely used to assess the stability of zonal flows in atmospheres and oceans, even though it is based on linearised
quasi-geostrophic theory for perturbations to a purely axisymmetric basic state. As an instability condition it only represents a
necessary condition but not a sufficient one to guarantee instability if it is met.

710 The more sophisticated stability theorem of Arnol'd (1966) and more recent extensions (e.g. see Mu and Wu, 2001, for a
review) offer a more refined approach which takes some account of nonlinearity and non-zonal structure, although with some
non-trivial restrictions. It is most well developed for quasi-geostrophic flows but a full treatment, e.g. of the primitive equations
of dynamical meteorology, is still lacking. Much of the most well developed theory also applies strictly to a Hamiltonian
system, neglecting forcing and dissipation processes that may maintain the flow in a quasi-equilibrium state. Some exceptions
715 exist (e.g. Andrews, 1984a, b) but they are still quite rare.

5.2 Equilibration and adjustment

Linear stability theory generally applies to the behaviour of small amplitude perturbations, but real flows are more typically observed when any incipient instabilities have grown to large amplitude and saturated. Precisely how equilibration is achieved may depend strongly on the structure of the flow and how it is maintained and/or dissipated. Even weakly nonlinear theories (e.g. Pedlosky, 1970, 1971; Drazin, 1970) show that growing instabilities act to reduce the vigour of the instability and its growth rate by modifying the structure of the initial background flow to make it less strongly unstable. For barotropically or baroclinically unstable flows, this commonly entails a combination of reducing the strength of any gradients of PV or buoyancy in the base state via a pattern of eddy fluxes, but the structure of the end state may depend on a variety of factors and constraints, including the strength and distribution of any forcing.

Where the forcing and dissipation is relatively weak, such that the characteristic timescale τ_H for the system to respond to changes in forcing is much longer than any advective overturning timescale $\tau_A \sim L/U$, fluid motions may be able effectively to nearly neutralise the instability so that, on timescales $O(\tau_H)$ or longer, the flow stays close to a marginally unstable state. This scenario may offer the possibility of parameterizing the effects of strong barotropic or baroclinic instabilities on large-scale flows through forms of barotropic or baroclinic adjustment (by analogy with convective adjustment). However, simply posing a requirement to seek a marginally unstable state is usually insufficient to determine or predict from first principles a unique equilibrated state unless additional constraints or information are available, such as an applicable optimisation principle (e.g. for entropy production, potential enstrophy or a generalised free energy, e.g. Lucarini (2009), Hollerbach et al. (2004), Bouchet and Venaille (2012)).

Such an adjustment principle may have useful applicability to the interpretation of [at least some](#) observed flow systems, however. Dowling (1993) and Stamp and Dowling (1993), for example, have suggested that the zonal mean winds in the observable cloud level atmospheres of both Jupiter and Saturn may approach a state that is just marginally unstable with respect to a form of Arnol'd's second stability theorem, a hypothesis that seemed to be confirmed in more quantitative observations of potential vorticity (Read et al., 2006, 2009a). This was used by Read et al. (2009b) to infer the existence of a unique reference frame on each planet which [represents the the frame in which the gravest/deepest Rossby wave trains are just able to propagate at the same phase speed and thus couple together and interact via over-reflection. This frame](#) corresponded closely on Jupiter to that of its magnetic field and deep interior. For Saturn, however, it differed significantly from the magnetic rotation periods inferred from measurements from the Voyager or Cassini spacecraft, but agreed more closely with alternative estimates of Saturn's interior rotation from measurements of its gravity field or perturbations in its rings (Anderson and Schubert, 2007; Mankovitch et al., 2019).

5.3 Baroclinic atmospheres

For baroclinic atmospheres, such as that of the Earth, the possibility persists that its zonal mean circulation is at least partly controlled by a form of baroclinic adjustment (Stone, 1978), although debates continue to the present day as to whether this is a valid assumption and/or precisely what principle is being optimised during equilibration (e.g. Zurita-Gotor and Lindzen,

2007; Schneider, 2007). Flaws in Stone’s original argument, that hinges on a tendency for the flow to deviate only slightly
 750 from a marginally unstable state, were discussed above, leading to alternative suggestions that the flow evolves towards a state
 with reduced nonlinear interactions (e.g. Schneider and Walker, 2006). Such a hypothesis is consistent with the observation
 that equilibrated flows in both the laboratory and in the atmospheres of Earth and some other planets exhibit the property that
 potential vorticity $Q \simeq Q(\psi)$.

The tendency shown in both laboratory experiments and simple GCM simulations for the total meridional heat transport
 755 (eddy plus zonal mean) to be largely independent of rotation rate (see Sections 2.2 and 3) over the range in $\mathcal{B}u$ and/or $\mathcal{R}o_T$
 from $O(1)$ to ~ 0.05 is somewhat remarkable. It is apparently due to a mutual compensation between decreasing heat transport
 by the zonal mean flow and increasing eddy heat transport with Ω until the latter reaches a peak around $\mathcal{R}o_T \simeq 0.1$. The
 resulting impact on the large-scale zonal mean flow is for the thermal structure to remain fairly static over this range of Ω ,
 which would imply that the mean slope of the isentropes in the latitude-height plane at mid-latitudes also remains largely
 760 independent of Ω .

This is consistent with a roughly constant value of the criticality parameter

$$\xi = \frac{f}{\beta} \frac{\partial \bar{\theta} / \partial y}{\Delta \theta_v} = - \frac{f}{\beta H} \frac{\partial \bar{\theta} / \partial y}{\partial \bar{\theta} / \partial z}, \quad (16)$$

where $\Delta \theta_v$ is a vertical contrast in potential temperature and H is a vertical length scale, often taken to be the pressure scale
 height of an atmosphere. ξ is thus a measure of the mean slope of the isentropes, which many studies have indicated remains
 765 close to $O(1)$ over a wide range of parameters centred around those for Earth (e.g. Stone, 1978; Schneider and Walker, 2006;
 Zurita-Gotor and Lindzen, 2007). A number of different implications for $\xi \simeq 1$ have been noted, including an absence of a
 significant inverse kinetic energy cascade, since ξ can also be interpreted as measuring the squared ratio of the Rhines scale

$$L_R = \left(\frac{U}{\beta} \right)^{1/2}, \quad (17)$$

(based on the zonal mean thermal wind scale $U \sim U_T$) which can estimate the large-scale limit to the main inverse cascade, to
 770 the Rossby deformation radius L_D (although L_R is more conventionally estimated using $U \sim (EKE)^{1/2}$ which can differ sig-
 nificantly from U_T under some conditions). This would also suggest a suppression of nonlinear wave-wave interactions, which
 would be consistent with equilibration towards a state in which $\mathbf{u} \cdot \nabla Q \simeq 0$, implying that $Q \simeq Q(\psi)$ in a quasi-geostrophic
 flow (where ψ is the horizontal stream function). Something close to this configuration of Q and ψ occurs within the regular
 baroclinic wave regime in rotating annulus experiments (e.g. Read et al., 1986), and even the Earth’s atmosphere seems to
 775 approach such a state to some extent (e.g. Butchart et al., 1989).

Recent work (e.g. Vallis, 1988; Zurita-Gotor, 2008; Jansen and Ferrari, 2012) has questioned whether the preferred value of
 ξ is universally around $O(1)$ for all situations, noting that conditions in the oceans, for example, may favour equilibrated values
 of $\xi \gg 1$ (Jansen and Ferrari, 2012, 2013). Jansen and Ferrari (2013) suggests that the equilibrated value of ξ in a baroclinic
 atmosphere is likely to be set by a balance between differential heating through radiative forcing (with radiative adjustment
 780 timescale τ_R) and eddy heat fluxes, parameterised via a diffusivity (for PV), D . This leads to an estimate of the isentropic slope
 $s \sim H / \sqrt{\tau_R D}$ along a lengthscale $\ell_{diff} \sim \sqrt{\tau_R D}$, over which eddies can transport PV and heat within the radiative timescale

τ_R , and thus

$$\xi \sim \frac{sa}{H} \sim \frac{a}{\sqrt{\tau_R D}} \quad (18)$$

(e.g. Jansen and Ferrari, 2012, 2013). If $\ell_{diff} \ll a$, therefore, values of $\xi \gg 1$ may occur, whereas if $\ell_{diff} \gtrsim a$, then heat transfer by eddies is efficient enough to homogenise potential temperature horizontally, relaxing the isentropic slope such that $\xi \leq 1$.

Taking a broader view across a wide range of parameter space and different physical systems, however, the discussion above in Sections 2 – 3 suggests that a more complete explanation for many of the recently published results on criticality and baroclinic adjustment may be related to the extent to which variations in heat transports, not only due to eddies but also to the zonal mean circulation, may mutually compensate each other as parameters are varied so that the total heat transport is maintained at a nearly constant rate. This compensation may be most effective within the typical range $0.05 < \mathcal{R}o_T \leq 1$, within which eddy heat transports increase with decreasing $\mathcal{R}o_T$ while zonal mean transport decreases. Beyond this limit (i.e. $\mathcal{R}o_T < 0.05$), both components of heat transport decrease as $\mathcal{R}o_T$ decreases, and the zonal mean thermal structure might be expected to relax gradually towards the radiative-convective equilibrium state as the transport by the zonal mean circulation becomes small compared to the eddy transport and the eddy diffusivity, D , decreases. The very high rotation rate limit of ξ would then depend upon the slope of the isentropes in this radiative-convective equilibrium state, but could well lie close to $O(1)$ for Earth-like atmospheric conditions. For other systems, however, such as in the Earth’s oceans, the criticality ξ has been suggested to be $\gg 1$ (e.g. Vallis, 1988; Jansen and Ferrari, 2012), and a similar criticality may well apply in the atmospheres of the gas giants (Jupiter and Saturn), where D is likely very small and the natural slope of isentropes in the convectively active troposphere is very steep. The existence or otherwise of a genuine baroclinic adjustment regime, however, depends upon the existence of a range of parameters over which eddy heat transports strengthen with Ω so can compensate for changes in the zonal mean transport. While this seems to be plausible for Earth-like atmospheres, it may not be completely general and should be investigated more thoroughly in future work.

5.4 Planetary atmospheres

The possible existence of both baroclinic and barotropic adjustment regimes immediately raises the question as to whether any of the known planetary atmosphere circulations are actually in one of these regimes. Given the limits on $\mathcal{R}o_T$ for the baroclinic adjustment regime as described here, the Earth itself would seem to lie close to the high Ω margins of this regime, which would be consistent with recent work highlighting its possible effects on the circulation. Mars, with a value $\mathcal{R}o_T \simeq 0.2$ (see Table 2) would also appear at first sight to be firmly within the baroclinic adjustment regime. However, it might not fulfill the other condition for the existence of a baroclinic adjustment regime, namely, that forcing timescales $\tau_S \gg \tau_A$. Radiative adjustment timescales on Mars are typically only around $\tau_R \simeq 1$ day, which is comparable to $\tau_A = L/U$ based on typical horizontal velocities and the planetary radius. Radiative forcing is therefore relatively much stronger on Mars than on Earth, which might suggest from Eq (18) that baroclinic adjustment would lead to an adjusted state with $\xi > 1$. Under these strongly forced/damped conditions, however, flow configurations that appear to be vigorously unstable, based on criteria derived for

815 non-dissipative flows, may actually be relatively stable in practice. This was illustrated recently in the work of Seviour et al. (2017), for example, who noted that zonal mean PV at high latitudes on Mars often took the form of a persistent annular band, across which were strong reversals in the sign of $\partial Q/\partial y$. Seviour et al. (2017) were able to reproduce this behaviour in a simple model in which zonal mean forcing was applied by relaxing to a similar PV distribution with a short relaxation timescale. Baroclinic adjustment concepts are therefore unlikely to apply to Mars in any useful sense.

820 The cases of slowly rotating planets, such as Venus or Titan in the Solar System, would also seem to place them outside of the limits on \mathcal{R}_{OT} for baroclinic adjustment to be effective, [since heat transports are then dominated by the zonal mean circulation and not by eddies](#). Their large scale thermal structure is typically characterised by very weak horizontal thermal gradients compared to their stratification, so isentropic slopes are very small. This would indicate that $\xi \ll 1$ with relatively weak eddy heat transports, at least on the planetary scale. The strong super-rotation of the atmospheres of Venus (and Titan?),
825 however, seems to allow for a role for baroclinic instabilities in atmospheric layers that rotate fast enough to be in a quasi-geostrophic regime, with local values of $\mathcal{R}_{OT} \sim 0.1$ (see Table 2). Radiative relaxation timescales near the cloud tops of Venus are also relatively long compared with advective overturning timescales, so baroclinic adjustment effects might be relevant to the circumpolar regions of Venus’s cloudy layers, a possibility that deserves further investigation.

Jupiter would seem to be well outside its baroclinic adjustment regime, based simply on its values of \mathcal{R}_{OT} and $\mathcal{B}u$ (see Table
830 2). Recent work has demonstrated that Jupiter exhibits a substantial upscale cascade of kinetic energy at its cloud tops, both spectrally local and non-local (Young and Read, 2017). This is consistent with estimates that suggest that L_R is significantly larger than L_D on Jupiter (Galperin et al., 2014), indicating that Jupiter’s weather layer is significantly super-critical with an effective value for ξ of around 5. This would seem to be because the relevant scale for U in L_R is much larger than the thermal wind scale U_T , with much of the kinetic energy at Jupiter’s cloud tops residing in its zonal jets. Such jets may well be
835 predominantly barotropic in structure, at least within the weather layer.

It is perhaps unsurprising, therefore, that barotropic adjustment seems to be a useful concept in the context of the gas giant planets, Jupiter and Saturn, as discussed above, with the circulation in the weather layers remaining close to a marginally unstable state, as determined by the Arnol’d II criterion. It seems reasonable that the weather layers of Uranus and Neptune might also be in a similar regime, which could readily be investigated in both observations and models given the availability
840 of suitable data. The quality and coverage of the original data from the Voyager encounter may not be adequate to verify these ideas by themselves, although the accuracy and latitude coverage of cloud motion tracking has improved significantly in recent years from Earth-based (ground and space) telescopes (e.g. Garcia-Melendo and Sánchez-Lavega, 2001). These issues should certainly be addressed in the context of future missions currently being planned for the ice giants.

In the context of the ever expanding discoveries of extrasolar planets, it seems highly likely that many of them will host
845 atmospheres within which baroclinic or barotropic instabilities may play an active role. Future investigations of their possible climates and atmospheric circulations will therefore need to take fairly full account of the influence of these instabilities, for which equilibration and adjustment concepts may surely prove useful in at least some cases.

Data availability. Data on the Martian atmosphere can be obtained from the Mars Climate Database website <http://www-mars.lmd.jussieu.fr/mars/access.html>. Data from the model simulations of Young et al. (2019a) can be obtained from Oxford University's Research Archive at <https://doi.org/10.5287/bodleian:PyYbbxpk2>. Data from laboratory experiments reported in Section 2 can be obtained from the authors on request.

Author contributions. PLR supervised much of the research reported from the University of Oxford, analysed some of the laboratory data and was the lead author of the paper. NL, FT-V, YW and RY carried out computations and analysis for model simulations and DK computed the energy budget reported in Section 4.3. Other authors also contributed to writing the text.

855 *Competing interests.* We confirm that we have no competing interests arising from the work described in this article.

Acknowledgements. PLR, FT-V and RMBY acknowledge support from the UK Science and Technology Research Council during the course of this research under grants ST/K502236/1, ST/K00106X/1 and ST/I001948/1. This article originated from a lecture prepared by PLR for the Summer School on Waves, Instability and Turbulence in Geophysical and Astrophysical Flows in Cargèse, Corsica in July 2019.

References

- 860 Aguiar, A. C. B., Read, P. L., Wordsworth, R. D., Salter, T., and Yamazaki, Y. H.: A laboratory model of Saturn's North Polar Hexagon, *Icarus*, 206, 755–763, 2010.
Anderson, J. D. and Schubert, G.: Saturn's gravitational field, internal rotation, and interior structure, *Science*, 317, 1384–1387, 2007.
Andrews, D. G.: On the existence of nonzonal flows satisfying sufficient conditions for stability, *Geophys. Astrophys. Fluid Dyn.*, 28, 243–256, 1984a.
- 865 Andrews, D. G.: On the stability of forced non-zonal flows, *Quart. J. R. Meteor. Soc.*, 110, 657–662, 1984b.
Antuñano, A., del Río-Gaztelurrutia, T. and. Sánchez-Lavega, A., Read, P. L., and Fletcher, L. N.: Potential Vorticity of Saturn's Polar Regions: Seasonality and Instabilities, *J. Geophys. Res.*, 124, 186–201, <https://doi.org/10.1029/2018JE005764>, 2019.
Arnol'd, V. I.: On an a priori estimate in the theory of hydrodynamical stability. [In Russian.], *Izv. Vyssh. Ucheb. Zaved. Matematika*, 54, 3–5, 1966.
- 870 Barnes, J. R.: Linear baroclinic instability in the Martian atmosphere, *J. Atmos. Sci.*, 41, 1536–1550, 1984.
Bell, M. J. and White, A. A.: The stability of internal baroclinic jets: some analytical results, *J. Atmos. Sci.*, 45, 2571–2590, 1988.
Boer, G. J. and Lambert, S.: The energy cycle in atmospheric models, *Clim. Dyn.*, 30, 371–390, 2008.
Bouchet, F. and Venaille, A.: Statistical mechanics of two-dimensional and geophysical flows, *Phys. Rep.*, 515, 227–295, <https://doi.org/10.1016/j.physrep.2012.02.001>, 2012.
- 875 Bowden, M.: An experimental investigation of heat transfer in rotating fluids, Ph.D. thesis, Durham University, UK, 1961.
Branscombe, L. E.: The Charney Baroclinic Instability Problem: Approximate Solutions and Modal Structures, *J. Atmos. Sci.*, 40, 1393–1409, 1983.
Butchart, N., Haines, K., and Marshall, J. C.: A theoretical and diagnostic study of solitary waves and atmospheric blocking, *J. Atmos. Sci.*, 46, 2063–2078, 1989.
- 880 Charney, J. G. and Stern, M. E.: On the Stability of Internal Baroclinic Jets in a Rotating Atmosphere, *J. Atmos. Sci.*, 19, 159–172, 1962.
Conrath, B. J., Gierasch, P. J., and Nath, N.: Stability of zonal flows on Jupiter, *Icarus*, 48, 256–282, 1981.
Dowling, T.: A relationship between potential vorticity and zonal wind on Jupiter, *J. Atmos. Sci.*, 50, 14–22, 1993.
Dowling, T.: Dynamics of Jovian atmospheres, *Ann. Rev. Fluid Mech.*, 27, 293–334, 1995.
Drazin, P. G.: Non-linear baroclinic instability of a continuous zonal flow, *Quart. J. R. Meteor. Soc.*, 96, 667–676, 1970.
- 885 Eady, E. T.: Long Waves and Cyclone Waves, *Tellus*, 1, 33–52, 1949.
Flór, J.-B., Scolan, H., and Gula, J.: Frontal instabilities and waves in a differentially rotating fluid, *J. Fluid Mech.*, 685, 532–542, <https://doi.org/10.1017/jfm.2011.338>, 2011.
Fraedrich, K., Kirk, E., Luksch, U., and Lunkeit, F.: The Portable University Model of the Atmosphere (PUMA): Storm track dynamics and low frequency variability, *Meteorol. Zeitschrift*, 14, 735–745, 2005.
- 890 Frisius, T., Lunkeit, F., Fraedrich, K., and James, I. N.: Storm-track organization and variability in a simplified global circulation model, *Quart. J. R. Meteor. Soc.*, 124, 1019–1043, 1998.
Früh, W.-G. and Read, P. L.: Experiments on a barotropic rotating shear layer. Part 1. Instability and steady vortices, *J. Fluid Mech.*, 383, 143–173, 1999a.
Früh, W.-G. and Read, P. L.: Flow field and point velocity measurements in a barotropically unstable shear layer, *Phys. Chem. Earth (B)*, 24, 461–466, 1999b.

- Galperin, B., Young, R., Sukoriansky, S., Dikovskaya, N., Read, P., Lancaster, A., and Armstrong, D.: Cassini observations reveal a regime of zonostrophic macroturbulence on Jupiter, *Icarus*, 229, 295–320, 2014.
- Garcia-Melendo, E. and Sánchez-Lavega, A.: A study of the stability of Jovian zonal winds from HST images: 1995–2000, *Icarus*, 152, 316–330, 2001.
- 900 Geisler, J., Pitcher, E., and Malone, R.: Rotating-Fluid experiments with an atmospheric general circulation model, *J. Geophys. Res.*, 88, 9706–9716, <https://doi.org/10.1029/JC088iC14p09706>, 1983.
- Gierasch, P. J., Ingersoll, A. P., and Pollard, D.: Baroclinic instabilities in Jupiter’s zonal flow, *Icarus*, 40, 205–212, 1979a.
- Gierasch, P. J., Thomas, P., French, R., and Veverka, J.: Spiral clouds on Mars: a new atmospheric phenomenon, *Geophys. Res. Lett.*, 6, 405–408, 1979b.
- 905 Gierasch, P. J., Ingersoll, A. P., Banfield, D., Ewald, S. P., Helfenstein, P., Simon-Miller, A., Vasavada, A., Breneman, H. H., Senske, D. A., and the Galileo Imaging Team: Observation of moist convection in Jupiter’s atmosphere, *Nature*, 403, 628–630, 2000.
- Guillot, T.: The Interiors of Giant Planets: Models and Outstanding Questions, *Ann. Rev. Earth Planet Sci.*, 33, 493–530, <https://doi.org/10.1146/annurev.earth.32.101802.120325>, 2005.
- Gula, J., Plougonven, R., and Zeitlin, V.: Ageostrophic instabilities of fronts in a channel in a stratified rotating fluid, *J. Fluid Mech.*, 627, 485–507, <https://doi.org/10.1017/S0022112009006508>, 2009.
- 910 Hayashi, Y.-Y. and Young, W. R.: Stable and unstable shear modes of rotating parallel flows in shallow water, *J. Fluid Mech.*, 184, 477–504, 1997.
- Hide, R.: Some laboratory experiments on free thermal convection in a rotating fluid subject to a horizontal temperature gradient and their relation to the theory of the global atmospheric circulation, in: *The Global Circulation of the Atmosphere*, edited by Corby, G. A., pp. 196–221, Royal Meteorological Society, London, 1969.
- 915 Hide, R. and Mason, P. J.: Sloping convection in a rotating fluid, *Adv. Phys.*, 24, 47–100, 1975.
- Hide, R. and Titman, C. W.: Detached shear layers in a rotating fluid, *J. Fluid Mech.*, 29, 39–60, 1967.
- Hollerbach, R., Futterer, B., More, T., and Egbers, C.: Instabilities of the Stewartson layer Part 2. supercritical mode transitions, *Theoret. Comput. Fluid Dyn.*, 19, 197–204, <https://doi.org/10.1007/s00162-004-0125-5>, 2004.
- 920 Hunt, B. G.: The influence of the Earth’s rotation rate on the general circulation of the atmosphere, *J. Atmos. Sci.*, 36, 1392–1408, 1979.
- Ingersoll, A., Gierasch, P., Banfield, D., and Vasavada, A.: Moist convection as an energy source for the large-scale motions in Jupiter’s atmosphere. Galileo Imaging Team, *Nature*, 403, 630–2, <https://doi.org/10.1038/35001021>, 2000.
- Ingersoll, A. P. and Cuzzi, J. N.: Dynamics of Jupiter’s cloud bands, *J. Atmos. Sci.*, 26, 981–985, 1969.
- Ingersoll, A. P., Beebe, R. F., Mitchell, J. L., Garneau, G. W., Yagi, G. M., and Müller, J.-P.: Interaction of eddies and mean zonal flow on 925 Jupiter as inferred from Voyager 1 and 2 images, *J. Geophys. Res.*, 86, 8733–8743, 1981.
- James, I. N.: *Introduction to Circulating Atmospheres*, Cambridge University Press, Cambridge, UK, 1994.
- Jansen, M. and Ferrari, R.: Macroturbulent Equilibration in a Thermally Forced Primitive Equation System, *J. Phys. Oceanogr.*, 69, 695–713, 2012.
- Jansen, M. and Ferrari, R.: Equilibration of an Atmosphere by Adiabatic Eddy Fluxes, *J. Atmos. Sci.*, 70, 2948–2962, 2013.
- 930 Kaspi, Y. and Showman, A. P.: Atmospheric dynamics of terrestrial exoplanets over a wide range of orbital and atmospheric parameters, *Astrophys. J.*, 804, 60, 2015.

- Kaspi, Y., Galanti, E., Hubbard, W. B., Stevenson, D. J., Bolton, S. J., Iess, L., Guillot, T., Bloxham, J., Connerney, J. E. P., Cao, H., Durante, D., Folkner, W. M., Helled, R., Ingersoll, A. P., Levin, S. M., Lunine, J. I., Miguel, Y., Militzer, B., Parisi, M., and Wahl, S. M.: Jupiter's atmospheric jet streams extend thousands of kilometres deep, *Nature*, 555, 223–226, <https://doi.org/10.1038/nature25793>, 2018.
- 935 Kong, D., Zhang, K., Schubert, G., and Anderson, J. D.: Origin of Jupiter's cloud-level zonal winds remains a puzzle even after Juno, *PNAS*, 115, 8499–8504, <https://doi.org/10.1073/pnas.1805927115>, 2018.
- Lebonnois, S., Hourdin, F., Eymet, V., Cresspin, A., Fournier, R., and Forget, F.: Superrotation of Venus' atmosphere analyzed with a full general circulation model, *J. Geophys. Res.*, 115, E06 006, <https://doi.org/10.1029/2009JE003458>, 2010.
- Lebonnois, S., Lee, C., Yamamoto, M., Dawson, J., Lewis, S. R., Mendonca, J., Read, P., Parish, H. F., Schubert, G., Bengtsson, L., Grin-
 940 spoon, D., Limaye, S. S., Schmidt, H., Svedhem, H., and Titov, D. V.: Models of Venus' atmosphere, in: *Towards Understanding the Climate of Venus: Applications of Terrestrial Models to Our Sister Planet*, edited by Bengtsson, L., Bonnet, R.-M., Grinspoon, D., Koumoutsaris, S., Lebonnois, S., and Titov, D., pp. 129–156, Springer, New York, 2013.
- Lebonnois, S., Sugimoto, N., and Gilli, G.: Wave analysis in the atmosphere of Venus below 100-km altitude, simulated by the LMD Venus GCM, *Icarus*, 278, 38–51, <https://doi.org/10.1016/j.icarus.2016.06.004>, 2016.
- 945 Lee, C. and Richardson, M. I.: A general circulation model ensemble study of the atmospheric circulation of Venus, *J. Geophys. Res.*, 115, E04 002, <https://doi.org/10.1029/2009JE003490>, 2010.
- Lewis, G. M. and Nagata, W.: Linear stability analysis for the differentially heated rotating annulus, *Geophys. Astrophys. Fluid Dyn.*, 98, 129–152, 2004.
- Lewis, G. M., Périnet, N., and van Veen, L.: Primary flow transitions in the baroclinic annulus: Prandtl number effects, in: *Modelling Atmospheric and Oceanic Flows: Insights from Laboratory Experiments and Numerical Simulations*, edited by von Larcher, T. and Williams, P. D., pp. 45–60, American Geophysical Union and Wiley, Hoboken, NJ, USA, 2015.
- 950 Lewis, S. R., Collins, M., Read, P. L., Forget, F., Hourdin, F., Fournier, R., Hourdin, C., Talagrand, O., and Huot, J.-P.: A Climate Database for Mars, *J. Geophys. Res.*, 104, 24 177–24 194, 1999.
- Lewis, S. R., Dawson, J., Lebonnois, S., and Yamamoto, M.: Modeling efforts, in: *Towards Understanding the Climate of Venus: Applications of Terrestrial Models to Our Sister Planet*, edited by Bengtsson, L., Bonnet, R.-M., Grinspoon, D., Koumoutsaris, S., Lebonnois, S., and
 955 Titov, D., pp. 102–128, Springer, New York, 2013.
- Lewis, S. R., Mulholland, D. P., Read, P. L., Montabone, L., Wilson, R. J., and Smith, M. D.: The solsticial pause on Mars: 1. A planetary wave reanalysis, *Icarus*, 264, 456–464, <https://doi.org/10.1016/j.icarus.2015.08.039>, 2016.
- Lian, Y. and Showman, A. P.: Deep jets on gas-giant planets, *Icarus*, 194, 597–615, <https://doi.org/10.1016/j.icarus.2007.10.014>, 2008.
- 960 Lian, Y. and Showman, A. P.: Generation of equatorial jets by large-scale latent heating on the giant planets, *Icarus*, 207, 373–393, <https://doi.org/10.1016/j.icarus.2009.10.006>, 2010.
- Lindzen, R. S.: Instability of a plane-parallel shear flow (Toward a mechanism for how it works), *PAGEOPH*, 126, 103–121, 1988.
- Liu, J. and Schneider, T.: Mechanisms of Jet Formation on the Giant Planets, *J. Atmos. Sci.*, 67, 3652–3672, <https://doi.org/10.1175/2010JAS3492.1>, 2010.
- 965 Lorenz, E. N.: Available Potential Energy and the Maintenance of the General Circulation, *Tellus*, 7, 157–167, 1955.
- Lorenz, E. N.: *The Nature and Theory of the General Circulation of the Atmosphere*, WMO, No. 218, T. P. 115, Geneva, Switzerland, 1967.
- Lucarini, V.: Thermodynamic efficiency and entropy production in the climate system, *Phys. Rev. E*, 80, 021 118, <https://doi.org/10.1103/PhysRevE.80.021118>, 2009.

- Mankovitch, C., Marley, M. S., Fortney, J. J., and Movshovitz, N.: Cassini ring seismology as a probe of Saturn's interior. I. rigid rotation, *Astrophys. J.*, 871, 1, <https://doi.org/10.3847/1538-4357/aaf798>, 2019.
- Margules, M.: Über die Energie der Stürme, *Jahrb. Kais.-kön Zent. für Met. und Geodynamik*, Vienna, pp. 1–26, 1903.
- Mendonça, J. M. and Read, P. L.: Exploring the Venus global super-rotation using a comprehensive general circulation model, *Plan. Space Sci.*, 134, 1–18, 2016.
- Merlis, T. M. and Schneider, T.: Atmospheric dynamics of Earth-like tidally locked aquaplanets, *J. Adv. Model. Earth Syst.*, 2, 13, 2010.
- 975 Mitchell, J. L. and Vallis, G. K.: The transition to superrotation in terrestrial atmospheres, *J. Geophys. Res.*, 115, E12 008, 2010.
- Montabone, L., Marsh, K., Lewis, S. R., Read, P. L., Smith, M. D., Holmes, J., Spiga, A., Lowe, D., and Pamment, A.: The Mars Analysis Correction Data Assimilation (MACDA) Dataset V1.0, *Geosci. Data J.*, 1, 129–139, <https://doi.org/10.1002/gdj3.13>, 2014.
- Mu, M. and Wu, Y.-H.: Arnol'd nonlinear stability theorems and their application to the atmosphere and oceans, *Surveys in Geophys.*, 22, 383–426, 2001.
- 980 Niino, H. and Misawa, N.: An experimental and theoretical study of barotropic instability, *J. Atmos. Sci.*, 41, 1992–2011, 1984.
- Palotai, C., Dowling, T. E., and Fletcher, L. N.: 3D Modeling of interactions between Jupiter's ammonia clouds and large anticyclones, *Icarus*, 232, 141–156, <https://doi.org/10.1016/j.icarus.2014.01.005>, 2014.
- Pedlosky, J.: Finite-amplitude baroclinic waves, *J. Atmos. Sci.*, 27, 15–30, 1970.
- Pedlosky, J.: Finite-amplitude baroclinic waves with small dissipation, *J. Atmos. Sci.*, 28, 587–597, 1971.
- 985 Peixoto, J. and Oort, A.: The Annual Distribution of Atmospheric Energy on a Planetary Scale, *J. Geophys. Res.*, 20, 2149–2159, <https://doi.org/10.1029/JC079i015p02149>, 1974.
- Read, P. L.: A combined laboratory and numerical study of heat transport by baroclinic eddies and axisymmetric flows, *J. Fluid Mech.*, 489, 301–323, 2003.
- Read, P. L.: Dynamics and circulation regimes of terrestrial planets, *Plan. Space Sci.*, 59, 900–914, 2011.
- 990 Read, P. L. and Lebonnois, S.: Superrotation on Venus, on Titan, and Elsewhere, *Ann. Rev. Earth Planet. Sci.*, 46, 175–202, <https://doi.org/10.1146/annurev-earth-082517-010137>, 2018.
- Read, P. L., Rhines, P. B., and White, A. A.: Geostrophic scatter diagrams and potential vorticity dynamics, *J. Atmos. Sci.*, 43, 3226–3240, 1986.
- Read, P. L., Gierasch, P. J., Conrath, B. J., Simon-Miller, A., Fouchet, T., and Yamazaki, Y. H.: Mapping potential-vorticity dynamics on Jupiter. I: Zonal-mean circulation from Cassini and Voyager 1 data, *Quart. J. R. Meteor. Soc.*, 132, 1577–1603, <https://doi.org/10.1256/qj.05.34>, 2006.
- 995 Read, P. L., Conrath, B., Fletcher, L., Gierasch, P., Simon-Miller, A., and Zuchowski, L.: Mapping potential-vorticity dynamics on Saturn: Zonal-mean circulation from Cassini and Voyager data, *Plan. Space Sci.*, 57, 1682–1698, <https://doi.org/10.1016/j.pss.2009.03.004>, 2009a.
- Read, P. L., Dowling, T. E., and Schubert, G.: Saturn's rotation period from its planetary wave configuration, *Nature*, 460, 608–610, <https://doi.org/10.1038/nature08194>, 2009b.
- 1000 Read, P. L., Pérez, E. P., Moroz, I. M., and Young, R. M. B.: General Circulation of Planetary Atmospheres: Insights from Rotating Annulus and Related Experiments, in: *Modelling Atmospheric and Oceanic Flows: Insights from Laboratory Experiments and Numerical Simulations*, edited by von Larcher, T. and Williams, P. D., pp. 9–44, American Geophysical Union and Wiley, Hoboken, NJ, USA, 2015.
- Read, P. L., Tabataba-Vakili, F., Wang, Y., Augier, P., Lindborg, E., Valeanu, A., and Young, R. M. B.: Comparative terrestrial atmospheric circulation regimes in simplified global circulation models. Part II: Energy budgets and spectral transfers, *Quart. J. R. Meteor. Soc.*, 144, 2558–2576, <https://doi.org/10.1002/qj.3351>, 2018.
- 1005

- Ripa, P.: General stability conditions for zonal flows in a one-layer model on the β -plane or the sphere, *J. Fluid Mech.*, 126, 463–489, 1983.
- Sakai, S.: Rossby–Kelvin instability: a new type of ageostrophic instability caused by a resonance between Rossby waves and gravity waves, *J. Fluid Mech.*, 202, 149–176, 1989.
- 1010 Salyk, C., Ingersoll, A. P., Lorre, J., Vasavada, A., and Del Genio, A. D.: Interaction between eddies and mean flow in Jupiter’s atmosphere: analysis of Cassini imaging data, *Icarus*, 185, 430–442, <https://doi.org/10.1016/j.icarus.2006.08.007>, 2006.
- Sánchez-Lavega, A., Lebonnois, S., Imamura, T., Read, P., and Luz, D.: The Atmospheric Dynamics of Venus, *Space Sci. Rev.*, 212, 1541–1616, <https://doi.org/10.1007/s11214-017-0389-x>, 2017.
- Schneider, T.: The thermal stratification of the extratropical troposphere, in: *The Global Circulation of the Atmosphere*, edited by Schneider, T. and Sobel, A. H., pp. 47–77, Princeton University Press, Princeton, USA, 2007.
- 1015 Schneider, T. and Walker, C. C.: Self-organization of atmospheric macroturbulence into critical states of weak nonlinear eddy-eddy interactions, *J. Atmos. Sci.*, 63, 1569–1586, 2006.
- Scott, R. K. and Dunkerton, T. J.: Vertical structure of tropospheric winds on gas giants, *Geophys. Res. Lett.*, 44, 3073–3081, <https://doi.org/10.1002/2017GL072628>, 2017.
- 1020 Seviour, W. J. M., Waugh, D. W., and Scott, R. K.: The stability of Mars’s annular polar vortex, *J. Atmos. Sci.*, 74, 1533–1547, <https://doi.org/10.1175/JAS-D-16-0293.1>, 2017.
- Sommeria, J., Meyers, S. D., and Swinney, H. L.: Experiments on vortices and Rossby waves in eastward and westward jets, in: *Nonlinear Topics in Ocean Physics*, edited by Osborne, A. R., pp. 227–269, North Holland, Amsterdam, 1991.
- Spiga, A., Guerlet, S., Millour, E., Indurain, M., Meurdesoif, Y., Cabanes, S., Dubos, T., Leconte, J., Boissinot, A., Lebonnois, S., Sylvestre, M., and Fouchet, T.: Global climate modeling of Saturn’s atmosphere. Part II: Multi-annual high-resolution dynamical simulations, *Icarus*, 335, 113–137, 2020.
- 1025 Stamp, A. P. and Dowling, T.: Jupiter’s winds and Arnol’d’s second stability theorem: slowly moving waves and neutral stability, *J. Geophys. Res.*, 98, 18 847–18 855, 1993.
- Stone, P.: Baroclinic Adjustment., *J. Atmos. Sci.*, 35, 561–571, [https://doi.org/10.1175/1520-0469\(1978\)035<0561:BA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035<0561:BA>2.0.CO;2), 1978.
- 1030 Sugimoto, N., Takagi, M., and Matsuda, Y.: Baroclinic instability in the Venus atmosphere simulated by GCM, *J. Geophys. Res.*, 119, 1950–1968, <https://doi.org/10.1002/2014JE004624>, 2014.
- Tabataba-Vakili, F., Read, P. L., Lewis, S. R., Montabone, L., Ruan, T., Wang, Y., Vaeleanu, A., and Young, R. M. B.: A Lorenz/Boer energy budget for the atmosphere of Mars from a “reanalysis” of spacecraft observations, *Geophys. Res. Lett.*, 42, 8320–8327, <https://doi.org/10.1002/2015GL065659>, 2015.
- 1035 Vallis, G. K.: Numerical studies of eddy transport properties in eddy-resolving and parametrized models, *Quart. J. R. Meteor. Soc.*, 114, 183–204, 1988.
- Vallis, G. K.: *Atmospheric and Oceanic Fluid Dynamics - Fundamentals and Large-Scale Circulation* (2nd Edition), Cambridge University Press, Cambridge, UK, 2017.
- Wang, Y., Read, P. L., Tabataba-Vakili, F., and Young, R. M. B.: Comparative terrestrial atmospheric circulation regimes in simplified global circulation models. Part I: From cyclostrophic super-rotation to geostrophic turbulence, *Quart. J. R. Meteor. Soc.*, 144, 2537–2557, <https://doi.org/10.1002/qj.3350>, 2018.
- 1040 Williams, G. and Holloway, J.: The range and unity of planetary circulations, *Nature*, 297, 295–299, 1982.
- Young, R. E., Houben, H., and Pfister, L.: Baroclinic instability in the Venus atmosphere, *J. Atmos. Sci.*, 41, 2310–2333, 1984.

- Young, R. M. B.: The Lorenz energy cycle in simulated rotating annulus flows, *Phys. Fluids*, 26, 056 602, <https://doi.org/10.1063/1.4873921>, 1045 2014.
- Young, R. M. B. and Read, P. L.: Forward and inverse kinetic energy cascades in Jupiter's turbulent weather layer, *Nature Phys.*, 13, 1135–1140, <https://doi.org/10.1038/NPHYS4227>, 2017.
- Young, R. M. B., Read, P. L., and Wang, Y.: Simulating Jupiter's weather layer. Part I: Jet spin-up in a dry atmosphere, *Icarus*, 326, 225–252, 2019a.
- 1050 Young, R. M. B., Read, P. L., and Wang, Y.: Simulating Jupiter's weather layer. Part II: Passive ammonia and water cycles, *Icarus*, 326, 253–268, 2019b.
- Zurita-Gotor, P.: The sensitivity of the isentropic slope in a primitive equation dry model, *J. Atmos. Sci.*, 65, 43–65, 2008.
- Zurita-Gotor, P. and Lindzen, R. S.: Theories of baroclinic adjustment and eddy equilibrations, in: *The Global Circulation of the Atmosphere*, edited by Schneider, T. and Sobel, A. H., pp. 22–46, Princeton University Press, Princeton, USA, 2007.