

# The Impact of Entrained Air on Ocean Waves

Juan M. Restrepo<sup>1</sup>, Alex Ayet<sup>2,3</sup>, and Luigi Cavaleri<sup>4</sup>

<sup>1</sup>Computer Science and Mathematics Division, Oak Ridge National Laboratory, Oak Ridge TN 37831 USA

<sup>2</sup>Ifremer, CNRS, IRD, Univ. Brest/ Laboratoire d'Océanographie Physique et Spatiale (LOPS), IUEM, Brest, France

<sup>3</sup>LMD/IPSL, CNRS, École Normale Supérieure, PSL Research University, Paris, France

<sup>4</sup>ISMAR, Istituto di Scienze Marine, Arsenale - Tesa 104, Castello 2737/F, 30122 Venice Italy

**Correspondence:** Juan M. Restrepo (restrepojm@ornl.gov)

**Abstract.** We make a physical-mathematical analysis of the implications that the presence of a large number of tiny bubbles may have, when present, on the thin upper layer of the sea. In our oceanographic example the bubbles are due to an intense rain It was found that the bubbles increase momentum dissipation in the near surface, as well as affecting the surface tension force. For short waves, the implications of increased vorticity are: momentum exchanges between wave and mean flow and  
5 modifications to the wave dispersion relation. For the direct effect we have analyzed, the implications are estimated non-significant when compared to other processes of the ocean. However, we hint to the possibility that our analysis may be useful in other areas of research or practical application.

*Copyright statement.*

## 1 Introduction

10 Our interest is on the consequences of an intense rain on the small scale roughness of the ocean surface. More specifically, we analyze how the presence of tiny, rain induced air bubbles in the thin upper layer of the ocean affects the high frequency tail of the gravity wave spectrum, including capillary waves. Although tiny, both in length and height, with respect to the majestic motion of the surface, these waves are important for both physical and practical reasons. Physically, they control large part of the exchanges between ocean and atmosphere: they act as roughness elements for atmospheric turbulence and modulate the  
15 interaction between longer waves, currents, and wind (e.g. Ayet et al., 2020). At the same time, their presence is what allows the scatterometers to measure globe-wide the winds on the ocean (Kudryavtsev et al., 1999).

The disappearance of capillary sea waves with rain is not new. Known to mariners since their early attempts (a reduced small scale roughness leads also to reduced breaking), the subject has also been studied in recent times by, e.g.

, see Le Méhauté and Khangaonkar (1990) and Cavaleri et al. (2018). The former ones focused on the mechanical action  
20 of rain on the dynamics of the wave orbital motion. The latter ones attention, taking the wavelets attenuation as a de facto evidence, focused more on the implications for air-sea interactions. In this work we try to clarify one of the reasons why rain attenuates surface wavelets. Granted, if strong enough, the rain drops mechanical action and induced turbulence, we focus our

attention on the rain induced presence of a large number of tiny air bubbles just below the surface (a few centimeters) and on how they affect the wave motion. To avoid the dominance of other processes, we exclude stormy conditions, considering either only swell or calm sea cases, but with enough wind to lead to waves on the surface. As for rain, just to frame the order of magnitude, under an intense rain with, e.g.,  $100 \text{ mm h}^{-1}$ , assuming 4 mm diameter rain drops (volume  $3 \times 10^{-6} \text{ m}^3$ ), this translates into 100 drops per  $\text{m}^2$  per second. Wolf (2001) suggests that under these conditions air injected into the sea can be a few  $\text{cm}^3 \text{ m}^{-2} \text{ s}^{-1}$ . Compared to other mechanisms that introduce air into the sea, one can conclude that rain contribution is very small. True in general as quantitative terms, we consider the process interesting from the physical point of view and with potential application in other practical fields.

On this basis we make a physical-mathematical analysis of the ensuing processes. In Section 2, we propose a model that suggests how the entrainment of air leads to an increase of the effective viscosity of the upper ocean. We then use results from ocean acoustics (see Oguz and Prosperetti (1990) and Prosperetti and Oguz (1993)) to compute the distribution of air bubbles for a given rain rate (Section 3). How changes of effective viscosity affect gravity waves is taken up in Section 4. It is found that under vigorous rains the air concentration due to air bubbles near the surface is capable of producing a damping effect akin to (idealized) contamination of a free surface. Discussion of the results and conclusions, and possible extensions of the present work appear in Sections 5 and 6.

## 2 The Dynamical Approach And The Presence Of Bubbles

To determine the impact of air bubbles on the thin upper level of the ocean, we perform homogenization on the Navier-Stokes equations. Homogeneization is a well-established technique in modeling transport in complex media endowed with statistical homogeneity in the media, as viewed at large scales (see Babuska (1976); Bensousan et al. (1978); Cioranescu and Donato (1999)).

We assume that air bubbles are distributed uniformly in the transverse direction, within the vicinity of the ocean surface. For our specific example we explore how the presence of bubbles affects the effective viscosity of the fluid averaged over a cell  $\Omega$ , of size  $\ell^3$  (sub-wave scale), over which the distributions of the density and viscosity of the combined water and air bubble mixture are statistically stationary. The typical sub-wave speed is  $u_\Omega$ . The ratio of the bubble radii to the averaging  $\ell$ , defines for us a small parameter  $\epsilon \ll 1$ .

Velocity and position are denoted by  $\mathbf{u} = (u, v, w)$ ,  $\mathbf{r} = (x, y, z)$ , respectively. The free surface is denoted by  $\eta$  and  $z = 0$  corresponds to the quiescent sea level. Gravity is  $g\hat{\mathbf{z}}$  and the vector  $\hat{\mathbf{z}}$  points upwards, along increasing  $z$ . Velocity is scaled by  $u_\Omega$ , length by  $\ell$ , time by  $\ell/u_\Omega$ , density by  $\rho_w$ , the density of water, pressure by  $\rho_w u_\Omega^2$ . We then define a Reynolds number  $\alpha = u_\Omega \ell / \nu_w$ , where  $\nu_w$  is the kinematic viscosity of water. We also define a Froude number  $1/\sqrt{\gamma} = u_\Omega / \sqrt{g\ell}$ . The scaling leads to

$$\alpha \mathbf{u}_t + \alpha \mathbf{u} \cdot \nabla \mathbf{u} = -\alpha \gamma \nabla \Pi + \nabla \cdot [D(\mathbf{r})\Xi], \quad (1)$$

$$\nabla \cdot \mathbf{u} = 0, \quad (2)$$

55 where  $\Pi$  is pressure and  $\Xi$  the stress tensor  $\Xi = \nabla \mathbf{u} + [\nabla \mathbf{u}]^\top$ .  $D$  is the proportionality tensor.

Let  $\mathbf{R} = (X, Y, Z)$  be the large-scale space variable, such that  $\nabla = \nabla + \epsilon \nabla_{\mathbf{R}}$ , and assume slow time  $\partial_t = \delta \partial_T$ . Also, assume that  $\Pi(\mathbf{r}, \mathbf{R}, T) = p - z + \epsilon^2 p_0 + \epsilon^4 p_1 + \dots$ ,  $\mathbf{u} = \epsilon(\mathbf{u}_0 + \epsilon \mathbf{u}_1 + \epsilon^2 \mathbf{u}_2 \dots)$  and  $\eta = \alpha(\eta_0 + \epsilon \eta_1 + \epsilon^2 \eta_2 \dots)$ . The orders are  $\alpha = \mathcal{O}(\epsilon^2)$ ,  $\delta = \mathcal{O}(\epsilon)$ , and  $\gamma = \mathcal{O}(\epsilon^{-1})$ .

60 In the following we derive the fluid mechanics equations for averaged dynamic quantities (see related work, Caflich et al. (1985)), appropriate at wave spatio-temporal scales, for example. Our goal is to find effective equations valid in a composite media, under the assumptions made above. The equations and the dynamic quantities will agree with the non-averaged ones, when the media has a single species density  $\rho_w$  and the tensor  $D = \nu_w \delta_{ij}$ . Here we introduce the inhomogeneity as tiny air bubbles, uniformly distributed in the transverse direction in a matrix of water. The source of the bubbles is rain and the connection between the bubble density in the near surface and the rain rate will be suggested in a later section. Anticipating some of  
 65 the results, the momentum and continuity equations for the averaged dynamic quantities are similar to the point versions. However, material properties, such as the density and the tensor  $D$  are different for the averaged cells, affecting quantitatively the momentum predictions.

Collecting by orders in  $\epsilon$ , the momentum equation reads,

–  $\mathcal{O}(\epsilon)$ :

70 
$$\nabla \cdot [D(\mathbf{r})\Xi_0] = 0.$$

At this microscale, pressure gradients in the cell are negligible, as are variations in velocity within the cell. We are also assuming stationary conditions. Upon integrating in  $\mathbf{r}$  over and invoking periodicity it is clear that  $\Xi_0 = \Xi_0(\mathbf{R}, t, T)$  and thus  $\mathbf{u}_0 = \mathbf{u}_0(\mathbf{R}, t, T)$ .

–  $\mathcal{O}(\epsilon^2)$ :

75 
$$\nabla \cdot [D(\mathbf{r})\Xi_1] + \nabla \cdot [D(\mathbf{r})\Xi_0] + \nabla_{\mathbf{R}} \cdot [D(\mathbf{r})\Xi_0] = 0. \quad (3)$$

The last term above is zero (we made use of  $\nabla \cdot (\nabla \Xi)^T = \nabla \cdot (\nabla \cdot \Xi)$ ). Integration by parts of (3) and using periodicity,  $\Xi_1(\mathbf{r}, \mathbf{R}, t, T) = -\Xi_0 + D^{-1}(\mathbf{r})C(\mathbf{R}, t, T)$ , where  $C$  is a tensor that is independent of  $\mathbf{r}$ . Periodicity and integration in  $\mathbf{r}$  over  $\Omega$  of every differential term in  $\Xi_1$  implies that

80 
$$0 = -\Omega \Xi_0 + \int_{\Omega} D^{-1}(\mathbf{r}) d\mathbf{r} C(\mathbf{R}, t, T). \quad (4)$$

Solving for the tensor from this equation and integrating with respect to  $\mathbf{r}$  one obtains

$$C(\mathbf{R}, t, T) = \left[ \frac{1}{\Omega} \int_{\Omega} D^{-1}(\mathbf{r}) d\mathbf{r} \right]^{-1} \Xi_0.$$

Define the tensor

$$\chi = \left[ \frac{1}{\Omega} \int_{\Omega} D^{-1}(\mathbf{r}) d\mathbf{r} \right]^{-1}. \quad (5)$$

Returning to (3),

$$85 \quad \Xi_1 = -\Xi_0 + D^{-1}(\mathbf{r})\chi\Xi_0, \quad (6)$$

which prescribes the first order correction to the stress tensor, which could be obtained explicitly if the zeroth order is known ( $D^{-1}(\mathbf{r})$  is an observed quantity). Taking the divergence of (6)

$$\nabla_{\mathbf{R}} \cdot (D\Xi_1) = -\nabla_{\mathbf{R}} \cdot (D(\mathbf{r})\Xi_0) + \nabla_{\mathbf{R}} \cdot (\chi\Xi_0). \quad (7)$$

–  $\mathcal{O}(\epsilon^3)$ : the momentum balance reads

$$90 \quad \frac{\partial \mathbf{u}_0}{\partial T} + \mathbf{u}_0 \cdot \nabla_{\mathbf{R}} \mathbf{u}_0 + \mathbf{u}_0 \cdot \nabla \mathbf{u}_1 + \nabla_{\mathbf{R}} p_0 = \\ + \nabla \cdot [D(\mathbf{r})\Xi_1] + \nabla_{\mathbf{R}} \cdot [D(\mathbf{r})\Xi_0] \\ + \nabla \cdot [D(\mathbf{r})\Xi_2] + \nabla_{\mathbf{R}} \cdot [D(\mathbf{r})\Xi_1]. \quad (8)$$

Using (7) in the last term in (8) and averaging all quantities over  $\Omega$  (i.e., integrating over  $\mathbf{r}$  using periodicity in the cells), leads to

$$95 \quad \frac{\partial \mathbf{u}_0}{\partial T} + \mathbf{u}_0 \cdot \nabla_{\mathbf{R}} \mathbf{u}_0 + \nabla_{\mathbf{R}} p_0 = \nabla_{\mathbf{R}} \cdot \{ \chi (\nabla_{\mathbf{R}} \mathbf{u}_0 + [\nabla_{\mathbf{R}} \mathbf{u}_0]^\top) \}. \quad (9)$$

The homogenized incompressibility condition is


$$\nabla_{\mathbf{R}} \cdot \mathbf{u}_0 = 0. \quad (10)$$

At this point we have a set of (closed) mean field equations for momentum and mass conservation, valid at the wave scales  $(\mathbf{R}, T)$ . The mean field velocity and pressure are well defined assuming homogeneity conditions at the small scale are valid.  
100 The effective tensor of proportionality in the stress tensor term  $\chi$  is formally obtained by knowing the microscale composition, density distribution and dynamic viscosities. In practice it is obtained by examining the stress/strain relation for the composite fluid.

Since the above derivation is not explicitly connected to rain, one can imagine that the same approach applies to similar situations where, for whichever reason, a large number of bubbles is distributed in the enclosing medium. We will touch further  
105 this point in the final Discussion.

The tensor  $\chi$  takes the value of the ocean, changing it when bubbles appear for whatever reason. In what follows we limit ourselves to the simplest possible case:  $\chi = \nu_w \delta_{ij}$ , when bubbles are not present, where  $\delta_{ij}$  is the 3-space dimensional Kronecker Delta. For our specific example we use (5) assuming a matrix consisting of a homogeneous concentration of air bubbles, with diffusion constant  $\nu_a$ , in a background fluid. In this very simple case, the enhanced value of  $\chi$  becomes

$$110 \quad \chi = K \delta_{ij}, \text{ with } K = \frac{\nu_w}{1 - \Theta(1 - N_\nu)}, \quad (11)$$



**Figure 1.** Relative effective momentum dissipation  $K/\nu_w$ , as a function of volumetric ratio of air to water  $\Theta$ , using (11) . Assumed here, the ratio  $N_\nu := \nu_w/\nu_a = 1/15$ . The range of  $\Theta$  has been limited to that applicable to oceanic conditions (over the whole range, the relative effective momentum dissipation is a monotonically increasing function).

where  $N_\nu = \frac{\nu_w}{\nu_a}$ , and  $\Theta$  connotes the volumetric ratio of air to water. For a given  $\Theta > 0$ ,  $N_\nu < 1$  leads to an increased effective momentum dissipation, and, when  $N_\nu > 1$ , the effective dissipation is lower. For air or oil bubbles,  $N_\nu < 1$ .

With a fixed ratio  $N_\nu = 1/15$ , which is roughly the ratio of viscosities of water and air, Figure 1 shows how the effective dissipation coefficient  $K$  changes, as a function of the volumetric ratio  $\Theta$ . The relationship is nearly linear for the small volume  
 115 fractions of the oceanic case. If we adopt the specific form of  $\chi$  as in (11), the dissipation term in (9) becomes  $K \nabla_{\mathbf{R}}^2 \mathbf{u}_0$  when bubbles are present and  $\nu_w \nabla_{\mathbf{R}}^2 \mathbf{u}_0$ , when there are no air bubbles. Equation (11) leads us to conclude that the effective momentum dissipation of water with air bubbles will be different from (actually larger than)  $\nu_w$ . Physically, there is an increase in the rotational component of the flow and an associated increase in energy dissipation.

### 3 Air Entrainment due to Rain

120 In this section we assume a statistical distribution for rain drops and derive the consequent density of air bubbles in the ocean surface layer.

#### 3.1 Rain Distribution

The density of rain drops is assumed to follow the Marshall-Palmer distribution (see Marshall (1948)). The density  $N_r$  of rain drops of radius  $r$  [m] per unit volume [number  $\text{m}^{-3} \text{m}^{-1}$ ] is

$$125 \quad N_r(r, R) = 2N_0 \exp(-\Lambda(R)r), \quad (12)$$

where  $N_0 = 8 \times 10^6 \text{ m}^{-3} \text{ m}^{-1}$ ,  $\Lambda(R) = 8.2R^{-0.21} \times 10^3 [\text{m}^{-1}]$ , and  $R [\text{mm h}^{-1}]$  is the rain rate. Using  $N_r$  we can then define the drop rate density (DRD), which describes the rate of falling rain drops of radius  $r$  per surface area [number  $\text{m}^{-2} \text{s}^{-1} \text{m}^{-1}$ ]. Namely,

$$\text{DRD}(r, R) = w_r(r)N_r(r, R), \quad (13)$$

130 where  $w_r$  is the terminal velocity of drops in the air (see Snyder (1990)). The terminal velocity is computed following the third order polynomial estimates of Dingle and Lee (1972).

### 3.2 Bubble Production

To complete our quantitative example we need to relate the number of bubbles in the upper layer of the ocean to rain. Falling rain drops generate subsurface air bubbles in the neighborhood of the sea surface (see Prosperetti and Oguz (1993) for a review). Small rain drops produce small air bubbles with a very narrow distribution of radii. Intermediate rain drops, with a radius between 0.55 - 1.1 mm, were shown not to produce air bubbles (see Medwin et al. (1992)). For these, the impinging rain drops do not have the kinetic energy necessary to produce the requisite conical crater and jetting of the sea surface that engulfs air. Large rain drops, with radii larger than 1.1 mm, create a crater and a canopy on the sea surface, which by collapsing produces a downward liquid jet at the bottom of the crater, followed by the generation of an air bubble. At high rain rates the larger rain drops are responsible for the bulk of the gas injection, creating bubbles with a varied distribution of radii.

The production of air bubbles by rain can be measured by acoustic means (see Prosperetti and Oguz (1993)). Oguz and Prosperetti (1990) classify air bubble production by falling rain drops in two regimes. Small rain drops, of radii between 0.41 and 0.55 mm, create *type I* air bubbles. These have a radius of 0.22 mm and are created at the bottom of a conical crater created on the ocean surface by the impinging rain drop. These air bubbles have a narrow acoustic spectrum with a spectral peak at 14 Hz. The acoustic spectrum for these type I air bubbles was found to be insensitive to the rain rate. Medwin et al. (1990) observed that when rain hits the ocean surface at an angle, owing to strong winds or very steep wave conditions, the acoustic spectrum peak shifts downward and there is a broadening of the spectrum at higher frequencies. The type I air bubbles do not contribute significantly to total submerged gas volume and their contribution will be ignored in what follows.


Rain drops with a radius greater than 1.1 mm produce *type II* air resonant bubbles of varying radius. The relationship between the rain drop radius  $r$  [m] and the peak acoustic emission frequency  $f_0$  [kHz] due to trapped air bubbles was empirically determined (see Medwin et al. (1992)) as

$$f_0 = \frac{160}{8r^3 \times 10^9} + 0.6. \quad (14)$$

The relation between the near surface air bubble radius  $a$  [m] and the peak acoustic emission follows from the Rayleigh-Plesset equation (Leighton, 1994, p. 306). For large bubbles considered here, it can be simplified to Minnaert's formula (see Minnaert (1933); Plesset and Prosperetti (1977)),

$$f_0 = \frac{1}{2\pi a \times 10^3} \sqrt{\frac{3\gamma P}{\rho_0}}, \quad (15)$$

where  $\gamma = 1.4$  is the ratio of specific heats of the bubble gas,  $P$  is the ambient pressure (surrounding the bubble), and  $\rho_0 = 1030 \text{ kg m}^{-3}$  the density of sea water. Close to the surface, the ambient pressure is approximately the surface pressure  $P = 1.01 \times 10^5 \text{ Pa}$  and Equation (15) further simplified as  $f_0 = 3.25 \times 10^{-3}/a$  (see (Medwin and Clay, 1997)).



**Figure 2.** Probability of air bubble creation by large rain drops as a function of rain drop radius  $r$  (from Medwin et al. (1992)).

160 The relation between the entrained air bubble radius  $a(r)$  [m] and the incident rain drop radius  $r$ , for large rain drops, thus reads, for  $r > 1.1 \times 10^{-3}$  m,

$$a(r) = 3.25 \times 10^{-3} \left( \frac{160}{8r^3 \times 10^9} + 0.6 \right)^{-1}. \quad (16)$$

As an example, rain drops of radius 1.1-2.3 mm produce *type II* air bubbles of radius 0.2-1.3 mm, respectively.

Small rain drops ( $r < 0.55$  mm) almost always produce type I air bubbles. However, this is not the case for the larger rain  
 165 drops. The distribution of the number of air bubbles produced, as a function of the rain drop radius  $r$ , was found by Medwin et al. (1992), and is depicted in Figure 2 as a probability distribution.

### 3.3 Bubble Distribution in the Wave Boundary Layer

[Tying back to Section 2](#), by finding the bubble distribution in the wave boundary layer, we can estimate the dimensionless volume fraction  $\Theta$ . More precisely, the volume fraction reads  $\Theta = \int V(a)N(a)da$ , for bubbles of radius  $a$  with density  $N(a)$   
 170 [number  $m^{-3} m^{-1}$ ] that occupy a volume  $V(a) = (4/3)\pi a^3$ . The aim of the final part of this section is to link  $\Theta$  to the rain precipitation rate.

In so doing we will make some approximations. First, bubbles are injected at a very limited depth by rain drops, and this depth may vary with bubble size (see Ho et al. (2000)). Secondly, a homogeneous air bubble distribution in the layer between the depth of injection and the surface is an approximation, as accumulation of bubbles near the surface can be highly complex due to buoyant forces, damping, and the background turbulence (see e.g. Merlivat and Memery (1983)). We will specify how this is taken into account.  
 175

The bubble distribution  $N(a)$  can be described by an advection-diffusion equation (see Woolf and Thorpe (1991) and references therein, and a similar case for oil drops, see Moghimi et al. (2018)). This equation describes a balance between advection of bubbles by the vertical fluid velocity, attenuation of bubble density (i.e. diffusion, which includes the shrinking of bubble  
 180 radius during their lifetime, see Merlivat and Memery (1983)), the production of bubbles by rain and a sink term that captures the loss of air bubbles bursting at the surface or dispersed by oceanic turbulence.

Solutions of the advection-diffusion equation necessitate specification of the velocity and the dispersion, generating non-stationary descriptions that might also include mixing due to wave turbulence (see Restrepo et al. (2015); Moghimi et al. (2018)).

185 In the following, the advection-diffusion equation is simplified to estimate the air bubble concentration in a thin layer close to the surface, similarly to those in Keeling (1993). First, we assume that the air bubble concentration can be linearly related to a bulk bubble distribution (i.e. averaged over the thin layer), which we thus define as  $c_e N(a)$ . Second, a steady state is considered, in which the bulk bubble distribution follows from a balance between the incoming bubble flux due to rain and the outgoing bubble flux due to upward bubble advection and bursting at the surface:

$$190 \quad w_r(r)P_r(r)N_r(r) = c_e W[a(r)]N[a(r)], \quad (17)$$

where  $P_r(r)$  is the probability of production of an air bubble by a rain drop of radius  $r$  (see Figure 2), and  $W(a)$  is the upward vertical ascent speed of bubbles of radius  $a$ . From this equation, the bubble distribution reads  $N(a) = w_r N_r(r, R) P_r / [c_e W(a)]$ .

We assume as an estimate for the bubble upward speed  $W(a)$  the form

$$W(a) = \sqrt{\frac{\tau}{a\rho_0} + ga} \quad (18)$$

195 where  $\tau = 72.8 \times 10^{-3} \text{ N m}^{-1}$  is the air-sea surface tension (see Clift et al. (1978), p. 172). This form, predicted in Mendelson (1967), is valid for ellipsoidal bubbles with radius greater than 0.65 mm, whose upward path is not rectilinear.

Referring to the above approximations, the factor  $c_e$  can first be interpreted as a depth scaling parameter accounting for differences between the bulk estimate and the non-homogeneous bubble distribution  $N$ . A further approximation is that the effective vertical velocity of bubbles can be significantly lower than (18), up to 60% due to the presence of contaminants at the bubble surface (see Clift et al. (1978), Fig. 7.3). Hence the factor  $c_e$  is also to be interpreted as a decrease in the bubble vertical velocity, which then reads  $c_e W(a)$ .

Making use of (13),  $N(a) = (\text{DRD} \times P_r)(r) / c_e W(a)$ . The volumetric ratio  $\Theta = \int V(a)N(a)da$ , for bubbles of radius  $a$  with density  $N(a)$  that occupy volume  $V(a) = (4/3)\pi a^3$  then reads

$$\Theta(R) = \int_{1.1 \times 10^{-3}}^{2.3 \times 10^{-3}} \frac{4\pi}{3} a(r)^3 \frac{(\text{DRD} \times P_r)(r)}{c_e W(a)} dr, \quad (19)$$

205 where the limits of integration encompass the volumetric contributions of type II bubbles.

The obtained values of volume fraction are consistent with data from laboratory experiments of Ho et al. (2000) (see their Figure 7), which suggests that for rain rates of  $114 \text{ mm h}^{-1}$  the volume fraction near the air/water interface is between  $10^{-6}$  and  $10^{-5}$ , corresponding to  $c_e$  of the order of 0.1.

#### 4 The Effect of Rain on Gravity and Capillary Waves



210 The subject of mathematical models for wave dissipation that reproduce the observed dissipation rates observed in laboratory experiments is considered in Henderson et al. (2015) study. The focus is on the dissipation effects of small waves due to surface contamination, with air above it. Several wave damping models are compared to data. In this work a model due Jenkins and Jacobs (1997) is compared to experimental data of damping of small waves. The model posits a small (boundary) layer sitting over a deep water layer. The two layer model yields expressions valid under several limits. Our work can contribute  
 215 to this work by making definite how the equations in the upper layer are modified by the presence of air bubbles and how their density relates to rain, if this is the mechanism for their generation.

In what follows we will invoke a far less general, but consistent, way to address the question on how the presence of rain bubbles affect water waves. The derivation of infinitesimal amplitude gravity waves dynamics, starting from Navier Stokes, appears in Lamb (1916), Article 349. We revert back to dimensional quantities in what follows. The solution  $\mathbf{u}_0$  satisfies the  
 220 linear version of (9) and (10) and the two stress conditions at the surface. Namely,

$$K[\partial_X v_0 + \partial_Y u_0] = 0, \quad \text{at } z = \eta, \quad (20)$$

$$-\partial_T \phi_0 + \left(g - \frac{\tau}{\rho_*} \partial_{XX}\right) \eta_0 + 2K \partial_Y v_0 = 0, \quad \text{at } z = \eta, \quad (21)$$

where  $\phi_0$  is the mean field velocity potential and  $\rho_*$  is the density of water (no rain) or the reduced density (raining, with air bubbles in the water), and time (at wave scales) is  $T$ . At some distance  $\ell$  below the surface the conditions are

$$225 \quad \underline{\nu_w} [\partial_X v + \partial_Y u] = K [\partial_X v_0 + \partial_Y u_0], \quad \text{at } z = \eta - \ell, \quad (22)$$

$$\underline{-\partial_T \phi + 2\nu_w \partial_Y v} = -\partial_T \phi_0 + 2K \partial_Y v_0, \quad \text{at } z = \eta - \ell, \quad (23)$$

By combining the boundary and non-boundary solutions, the approximate solution of the system, assuming a vanishingly small upper layer thickness and vanishing velocity at depth, is

$$\begin{aligned} \phi &= A e^{kZ + i(kX - \sigma T)} \exp[-2Kk^2 T], \\ 230 \quad \psi &= \frac{2Kk^2 A}{\sigma} e^{kZ + i(kX - \sigma T)} \exp[-2Kk^2 T], \\ \eta &= \frac{kA}{\sigma} e^{i(kX - \sigma T)} \exp[-2Kk^2 T], \end{aligned} \quad (24)$$

where  $\psi$  is the streamfunction and  $\sigma$  the angular frequency. The solution (24) represents infinitesimal progressive waves traveling in the  $X$  direction.

The dispersion relation for the waves is

$$235 \quad \sigma^2 = gk + \frac{\tau}{\rho_*} k^3,$$

showing that changes in the surface tension due to the presence of air bubbles injected by the rain impacts the dispersion relation. The effective density is

$$\rho^* = \frac{1}{\Omega} \int_{\Omega} \rho(\mathbf{r}) d\mathbf{r},$$

which in our simplified accounting would be  $\rho^* = \Theta\rho_a + (1 - \Theta)\rho_w$ .

240 The effective wave dissipation  $2K$  appears in the exponential factor  $\exp[-2Kk^2T]$ . When the rainstorm is sufficiently intense (but the rain drops not too large), the wave dissipation  $2K$  will change from  $2\nu_w$  to a higher value, depending on how much air is injected by the impinging rain drops.


245 The dependence of the effective wave dissipation on the rain rate, using (11) and (19), is depicted in Figure 3. The surface tension effect increases as well affecting the dispersion relation. These effects are manifest primarily in capillary, high frequency wave components. The bubble model derived in Section 3 has one free parameter,  $c_e$ , which controls the upward flux of bubbles. The solid line shows the wave dissipation for  $c_e = 1$ , i.e. the situation for which the bulk estimation of the upward flux of bubbles is assumed to be correct. The dashed line shows the wave dissipation corresponding to the more realistic value  $c_e = 0.1$ , which matches the measured volume fraction of Ho et al. (2000). With respect to  $c_e = 1$ , it corresponds to a situation where the residence time of bubbles in the upper ocean layer is increased, due to e.g. near surface turbulence.

250 In both cases the increase in effective wave damping due to the injection of air, at least as estimated by simple considerations, is small compared to the values reported in the experiments in Peirson et al. (2013) (and in Tsimplis (1992)) a further proof of what stated in the introduction, i.e. that, for large rain rates and, as in the wave tank experiments, large rain drops, the mechanical effect is dominant for the immediate, albeit limited, attenuation of water waves. In these, using very high rain rates, the effective dissipation was found to increase by 3-10 times when rain is present, as compared to without rain.

255 Figure 3 also shows the sensitivity of the model to changes in  $\nu_w/\nu_a$ , which is much lower than to changes in the bubbles volume fraction, i.e. the tuning parameter  $c_e$ . The grey shadings are the intervals obtained by varying the air viscosity inside air bubbles ( $\nu_a$ ) from  $10^{-5}$  to  $10^{-4}$  (the lower and upper part of the shadings respectively), for a fixed  $c_e$  of 0.1. This range of variation of  $\nu_a$  corresponds to variations of the effective kinematic viscosity of air due to the mechanical effect of rain drops, as observed in Harrison and Veron (2017).

## 260 5 Discussion and conclusions

Stimulated by an oceanographic problem, we have developed a physical-mathematical approach to analyze how the presence of a large number of small bubbles affects the characteristics of the containing liquid. Granted the general method and having specified the necessary conditions for its application, we focused on a specific example: the presence of air bubbles in the thin upper layer of the ocean and their impact on ocean surface waves. We first derived the (upscaled) equations for mass and momentum conservation. The upscaled velocity and pressure are averages over subwave scales over which the material with heterogeneities assumes a stationary distribution. These equations revealed that the effective (upscaled) momentum dissipation is increased when bubbles are present with respect to the no-bubble case.



**Figure 3.** Relative effective wave dissipation  $2K/2\nu_w$ , as a function of rain rate  $R$  for two values of the unconstrained parameter  $c_e$ : 1 (solid) and 0.1 (dashed). The very small region with grey shading, below the upper curve, correspond to a variation of air diffusivity between  $10^{-5} \text{ m s}^{-2}$  (dashed line) and  $10^{-4} \text{ m s}^{-2}$ , for a fixed  $c_e = 0.1$ . The figure highlights that the model outcomes are more sensitive to the parameter  $c_e$  than to the value of the diffusivity  $\nu_a$ .

Under homogeneization conditions, the model yields a dynamic equation that has a higher effective diffusion constant with bubbles than without bubbles. This translates into an increased kinematic viscosity of the air-water mixture, which is directly  
 270 related to an increase in the damping of surface waves. The source for the tiny uniformly distributed bubbles is, in our case, rain,  
whose small drops generate the bubble distribution (large drops lead to splashing and turbulence in the upper layer, a different  
 process). For typical rain rates, we found that the resulting wave damping due to bubbles is not significant with respect to other  
damping sources, such as turbulence-induced dissipation due to the impact of large rain drops. We also found that the presence  
of bubbles affects the capillary waves dispersion relation, through a change in the effective surface tension felt by the waves.  
 275 The increase in wave damping (related to kinematic viscosity) when bubbles are present may appear counterintuitive at first  
sight, but results from kinematic viscosity being the ratio of the dynamic viscosity and the density of the fluid of interest. As  
bubbles are injected into water, the homogenized kinematic viscosity decreases, but so does the homogenized fluid density,  
with the overall effect being an increase in the homogenized kinematic viscosity. Physically, the presence of small bubbles  
at the typically small void ratios of non-breaking waves enhances momentum transfer between the waves and the mean flow.  
 280 The enhanced wave rotational component results in higher viscous effects. Moreover, since the density enters explicitly in the  
definition of the surface tension force, when the volume fraction of air to water is not zero the average density of the medium  
decreases, and thus with this decrease comes an increase in surface tension forces. Finally, the viscous effects and changes in  
the surface tension also lead to changes in the dispersion relation, and as a consequence, in the wave group velocity as well.

Many simplifications were made in the formulation of the rain/wave model. In the momentum equation we used the most  
 285 simplistic model for the tensor  $D(\mathbf{r})$ . The consequence are very meak momentum exchanges between rotational and irrotational  
components of the velocity, and this argues for a more vigorous research effort in determining more realistic models for the  
tensor. In particular, the tensor should account for all sources of small scale heterogeneity besides air bubbles. The model  
for the mechanism that ties the rain, the bubble presence in the layer and the momentum and mass conservation equations

below the free surface has three critical parameters: the rain rate  $R$ , the air-to-water volume fraction  $\Theta$  (through the flux  
290 strength parameter  $C_{e\tau}$ , associated with the upward flux of air bubbles from their injection depth to the surface) and the ratio of  
diffusivities of water and air,  $N_D$ . Given those approximations, the basic result is that the volume fraction of air to water would  
have to be exceptionally large for the effect to be significant, when compared to other damping mechanisms. However, granted  
the limited effect in the analyzed case, larger amounts of bubbles or, mutatis mutandis, of buoyant organic or hydrocarbons, are  
certainly possible. In these cases, that we have not explored, the damping effect can indeed be relevant and, within the specified  
295 approximations, quantifiable following the procedure we have outlined.

One last simplification is related to the fact that waves occur in a salty ocean. The density of the fresh water due to rain  
and that of the ocean are different. Both fresh water as well as air bubbles are thus affecting the composition of the ocean  
in the near surface and thus the local density and the tensor  $D(\mathbf{r})$ . The change in the water composition will also affect  
some aspects of the bubble distribution model presented earlier. The more complex composition will affect quantitatively the  
300 results, but it is presumed not to affect the qualitative conclusions presented. Another effect that has been ignored here is  
full considerations of the water/air interface stress conditions when rain is present. In Veron and Meusiciens (2016) a model is  
presented for how the impact of rain affects the boundary conditions (cf., their Eqs. 2.11, 2.14, and for waves 3.4). Namely the  
pressure at the free surface is composed of the atmospheric pressure plus a rain-induced pressure. The rain induced pressure  
is trivially space-averaged in order to enter the homogenized set of momentum/condition equations presented here. The  
305 consequences on the waves enter through Bernoulli's equation and affect the dispersion of the waves. These issues are presented  
in Veron and Meusiciens (2016) and not repeated here.

There are several processes where the mixture of a liquid containing a large number of bubbles or droplets has different char-  
acteristics. Examples are a frothy ocean, whatever the content, cavitating flows as it happens in ship propellers or in pressurized  
flows. One example we came across while preparing this article describes how foams are used to ameliorate unwanted ship  
310 motion due to sloshing of their holding tank contents (see Denkov et al. (2005); Kim et al. (2007)), as well as the stabilizing  
effect of free surface sloshing of bubbly drinks (see Cappello et al. (2015)). These foamy cases are non-trivial extensions of the  
homogenisation procedure we present in this paper. Consideration of chemistry, compressible effects, topology, is required  
and the intricate formulation of stress conditions at the interfaces would need to be derived. Nevertheless the procedure may  
play a constructive role in formulating a mean field description of the dissipation and consequent attenuation of the foam on  
315 sloshing motions.

## 6 Summary

We itemize here below our main conclusions:

1. A general methodology has been developed to quantify the effect of small-scale bubbles on the properties of the  
containing liquid. Averaged dynamic quantities can be defined, and for these, averaged momentum and mass conservation  
320 equations of general applicability can be derived.
2. A model was derived that permits an estimate of the amount of air injected by rain.

3. The presence of tiny air bubbles, that we assume uniformly distributed, changes the physical characteristics of the containing liquid, affecting momentum balances: momentum dissipation of irrotational motions is enhanced due to an increase in the rotational component of the flow.
- 325 4.
5. We have also specifically addressed the impact of the presence of air bubbles on gravity waves. We found that air bubbles increased the effective wave damping and hence, that rain has a damping effect on waves. Changes in the density near the free surface will also affect the surface tension. Since the generation of bubbles by rain is small in the natural setting, the damping effect is small, akin to damping of waves due to surface contamination. An increase in wave damping has implications to momentum exchanges between the waves and the mean flow. Wave damping changes as well as surface tension changes impact the wave dispersion relation as well as the group velocity. These effects are primarily relevant to the dynamics of small and capillary waves.
- 330
- 6.
7. We have cited a number of situations where, granted certain conditions, our approach can be applied for a general estimate of the overall increased dissipation.
- 335

*Acknowledgements.* The code used to generate the figures is freely available on "https://github.com/AAyet/rain\_on\_waves". The authors wish to thank the Kavli Institute of Theoretical Physics (KITP), at the University of California, Santa Barbara, for their hospitality and for supporting this research project. The KITP is supported in part by the National Science Foundation under Grant No. NSF PHY-1748958. AA is supported by DGA grant No. D0456JE075 and the French Brittany Regional Council. AA would like to thank B. Chapron for insightful

340 comments. JMR was supported by grant NSF/OCE #1434198.

## References

- Ayet, A., Chapron, B., Redelsperger, J.-L., Lapeyre, G., and Marié, L.: On the Impact of Long Wind-Waves on Near-Surface Turbulence and Momentum Fluxes, *Boundary-Layer Meteorology*, 174, 465–491, 2020.
- Babuska, I.: Homogenization approach in engineering, vol. 134, *Lecture notes in economics and mathematical systems*, 1976.
- 345 Bensousan, A., Lions, J. L., and Papanicolaou, G.: *Asymptotic Analysis for Periodic Structures*, North-Holland, 1978.
- Caffisch, R. E., Miksis, M. J., Papanicolaou, G., and Ting, L.: Wave propagation in bubbly liquids at finite volume fraction, *Journal of Fluid Mechanics*, 160, 1–14, 1985.
- Cappello, J., Sauret, A., Boulogne, F., Dressaire, E., and Stone, H. A.: Damping of liquid sloshing by foams: from everyday observations to liquid transport, *Physics of Fluids*, 27, 022 103, 2015.
- 350 Cavaleri, L., Baldock, T., Bertotti, L., Langodan, S., Olfateh, M., and Pezzutto, P.: What a Sudden Downpour Reveals About Wind Wave Generation, *Procedia IUTAM*, 2018.
- Cioranescu, D. and Donato, P.: *An Introduction to Homogenization*, vol. 17, *Oxford Lecture Series in Mathematics and Its Applications*, 1999.
- Clift, R., Grace, J. R., and Weber, M. E.: *Bubbles, drops, and particles*, Courier Corporation, 1978.
- 355 Denkov, N. D., Subramanian, V., Gurovich, D., and Lips, A.: Wall slip and viscous dissipation in sheared foams: Effect of surface mobility, *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 263, 129–145, 2005.
- Dingle, N. and Lee, Y.: Terminal fallspeeds of raindrops, *Journal of applied meteorology*, 11, 877–879, 1972.
- Harrison, E. L. and Veron, F.: Near-surface turbulence and buoyancy induced by heavy rainfall, *Journal of Fluid Mechanics*, 830, 602–630, 2017.
- 360 Henderson, D., Rajan, G., and Segur, H.: Dissipation of Narrow-Banded Surface Water Waves, in: *Hamiltonian Partial Differential Equations and Applications*, edited by Guyenne, P., Nicholls, D., and Sulem, C., vol. 75 of *Fields Institute Communications*, pp. 163–183, Springer, 2015.
- Ho, D., Ascher, W., Bliven, L. F., Schlosser, P., and Gordan, E.: On mechanisms of rain-induced air-water gas exchange, *Journal of Geophysical Research, Oceans*, 105, 24 045–24 057, 2000.
- 365 Jenkins, A. and Jacobs, S.: Wave damping by a thin layer of viscous fluid, *Physics of Fluids*, 15, 1256–1264, 1997.
- Keeling, R. F.: On the role of large bubbles in air-sea gas exchange and supersaturation in the ocean, *Journal of Marine Research*, 51, 237–271, 1993.
- Kim, Y., Nam, B. W., Kim, D. W., and Kim, Y. S.: Study on coupling effects of ship motion and sloshing, *Ocean Engineering*, 34, 2176–2187, 2007.
- 370 Kudryavtsev, V., Makin, V., and Chapron, B.: Coupled sea surface-atmosphere model: 2. Spectrum of short wind waves, *Journal of Geophysical Research: Oceans*, 104, 7625–7639, 1999.
- Lamb, H.: *Hydrodynamics*, Cambridge University Press, 1916.
- Le Méhauté, B. and Khangaonkar, T.: Dynamic interaction of intense rain with water waves, *Journal of Physical Oceanography*, 20, 1805–1812, 1990.
- 375 Leighton, T.: *The acoustic bubble*, Academic press, 1994.
- Marshall, J. S.: The distribution of raindrops with size., *J. Meteor.*, 5, 165–166, 1948.
- Medwin, H. and Clay, C. S.: *Fundamentals of acoustical oceanography*, Academic press, 1997.

- Medwin, H., Kurgan, A., and Nystuen, J. A.: Impact and bubble sound from raindrops at normal and oblique incidence, *The Journal of the Acoustical Society of America*, 88, 413–418, 1990.
- 380 Medwin, H., Nystuen, J. A., Jacobus, P. W., Ostwald, L. H., and Snyder, D. E.: The anatomy of underwater rain noise, *The Journal of the Acoustical Society of America*, 92, 1613–1623, 1992.
- Mendelson, H. D.: The prediction of bubble terminal velocities from wave theory, *AIChE Journal*, 13, 250–253, 1967.
- Merlivat, L. and Memery, L.: Gas Exchange Across an Air-Water Interface: Experimental Results and Modeling of Bubble Contribution to Transfer, *Journal of Geophysical Research*, 88, 707–724, 1983.
- 385 Minnaert, M.: XVI. On musical air-bubbles and the sounds of running water, *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 16, 235–248, <https://doi.org/10.1080/14786443309462277>, 1933.
- Moghimi, S., Ramírez, J., Restrepo, J. M., and Venkataramani, S.: Mass Exchange Dynamics of Surface and Subsurface Oil in Shallow-Water Transport, *Ocean Modelling*, 128, 1–12, 2018.
- Oguz, H. N. and Prosperetti, A.: Bubble entrainment by the impact of drops on liquid surfaces, *Journal of Fluid Mechanics*, 219, 143–179, 390 1990.
- Peirson, W. L., Beyá, J. F., Banner, M. L., Peral, J. S., and Azarmsa, S. A.: Rain-induced attenuation of deep-water waves, *Journal of Fluid Mechanics*, 724, 5–35, 2013.
- Plesset, M. S. and Prosperetti, A.: Bubble dynamics and cavitation, *Annual review of fluid mechanics*, 9, 145–185, <https://doi.org/10.1146/annurev.fl.09.010177.001045>, 1977.
- 395 Prosperetti, A. and Oguz, H. N.: The impact of drops on liquid surfaces and the underwater noise of rain, *Annual Review of Fluid Mechanics*, 25, 577–602, 1993.
- Restrepo, J. M., Ramírez, J. M., and Venkataramani, S. C.: An Oil Fate Model for Shallow Waters, *Journal of Marine Science and Engineering*, 3, 1504–1543, <https://doi.org/10.3390/jmse3041504>, 2015.
- Snyder, D. E.: Characteristics of sound radiation from large raindrops, Tech. rep., Naval Postgraduate School, Monterrey CA, 1990.
- 400 Tsimplis, M. N.: The Effect of Rain in Calming the Sea, *Journal of Physical Oceanography*, 22, 404–412, 1992.
- Veron, F. and Meusniers, L.: A kinetic model for particle surface interaction applied to rain falling on water waves, *Journal of Fluid Mechanics*, 796, 767–787, 2016.
- Wolf, D. K. and Thorpe, S. A.: Bubbles and the air-sea exchange of gases in near-saturation conditions, *Journal of Marine Research*, 49, 435–466, 1991.