

Brief Communication: Lower Bound Estimates for Residence Time of Energy in the Atmospheres of Venus, Earth, Mars and Titan

Javier Pelegrina¹, Carlos Osácar¹, and Amalio Fernández-Pacheco²

¹Facultad de Ciencias. Universidad de Zaragoza. 50009 Zaragoza (Spain)

²Facultad de Ciencias and BIFI. Universidad de Zaragoza. 50009 Zaragoza (Spain)

Correspondence: Carlos Osácar (cosacar@unizar.es)

Abstract.

The residence time of energy in a planetary atmosphere, τ , recently introduced and computed for the Earth's atmosphere (Osácar et al., 2020), is here extended to the atmospheres of Venus, Mars and Titan. τ is the timescale for the energy transport across the atmosphere. In the cases of Venus, Mars and Titan, these computations are ~~mere~~-lower bounds due to a lack of some energy data. If the analogy between τ and the solar Kelvin-Helmholtz scale is assumed, then τ would also be the time the atmosphere needs to return to equilibrium after a global thermal perturbation.

1 Introduction

When the inflow, F_i , of any substance into a box is equal to the outflow, F_o , then the amount of that substance in the box, M , is constant. This constitutes an equilibrium or steady state. Then, the ratio of the stock in the box to the flow rate (in or out) is called ~~the~~-residence time and is a timescale for the transport of the substance in the box.

$$t = \frac{M}{F}. \quad (1)$$

In equation (1) it is assumed that the substance is conserved. A good example of this type is the parameter defined in atmospheric chemistry (Hobbs, 2000) as the average residence time of each individual gas, defined as (Eq. 1). M is the total average mass of the gas in the atmosphere and F the total average influx or outflux, which in time average for the whole atmosphere are equal.

In this work we extend the substance that flows in the box from matter to energy and the residence time is

$$\tau = \frac{E}{F}, \quad (2)$$

where E is the total energy in the box (a planetary atmosphere) and F the energy flux that enters or leaves it.

Here, by using (2), we estimate the average residence time of energy in several planetary atmospheres. ~~But first it is worth recalling that several authors have previously considered the energy-residence time relation in other type of problems (Meilveen, 1992, 2010; Harte, 1988).~~

~~The planetary~~ Planetary atmospheres constitute steady state problems because the storage of energy in their interior is not systematically increasing or decreasing. Several authors have previously considered the energy-residence time relation in other type of problems (Mcilveen, 1992, 2010; Harte, 1988).

25 The structure of this communication is the following: Section 2 addresses the numerator of Eq. (2) E , while Section 3 deals with the denominator F . In Section 4 the residence time of energy is considered for the Sun. In Section 5, the radiative constant is introduced and compared with the atmospheric residence timescale. In Section 6 we comment on some final points.

2 Forms of energy in a planetary atmosphere

The most important forms of energy in an atmosphere are: the thermodynamic internal energy, U , the potential energy due to the planet's gravity, P , the kinetic energy, K , and the latent energy, L , related to the phase transitions.

In a planar atmosphere, in hydrostatic equilibrium and by using the state equation for an ideal gas, these magnitudes can be written as

$$U = \int_0^{\infty} c_v T(z) \rho(z) dz = \frac{c_v}{R} \int_0^{\infty} p(z) dz, \quad (3)$$

35
$$P = \int_0^{\infty} g z \rho(z) dz = \int_0^{\infty} p(z) dz, \quad (4)$$

In expressions (3) and (4), c_v is the specific heat at constant volume, R is the gas constant and $\rho(z)$ and $T(z)$ are the density and temperature of the mixture of gases of the atmosphere, respectively. E stands for the total energy in the atmosphere:

$$E = U + P + K + L. \quad (5)$$

The sum $S = U + P$ will be called dry static energy, then

40
$$E = S + K + L. \quad (6)$$

It is important to remark that S is much bigger than the sum $K + L$. For example, for the Earth (Peixoto and Oort, 1992)

$$\frac{S}{K + L} = \frac{150}{6} = 25. \quad (7)$$

~~In ideal gases, S can be expressed as~~

$$S = (1 + \frac{c_v}{R})P = \frac{c_p}{R}P,$$

45 ~~where c_p is the specific heat at constant pressure.~~

In the case of ~~the~~ Earth's atmosphere, the four terms U , P , K , and L are known (Peixoto and Oort, 1992), so we know E . However for the atmospheres of Venus, Mars and Titan we can only compute the terms U and P and estimate S but not

Table 1. Forms of energy in planetary atmospheres

	Venus	Earth	Mars	Titan
P (J m ⁻²)	1.24E+11	7.00E+08	6.05E+06	2.63E+09
U (J m ⁻²)	4.31E+11	1.80E+09	2.10E+07	6.79E+09
S (J m ⁻²)	5.55E+11	2.50E+09	2.71E+07	9.42E+09
K (J m ⁻²)	...	1.30E+06
L (J m ⁻²)	...	7.00E+07
E (J m ⁻²)	...	2.57E+09
C_p/R	4.47	3.5	4.37	3.58

Table 2. Earth's energy comparison

<u>Units 10⁶ J m⁻²</u>	<u>Peixoto and Oort (1992)</u>	<u>Hartmann (1994)</u>	<u>Δ(%)</u>
<u>P</u>	<u>693</u>	<u>700</u>	<u>0.17</u>
<u>U</u>	<u>1803</u>	<u>1800</u>	<u>-1.01</u>
<u>L</u>	<u>63.8</u>	<u>70</u>	<u>-9.72</u>
<u>K</u>	<u>1.23</u>	<u>1.3</u>	<u>-5.69</u>
<u>E</u>	<u>2561</u>	<u>2571</u>	<u>-0.39</u>
<u>S</u>	<u>2493</u>	<u>2500</u>	<u>-0.28</u>
<u>(E - S)/E(%)</u>	<u>2.539</u>	<u>2.773</u>	

E . We have carried out these computations by performing the numerical integration (4), using the vertical data $p(z)$ shown in (Sánchez-Lavega, 2011, page 212-227). The results of E or S for each planet are shown in Table 1.

50 **In case-**

For the Earth's atmosphere, the estimates of different authors are very similar. Table 2 compares values of Peixoto and Oort (1992) and Hartmann (1994). The last row corresponds to the difference between the total energy of the Earth's atmosphere (E) and its dry static energy (S). The kinetic and latent components can be neglected in a first approximation.

The sound velocity of an ideal gas is

$$55 \quad c = \sqrt{\gamma \frac{R^*}{M} T} \quad (8)$$

where R^* is the universal constant of gases and M is the molecular mass of the gas; $\gamma = C_p/C_v$ is the adiabatic constant and T the temperature. The sound velocity can be used to estimate the ratio between K and S .

$$\frac{K}{S} \approx \left(\frac{v}{c}\right)^2 \quad (9)$$

Table 3. Wind velocity in Mars

	<u>Day</u>	<u>Night</u>	<u>Storm</u>	<u>Max during storm</u>
<u>v (m/s)</u>	<u>7</u>	<u>2</u>	<u>17</u>	<u>26</u>
<u>$K/S \approx (v/c)^2$</u>	<u>0.0009</u>	<u>0.00007</u>	<u>0.0055</u>	<u>0.0129</u>

Table 4. Fluxes of energy and residence times in planetary atmospheres

	Venus	Earth	Mars	Titan
F_i (W m^{-2})	17292 <u>17292 ± 1715</u>	509.6 <u>561 ± 9.17</u>	49.49 <u>± 3.97</u>	6.88 <u>6.88</u>
F_o (W m^{-2})	17292 <u>17292 ± 1713</u>	509.6 <u>561 ± 5</u>	49.49 <u>± 4.239</u>	6.87 <u>6.87</u>
τ (days)	371 <u>371.48 ± 26.04</u>	57 <u>53.43 ± 0.42</u>	7 <u>6.87 ± 0.41</u>	15916

60 In the case of Mars, on surface $c = 228.73 \text{ m s}^{-1}$. Table 3 contains data of winds measured by Viking probes on the surface (Sheehan, 1996, p. 194). With these data, K can be neglected in Mars. In the case of Titan, Mitchell (2011) assumes that the kinetic energy can be neglected. Based on these figures, the kinetic energy can be omitted in a first approximation for Mars and Titan.

In case when S is not much bigger than $K + L$, our results will be a poor for τ would be a lower bound. Future observations will settle-determine these numbers.

65 **3 Energy fluxes absorbed and emitted by the planetary atmospheres -Residence and residence times**

The values of ~~these~~the energy fluxes for all planets have been deduced from Read et al. (2016). For each planet, F_i and F_o represent the inflow ~~or~~and outflow of energy absorbed or emitted by the atmospheres. The so called ‘Trenberth diagrams’ are particularly suited to the identification of these fluxes.

As an example, in the case of Venus (see ~~Read et al. (2016, Figure 3)~~Read et al. (2016, Figure 6)), the fluxes absorbed by the atmosphere (F_i) are: 135 W m^{-2} from incoming solar radiation (shortwave) absorbed in the middle atmosphere, 3 W m^{-2} ~~also~~ from incoming solar radiation absorbed by the lower atmosphere, ~~;~~ and 17154 W m^{-2} of longwave flux absorbed from surface. Thus, the total influx is 17292 W m^{-2} .

The emitted fluxes (F_o) are 17132 W m^{-2} of longwave radiation to surface and 160 W m^{-2} , ~~also~~of longwave radiation emitted from atmosphere to space. The total outflux value is 17292 W m^{-2} . Analogous calculations for the rest of planets give the values for F_i and F_o shown in Table 4.

These energy fluxes were computed by Read et al. (2016) through complex and detailed numerical models. Their results coincide well with observations and have little uncertainty, so its effect on the residence time of energy can be neglected. In any case, here we have computed that uncertainty value.

For Earth, quoting (Read et al., 2016, p. 704): *"the Earth's energy budget has been quantified in the most detail and to relatively high precision (...). Even so, a number of significant uncertainties persist (...). The incoming solar flux (or solar irradiance) is known to the highest accuracy at 340.2 Wm^{-2} (Kopp and Lean, 2011) and varies the least of all the other fluxes. For the other fluxes, estimates vary as to their likely uncertainty, from around 1 Wm^{-2} for some to around 10 Wm^{-2} for the least well-characterized quantities (...). Figure 1 summarizes the recent set of estimates obtained from combinations of remote sensing and in situ measurements, together with well-validated numerical model simulations (...). These represent some of the most comprehensive studies to date that include strenuous efforts to trace the uncertainties in all of the main fluxes. (...). Figure 1 thus represents the current state of the art in deriving such an energy budget for an entire planet."* Although (Read et al., 2016) do not give exact numbers for uncertainty of energy fluxes, their references herein do. We have computed the following uncertainty values: $F_{in} = 561 \pm 9.17 \text{ Wm}^{-2} \Rightarrow \tau = 53.43 \pm 0.87 \text{ d}$, and $F_{out} = 561 \pm 5 \text{ Wm}^{-2} \Rightarrow \tau = 53.43 \pm 0.48 \text{ d}$. We note how both fluxes and residence times are extremely similar and compatible. A weighted average would give us $\tau = 53.43 \pm 0.42 \text{ d}$.

When computing the energy fluxes of Mars, Read et al. (2016) use a detailed radiative transfer model *"suggesting an uncertainty in infrared fluxes of around 6–12%"*. By using the worst case scenario of a 12% uncertainty, we obtain $F_{in} = 49 \pm 3.97 \text{ Wm}^{-2} \Rightarrow \tau = 6.87 \pm 0.56 \text{ d}$, and $F_{out} = 49 \pm 4.23 \text{ Wm}^{-2} \Rightarrow \tau = 53.43 \pm 0.59$. This gives us $\tau = 6.87 \pm 0.41 \text{ d}$. These uncertainties are reflected in Table 4.

About the energy fluxes in Venus, Read et al. (2016) state: *"energy fluxes agree with available observations to around $\pm 10\%$ "*. However, they admit that *"the energy budget presented (...) should therefore be seen as a plausible scheme that is internally self-consistent and representative of a reasonably good radiative–dynamical model of the Venus atmosphere in equilibrium"*. Assuming an uncertainty of 10% in energy fluxes, $F_{in} = 17292 \pm 1715 \text{ Wm}^{-2} \Rightarrow \tau = 371.48 \pm 36.84 \text{ d}$, and $F_{out} = 17292 \pm 1713 \text{ Wm}^{-2} \Rightarrow \tau = 371.48 \pm 36.80 \text{ d}$. This gives $\tau = 371.48 \pm 26.04 \text{ d}$.

In Titan's energy fluxes, Read et al. (2016) do not state any bound on uncertainties. However, they say (Read et al., 2016, p.711) *"energy fluxes are consistent with the measurements of Li et al. (2011) to within a few per cent, although the internal and surface fluxes are not well constrained by observations."* We can assume that the energy fluxes they present and used here, are fairly accurate with low uncertainty.

With the total energy values, E or S (in Table 1) and F (Table 4), we estimate the value of residence time of energy in the atmosphere of each planet. However, as we stressed above, strictly speaking E is only known in the Earth's case. In the other three cases, the ratio (S/F) is a lower bound for the actual residence time.

$$\frac{S}{F} \leq \frac{E}{F} = \tau. \quad (10)$$

These results and their estimated uncertainties are shown in Table 4.

4 **Energy residence** Residence time of energy in the Sun

~~The Sun also constitutes a steady state for the energy. Thermonuclear energy is produced at the center and radiant energy is emitted across the surface as a blackbody at 6000 K. Temperatures in~~ Although the physics in the solar interior greatly differs

from that of a planetary atmosphere, we have considered convenient to introduce this section because of the parallelism that exists between the atmospheric τ and the solar Kelvin-Helmholtz timescale.

$$\tau_{KH} = \frac{GM_{\odot}^2}{R_{\odot}L_{\odot}} \sim 10^7 \text{ yr.} \quad (11)$$

115 Where G is the gravitational constant and M_{\odot} , R_{\odot} and L_{\odot} stand for the solar mass, radius and radiant flux.

The Sun is in a steady state for the energy. The temperatures in its interior are not systematically increasing or decreasing. In Stix (2003) it is shown that in solar physics the Kelvin-Helmholtz (KH (Kippenhahn and Weigert, 1994)) timescale timescale (KH) corresponds to both the time that takes a photon a photon takes from the core until it goes out to leaves the surface and the time necessary for the star to return to equilibrium after a global perturbation:-

$$120 \quad \tau_{KH} = \frac{GM_{\odot}^2}{R_{\odot}L_{\odot}} \sim 10^7 \text{ yr.}$$

As L_{\odot} is the luminosity of the Sun, equation (11) is also a form of expressing the-

As τ_{KH} is the ratio between stored energy and its flux, it also can be considered as a residence time of energy in our star. This is because-

$$\tau_{KH} = \frac{GM_{\odot}^2}{R_{\odot}L_{\odot}} \sim \frac{|\text{gravitational energy}|}{L_{\odot}}.$$

125 And as in a star like the Sun, the Virial Theorem (Kippenhahn and Weigert, 1994) links-

$$|\text{gravitational energy}| = 2 |\text{total energy}|,$$

hence:-

$$\tau_{KH} \sim 2 \frac{|\text{total energy}|}{L_{\odot}}.$$

130 the Sun (for details, see Osácar et al. (2020)). Furthermore, Spruit (2000) shows that KH is the longest timescale for any solar perturbations.

5 Radiative relaxation timescale

The most simple models that can be devised for the structure of the atmosphere are the static radiative ones. But if energy transfer with the surface is taken into account, the structure produced under radiative equilibrium cannot be maintained. Convection develops spontaneously and neutralizes the stratification introduced by radiative transfer. The new radiative-convective equilibrium produces two layers. Below a certain height, the thermal structure is controlled by convective overturning and constitutes the troposphere. In this layer, the vertical profile of temperature is adiabatic. In the layer above troposphere, which

constitutes the stratosphere, the thermal structure remains close to radiative equilibrium, because radiative transfer stabilizes the stratification.

In this context, In summary, if the concept of radiative relaxation timescale is introduced. It is defined as:

$$140 \quad \tau_R = \frac{c_p p / g}{4\sigma T_{\text{eff}}^3}.$$

See, for example Wells (2012). In this expression, p is pressure, c_p is the specific heat at constant pressure, g gravity, σ analogy between the solar KH and the atmospheric τ is assumed, then τ is not only the timescale for the energy transport in the atmosphere, but also the timescale the atmosphere needs to return to equilibrium after a global thermal perturbation. Furthermore, τ is the Stefan-Boltzmann constant and T_{eff} is the blackbody effective temperature of the planet longest timescale
145 for any atmospheric perturbation.

5 Radiative relaxation timescale

In general, if a state of an atmospheric state at equilibrium is perturbed, the atmosphere uses the most efficient **procedure mechanism** at hand to neutralize it. ~~The procedure~~ Typically, this mechanism can be convective, advective or radiative. The radiative relaxation timescale, τ_R , is the time it would take to relax the perturbation by radiating the energy excess in the infrared. This timescale is often found in the bibliography and can be compared with the energy residence time τ .
150

This timescale τ_R is computed by a perturbative method, see for example Wells (2012), and the result is

$$\tau_R = \frac{c_p p / g}{4\sigma T_{\text{eff}}^3}. \quad (12)$$

In this expression, p is pressure, c_p is the specific heat at constant pressure, g is gravity, σ is the Stefan-Boltzmann constant and T_{eff} is the blackbody effective temperature of the planet.

155 Defining T_r and p_r as the temperature and pressure of the level from which most of infrared photons are emitted to the space, and T_s and p_s as the temperature and pressure at surface, if we assume that in the troposphere the temperature profile is given by a dry adiabat, then we have

$$p_r = p_s \left(\frac{T_r}{T_s} \right)^{\frac{c_p}{R}}. \quad (13)$$

Assuming this hypothesis (Pierrehumbert, 2010), for the Earth, we obtain $p_r = 670$ mb. The Earth's actual p_r is somewhat lower than this estimate because the tropospheric temperature decays less strongly with height than the dry adiabat. The result for For this value of p_r , the value of τ_R is about 22 days.
160

Due to the factor p in the numerator of Eq. 12, the value of τ_R decreases rapidly with height. So Therefore, radiation is not an efficient **procedure mechanism** to neutralize perturbations in the low troposphere. There In that region τ_R is thus very long.

A clear example is We find a clear example of these phenomena in Venus, where τ_R varies from 116 days at 40 Km (lower
165 cloud deck) to 0.5 hr at 100 Km (Sánchez-Lavega et al., 2017).

If the above mentioned analogy between the atmospheric τ and the solar scale KH is assumed, τ is the time necessary to return to equilibrium after a global perturbation, whilst τ_R is the timescale corresponding to small perturbations and $\tau > \tau_R$.

6 Final comments

In our opinion, the concept of "Residence time of energy in a planetary atmosphere" is completely original. ~~This residence time~~ Here τ has been computed for the atmospheres of Venus, Earth, Mars and Titan. In the cases of Venus, Mars and Titan, they are ~~mere~~-lower bounds due to a lack of data about kinetic and latent energies. ~~In reference (Osácar et al., 2020) the value of τ for the Earth was computed by using the data provided by Hartmann (1994).~~

The analogy between τ and the KH solar timescale seems likely, although this does not constitute a proof.

The usual radiative timescales τ_R presented by other authors (e.g. Houghton (2002), Wells (2012), Sánchez-Lavega (2011)) are calculated assuming that a small perturbation is produced in the temperature, i.e. it is a perturbative ~~evaluation~~ computation. Furthermore, it depends on the existing values of pressure and temperature at the height where it is computed. Since about the 80% of radiative flux leaving an atmosphere comes from the cold top of the highest atmospheric opaque layer, we have estimated τ_R with T_{eff} at the height of maximum emission, p_r . The obtained radiative timescale is smaller than τ . On the contrary, in the computation of the residence time of energy τ in planetary atmospheres, only global averaged planetary parameters are used.

Data availability. The data of the energies used for the estimation of residence time in the Venus, Mars and Titan atmospheres were computed with p and T from Sánchez-Lavega (2011, page 212-227). Those for the Earth's atmosphere were extracted from Peixoto and Oort (1992). The fluxes of energy for all the cases were deduced from Read et al. (2016).

Author contributions. Amalio Fernández-Pacheco conceived the idea; Carlos Osácar, Javier Pelegrina and Amalio Fernández-Pacheco wrote the paper.

Competing interests. The authors declare no conflict of interests.

References

- Harte, J.: Consider a Spherical Cow, University Science Books, 1988.
- Hartmann, D. L.: Global Physics Climatology, Academic Press, 1994.
- 190 Hobbs, P.: Introduction to Atmospheric Chemistry., Cambridge University Press, 2nd. edn., 2000.
- Houghton, J.: The Physics of the Atmosphere, Cambridge University Press, 2002.
- Kippenhahn, R. and Weigert, A.: Stellar Structure and Evolution, Springer Verlag, 1994.
- Kopp, G. and Lean, J.: A new, lower value of total solar irradiance: Evidence and climate significance., *Geophys. Res. Lett.*, 38, L01 706, <https://doi.org/10.1029/2010GL045777>, 2011.
- 195 Mcilveen, R.: Fundamentals of Weather and Climate, Chapman and Hall, 1st. edn., 1992.
- Mcilveen, R.: Fundamentals of Weather and Climate, Oxford, 2nd. edn., 2010.
- Mitchell, J. L.: Titan's transport-driven methane cycle, *Astrophys. J. Letters*, 756:L26, 1–5, <https://doi.org/10.1088/2041-8205/756/2/L26>, 2011.
- Osácar, C., Membrado, M., and Fernández-Pacheco, A.: Brief communication: Residence time of energy in the atmosphere, *Nonlinear Processes in Geophysics*, 27, 235–237, <https://doi.org/10.5194/npg-27-235-2020>, 2020.
- 200 Peixoto, J. and Oort, H.: Physics of Climate, AJP, 1992.
- Pierrehumbert, R. T.: Principles of Planetary Climate, Cambridge University Press, 2010.
- Read, P. L., Barstow, J., Charnay, B., Chelvanithhilan, S., Irwin, P. G. J., Knight, S., Lebonnois, S., Lewis, S. R., Mendonça, J., and Montabone, L.: Global energy budgets and 'Trenberth diagrams' for the climates of terrestrial and gas giant planets, *Quarterly Journal of the Royal Meteorological Society*, 142, 703–720, <https://doi.org/10.1002/qj.2704>, 2016.
- 205 Sánchez-Lavega, A.: An Introduction to Planetary Atmospheres, Taylor and Francis, 2011.
- Sánchez-Lavega, A., Lebonnois, S., Imamura, T., Read, P., and Luz, D.: The Atmospheric Dynamics of Venus, *Space Sci Rev*, 202, 1541–1616, <https://doi.org/10.1007/s11214-017-0389-x>, 2017.
- Sheehan, W.: The Planet Mars: A History of Observation and Discovery, University of Arizona Press, 1996.
- 210 Spruit, H.: Theory of solar irradiance variations, *Space Science Reviews*, 94, 113–126, <https://doi.org/10.1023/A:1026742519353>, 2000.
- Stix, M.: On the time scale of the energy transport in the Sun, *Solar Physics*, 212, 3–6, 2003.
- Wells, N. C.: Physics of the Atmosphere and the Ocean, Wiley, 3rd edn., 2012.