# **Effects of Rotation and Topography on Internal Solitary Waves Governed by the Rotating-Gardner Equation**

Karl R. Helfrich<sup>1,\*</sup> and Lev Ostrovsky<sup>2,\*</sup>

<sup>1</sup>Department of Physical Oceanography, Woods Hole Oceanographic Institution, Woods Hole, MA USA <sup>2</sup>University of Colorado Boulder, Boulder, Colorado, and University of North Carolina at Chapel Hill, Chapel Hill, NC USA \*The authors contributed equally to this work.

Correspondence: Lev Ostrovsky (lev.ostrovsky@gmail.com)

**Abstract.** Nonlinear oceanic internal solitary waves are considered under the influence of the combined effects of saturating nonlinearity, the earth's rotation, and horizontal depth inhomogeneity. Here the basic model is the extended Korteweg–de Vries equation that includes both quadratic and cubic nonlinearity (the Gardner equation) with additional terms incorporating slowly varying depth and weak rotation. The complicated interplay between these different factors is explored using an approximate

5 adiabatic approach and then through numerical solutions of the governing variable-depth, rotating Gardner model. These results are also compared to analysis in the Korteweg–de Vries limit to highlight the effect of the cubic nonlinearity. The study explores several particular cases considered in the literature that included some of these factors to illustrate limitations. Solutions are made to illustrate the relevance of this extended Gardner model for realistic oceanic conditions.

# 1 Introduction

- 10 Oceanic internal waves are an important class of nonlinear wave processes. In particular, the internal solitary waves (ISW) are the most ubiquitous type of solitons in the natural environment. These waves often propagate for long distances over several inertial periods, and the effect of Earth's background rotation is potentially significant (e.g. Farmer et al., 2009; Grimshaw et al., 2014; Helfrich, 2007; Ostrovsky and Helfrich, 2019). The large ISWs in the South China Sea are prominent examples (Zhao and Alford, 2006; Alford et al., 2010). There are also numerous remote sensing images throughout the coastal oceans that show
- 15 multiple wave packets (e.g. Jackson, 2004), indicating that the ISWs persist over periods longer than the local inertial period. It is also known that rotation destroys internal solitons due to resonant radiation of inertia-gravity waves (terminal damping; see Grimshaw et al., 1998a). The theoretical modeling of such processes often uses the rotation-modified Korteweg–de Vries equation (rKdV) derived for nonlinear waves in rotating ocean (Ostrovsky, 1978). In application to the oceanic conditions, this equation may need additional modifications, specifically for the variable depth along the propagation path. This specific case

20 was considered in Grimshaw et al. (2014), Ostrovsky and Helfrich (2019), and Stepanyants (2019).

The rotation-induced solitary wave decay can be suppressed in certain ambient shear flows wherein the sign of the rotation coefficient is changed (Alias et al., 2014). In those cases the rKdV equation supports solitary wave solutions. This interesting situation is not conside this paper.

Another important feature of oceanic solitons is that in many cases they are strongly nonlinear so that the Korteweg-de Vries

- 25 approximation (KdV) involving only the quadratic nonlinearity is inapplicable (e.g. Apel et al., 2007; Helfrich and Grimshaw, 2008; Ostrovsky and Grue, 2003; Ostrovsky and Irisov, 2017). A better approximation of the real processes can be given by the Gardner equation, that extends KdV by adding a cubic nonlinear amplitude term. It is formally applicable we en the quadratic and cubic nonlinear terms are of parable small amplitude. In that situation higher equations for dispersion and nonlinear-dispersive terms are negligible. For example, this is the case in a two-layer fluid when the layer depths are nearly
- 30 equal. Gardner solutions preserve the qualitative features of strongly nonlinear waves such as the existence of a limiting soliton amplitude at which its length infinitely increases, whereas a KdV soliton amplitude can unlimitedly increase with its width tending to zero. Moreover, in many cases the Gardner equation gives a good quantitative approximation for very solitons, beyond the formal limits of its applicability (Stanton and Ostrovsky, 1998; Pelinovsky et al., 2000; Grimshaw et al., 2004) and thus has become the phenomenological model of choice. There are numerous studies using the Gardner equation in
- 35 applications to oceanic waves including its extension for the rotating ocean (e.g. Holloway et al., 1999; Obregon et al., 2018; Talipova et al., 2015).

In this paper we make the next step by adding a sloping bottom effect to the Gardner model with rotation. Correspondingly, the results of Grimshaw et al. (2014), Ostrovsky and Helfrich (2019), and Stepanyants (2019) can be significantly modified. Since the Gardner solitons behave differently from the KdV solitons upon propagation, the growth of their amplitude is limited,

40 whereas their length can grow unlimitedly if the depth variation allows the particular, we discuss an interplay between the effects of nonlinearity, rotation, and inhomogeneity. Some realistic estimates are also given.

he work by Karczewska and Rozmej (2020) considers the analogs of the KdV and Boussinesq equations for shallow-water waves with different relative orders of the small perturbing factors: nonlinearity, dispersion, and bottom slope the matically, various terms, such as nonlinear dispersive terms, can prevail over other perturbations were, as mentioned above these additional terms are considered to remain of smaller order and are not considered here.

# 2 Rotating-Gardner equation

45

A standard model for the evolution of large-amplitude oceanic internal solitary waves is the rotating-Gardner (rG), or extended-KdV equation with rotation and variable depth h(x) (Holloway et al., 1999)

$$\frac{\partial}{\partial x} \left[ \frac{\partial \eta}{\partial t} + \left( c + \alpha \eta + \nu \eta^2 \right) \frac{\partial \eta}{\partial x} + \beta \frac{\partial^3 \eta}{\partial x^3} + \frac{c}{2} \frac{Q_x}{Q} \eta \right] = \gamma \eta.$$
(1)

- 50 Here the wave amplitude function  $\eta(x,t)$  depends on the horizontal position x and time t. The linear long wave phase speed c is found from an eigenvalue problem for the structure function  $\Phi(z)$  of a specified vertical mode. Both c and  $\Phi(z)$  are slow functions of x. The x-dependent coefficients  $\alpha$ ,  $\nu$ ,  $\beta$ , Q, and  $\gamma$  of the quadratic nonlinear, cubic nonlinear, non-hydrostatic, inhomogeneous, and rotation terms, respectively, are found as integrals over the depth of  $\Phi$  or  $\Phi'$ , and the background stratification  $\overline{\rho}(z)$  and current  $\overline{u}(z)$ . They can be found in numerous publications (e.g. Holloway et al., 1999; Grimshaw et al., 2004)
- 55 and are summarized briefly in Appendix I. When considering a spatially inhomogeneous situation it is advantageous to switch

from the (x,t) system to the (s,x) system, where

$$s = \int_{0}^{x} \frac{dx}{c} - t.$$
 (2)

If additionally  $\zeta = Q^{1/2}\eta$  is introduced, then (1) becomes

$$\frac{\partial}{\partial s} \left[ \frac{\partial \zeta}{\partial x} + \frac{\alpha}{c^2 Q^{1/2}} \zeta \frac{\partial \zeta}{\partial s} + \frac{\nu}{c^2 Q} \zeta^2 \frac{\partial \zeta}{\partial s} + \frac{\beta}{c^4} \frac{\partial^3 \zeta}{\partial s^3} \right] = \gamma \zeta.$$
(3)

60 We note that here  $\zeta^2 = Q\eta^2$  is the wave action flux. Equation (3) can be shown to have two conserved quantities

$$M = \int \zeta \, ds \quad \text{and} \quad E = \int \zeta^2 \, ds, \tag{4}$$

where the integrals are over the full *s* domain (infinite or periodic). These are, respectively, mass and energy related quantities as discussed further in the next section.

Additionally, any initial condition to (3) with  $\gamma \neq 0$  must satisfy the zero mass requirement (Ostrovsky, 1978)

65 
$$\int_{T/2}^{-T/2} \zeta(s,0)ds = 0,$$
 (5)

where s = -t at x = 0, and T is the length of the time domain.

In the absence of rotation,  $\gamma = 0$ , (3) reduces to the Gardner equation that has the solitary wave solutions

$$\zeta = \frac{a}{1 + B\cosh[\sigma(s - \kappa x)]},\tag{6}$$

described by the parameter B. Here

70 
$$a = \frac{Q^{1/2}\alpha}{\nu}(B^2 - 1), \quad \sigma^2 = \frac{c^2\alpha a}{6\beta Q^{1/2}}, \quad \kappa = \frac{\beta\sigma^2}{c^4}.$$
 (7)

The amplitude, A, in terms of  $\eta (= \zeta Q^{-1/2})$  is

$$A = \frac{aQ^{-1/2}}{1+B} = \frac{\alpha}{\nu}(B-1).$$
(8)

From the mass constraint (5), a solitary wave initial condition requires the addition of a constant pedestal,

$$d = -\frac{1}{T} \int_{T/2}^{-T/2} \zeta(s,0) \ ds,$$

75 with  $\zeta$  given by (6).

There are three families of steady solitary wave solutions given by (6)-(8) (Grimshaw et al., 1999). When  $\nu < 0$ , solitary wave solutions require 0 < B < 1 and have polarity  $\alpha A > 0$ . They approach the sech<sup>2</sup> KdV solitary wave as  $B \rightarrow 1$  and as  $B \rightarrow 0$  the solution approaches the maximum amplitude,  $A_{max} = -\alpha/\nu$ , flat-top wave. When  $\nu > 0$ , solitary wave solutions

require  $B^2 > 1$ , and there are two solution branches. For B > 1,  $\alpha A > 0$  and there is no limit on the wave amplitude. The sech<sup>2</sup>

KdV wave is recovered as B → 1 from above and for B ≫ 1 the solution approaches the sech solution of the modified KdV equation (i.e. the Gardner equation with α = 0). The third branch occurs for B < -1 with the wave polarity αA < 0. In this case, solitary wave amplitude has a minimum amplitude A<sub>min</sub> = -2α/ν obtained as B → -1 from below. This limiting wave has an algebraic structure and for B ≪ -1 the solution again approaches the sech wave of modified KdV equation. For ν > 0 there are also localized pulsating traveling wave solutions (breathers) that have total negative mass between zero and the mass of the limiting solitary wave at B = -1 (Pelinovsky and Grimshaw, 1997).

### **3** Adiabatic evolution of solitary waves

Assuming that  $\zeta \to 0$  for  $|s| \to \infty$  and integrating (3) gives

$$\frac{\partial\zeta}{\partial x} + \frac{\alpha}{c^2 Q^{1/2}} \zeta \frac{\partial\zeta}{\partial s} + \frac{\nu}{c^2 Q} \zeta^2 \frac{\partial\zeta}{\partial s} + \frac{\beta}{c^4} \frac{\partial^3 \zeta}{\partial s^3} = -\gamma \int_s^\infty \zeta ds'.$$
(9)

Multiplication by  $\zeta$  and integration in s from  $-\infty$  to  $\infty$  gives

90 
$$\frac{d}{dx}\left(\int_{-\infty}^{\infty}\zeta^2 ds\right) = -2\gamma\int_{-\infty}^{\infty}\left(\int_{s}^{\infty}\zeta ds'\right)ds = -\gamma\left(\int_{-\infty}^{\infty}\zeta ds\right)^2.$$
 (10)

Solution of the rG equation (9) implies that the right side of (10) is zero and E is a constant. However, progress can be made if we start with a solitary wave from (6), assume that the inhomogeneous and rotational effects are very weak such that the evolving wave remains a solitary wave, but with slowly varying amplitude, and take the limits of integration to contain only the evolving solitary wave. With the solitary wave solution (6) written as

95 
$$\zeta = a\mathcal{F}(y), \quad \mathcal{F} = \frac{1}{1 + B\cosh(y)}, \text{ and } y = \sigma s,$$
 (11)

equation (9) gives

$$\frac{d}{dx}\left(\frac{a^2}{\sigma}\mathcal{I}_2\right) = -\gamma \frac{a^2}{\sigma^2}\mathcal{I}_1^2,\tag{12}$$

where

$$\mathcal{I}_n = \int_{-\infty}^{\infty} \mathcal{F}^n dy, \qquad n = 1, 2.$$
(13)

100 The result, (12), is a statement for the variation of the energy,

$$E_w = \frac{a^2}{\sigma} \mathcal{I}_2,\tag{14}$$

of the slowly evolving solitary wave. In the absence of rotation  $E_w$  is conserved; however, the wave mass,

$$M_w = \frac{a}{\sigma} \mathcal{I}_1,\tag{15}$$

is not necessarily conserved. Changes in  $M_w$  are compensated by the formation of a trailing shelf. For example, in the typical

105 case with ν < 0 of a solitary wave approaching a point of polarity reversal, α = 0, the wave mass increases in magnitude (Grimshaw et al., 1998b). Thus the trailing shelf must have the sign opposite to the wave polarity. When ν > 0, the situation is more subtle, but again any variations in M<sub>w</sub> are compensated by a shelf (Grimshaw et al., 1999; Nakoulima et al., 2004). In a homogeneous, rotating environment both E<sub>w</sub> and M<sub>w</sub> decrease with x and are compensated by the trailing wave radiation (Grimshaw et al., 1998a). With both inhomogeneity and rotation the energy will decrease with x, but the wave mass may
110 increase or decrease depending on the interplay between these two effects.

Note that for the KdV equation and with Q defined as in (A.3d),  $\rho_0 E_w$  is the solitary wave energy,  $\iint pu \, dz \, dt$ . Here p and u are the first-order, wave-induced pressure and horizontal velocity fields. With the addition of the cubic nonlinear term in the Gardner equation,  $E_w$ , is not exactly the wave energy, but is still a good measure of it. The actual wave mass is  $Q^{-1/2}M_w$ .

#### 3.1 Rotating-KdV equation

115 For later reference we first consider the adiabatic theory for the rotating-KdV equation ( $\nu = 0$ ) from Grimshaw et al. (2014). The solitary wave solution is found from (6) and (8) with B = 1 as

$$\zeta = a\mathcal{F}, \qquad \mathcal{F} = \operatorname{sech}^2(y), \qquad \sigma^2 = \frac{c^2 \alpha a}{12\beta Q^{1/2}}, \qquad a = AQ^{1/2}.$$
(16)

From (13),  $\mathcal{I}_1 = 2$  and  $\mathcal{I}_2 = 4/3$ , and (12) gives

$$\frac{d}{dx}\left(\frac{4}{3}\frac{a^2}{\sigma}\right) = -4\gamma \frac{a^2}{\sigma^2}.$$
(17)

120 This can be written as

$$\frac{d}{dx}\mathcal{A}^{3/2} = -3\gamma w^2 \mathcal{A}, \quad \mathcal{A} = wa, \quad w = \left(\frac{12\beta Q^{1/2}}{\alpha c^2}\right)^{1/3}.$$

When integrated this gives

$$\frac{A}{A_0} = \left(\frac{Q_0}{Q}\right)^{1/2} \frac{a}{a_0} = \left(\frac{Q_0}{Q}\right)^{1/2} \frac{w_0}{w} \left[1 - (w_0 a_0)^{-1/2} \int_0^x \gamma w^2 dx'\right]^2.$$
(18)

The 0 subscript indicates the initial value at x = 0.

125 In the absence of rotation,  $\gamma = 0$ , conservation of wave action gives

$$\frac{A}{A_0} = \left(\frac{Q_0}{Q}\right)^{1/2} \frac{w_0}{w} = \left(\frac{Q_0^2 \beta_0}{\alpha_0 c_0^2} \frac{\alpha c^2}{Q^2 \beta}\right)^{1/3}.$$
(19)

For a homogeneous, rotating environment (18) gives

$$\frac{A}{A_0} = \left[1 - \frac{x}{X_{eO}}\right]^2, \quad X_{eO} = \frac{c}{\gamma} \left(\frac{\alpha A_0}{12\beta}\right)^{1/2}.$$
(20)

The KdV solitary wave is completely extinguished by radiation of inertia-gravity waves in the finite distance  $X_{eO}$ .

#### 130 3.2 Rotating-Gardner equation

When  $\nu \neq 0$ , the Gardner solitary wave solution (11) and (13) give

$$\mathcal{I}_1 = \frac{4}{(B^2 - 1)^{1/2}} \,\mathcal{T}(B),\tag{21}$$

$$\mathcal{I}_2 = \frac{2}{B^2 - 1} - \frac{4}{(B^2 - 1)^{3/2}} \mathcal{T}(B),$$
(22)

where

135 
$$\mathcal{T}(B) = \tan^{-1}\left(\frac{B-1}{\sqrt{B^2-1}}\right) = \operatorname{sgn}(B) \tan^{-1}\sqrt{\frac{B-1}{B+1}}.$$
 (23)

Substituting (7), (21) and (22) into (12) gives

$$\frac{d}{dx} \left[ \left( \frac{24\beta \alpha^2 Q^2}{c^2 \nu^3} \right)^{1/2} \left[ (B^2 - 1)^{1/2} - 2\mathcal{T}(B) \right] \right] = -96 \frac{\gamma \beta Q}{c^2 \nu} \mathcal{T}^2(B).$$
(24)

This equation can be integrated to obtain B(x), hence A(x) from (8). Since  $B \to 1$  as  $\nu \to 0$ , the equation remains regular for cases with  $\nu(x)$  changing sign. Note also that it remains real for  $\nu < 0$  and 0 < B < 1 since  $\tan^{-1}(iy) = i \tanh^{-1}(y)$ .

140 While it is not necessary to make the sign of  $\nu$  explicit in integrating (24), for  $\nu < 0$  (0 < B < 1), it can be written as

$$\frac{\sqrt{1-B^2}}{B}\frac{dB}{dx} = \frac{1}{2}\frac{d}{dx}\left[\ln\left(\frac{\beta Q^2 \alpha^2}{c^2 |\nu|^3}\right)\right]\left(2\tanh^{-1}\sqrt{\frac{1-B}{1+B}} - \sqrt{1-B^2}\right) + 8\gamma\left(\frac{6\beta|\nu|}{c^2\alpha^2}\right)^{1/2}\left(\tanh^{-1}\sqrt{\frac{1-B}{1+B}}\right)^2.$$
(25)

and for  $\nu > 0$  ( $B^2 > 1$ )

$$\frac{\sqrt{B^2 - 1}}{B} \frac{dB}{dx} = -\frac{1}{2} \frac{d}{dx} \left[ \ln\left(\frac{\beta Q^2 \alpha^2}{c^2 \nu^3}\right) \right] \left(\sqrt{B^2 - 1} - 2 \tan^{-1}\left(\frac{B - 1}{\sqrt{B^2 - 1}}\right) \right) \\ - 8\gamma \left(\frac{6\beta\nu}{c^2\alpha^2}\right)^{1/2} \left[ \tan^{-1}\left(\frac{B - 1}{\sqrt{B^2 - 1}}\right) \right]^2.$$
(26)

145

As discussed above, in a non-rotating system the right-hand side of (24) is zero, and the conservation of wave energy gives

$$E_{w0} = \left(\frac{24Q^2\alpha^2\beta}{c^2\nu^3}\right)^{1/2} \left[ (B^2 - 1)^{1/2} - 2\mathcal{T}(B) \right],\tag{27}$$

where  $E_{w0}$  is a constant evaluated at x = 0. This can be solved to give B(x).

Radiation decay in a homogeneous environments (where  $c, \alpha, ...$  are constants) was recently considered by Obregon et al. (2018). Here we briefly redevelop their decay result in our variables for clarity. In a homogeneous environment the first term on the right-hand-side of (25) or (26) is zero. Then the distance,  $X_{eG}$ , to complete radiation decay in the rG equation is found by integration of (25) or (26), with the first term on the right side set to zero, from x = 0 to  $X_{eG}$ , where  $B = B_0$  and  $B_e$ , respectively. When  $A_0\alpha > 0$ ,  $B_0 > 0$ , the solitary wave decays from an initial amplitude  $A_0 = \alpha \nu^{-1}(B_0 - 1)$  to zero when



Figure 1.  $X_{eG}/X_{eO}$  as a function of  $B_0$  for the three Gardner solitary wave regimes:  $\nu < 0$  for  $0 < B_0 < 1$  and  $\nu > 0$  for  $B^2 > 1$ .

 $B_e = 1$ , regardless of the sign of  $\nu$ . While for  $\nu > 0$  and  $A_0 \alpha < 0$ ,  $B_0 < -1$  and the solitary wave decay can be followed only 155 to the limiting amplitude,  $A_{lim} = -2\alpha/\nu$ , where  $B_e = -1$ . These considerations give

$$\frac{X_{eG}}{X_{eO}} = \frac{\sqrt{2}}{8} \begin{cases} \mathcal{I}_B(B_0, 1) | B_0 - 1|^{-1/2}, & B_0 > 0, \\ \mathcal{I}_B(B_0, -1)(1 - B_0)^{-1/2}, & B_0 < -1, \end{cases}$$
(28)

where

$$\mathcal{I}_{B}(B_{0}, B_{e}) = \begin{cases}
\int_{B_{0}}^{B_{e}} \frac{\sqrt{1-B^{2}}}{B} \left( \tanh^{-1} \sqrt{\frac{1-B}{1+B}} \right)^{-2} dB, \quad \nu < 0, \\
-\int_{B_{0}}^{B_{e}} \frac{\sqrt{B^{2}-1}}{B} \left( \tan^{-1} \sqrt{\frac{B-1}{B+1}} \right)^{-2} dB, \quad \nu > 0.
\end{cases}$$
(29)

Again,  $X_{eO}$  is the rKdV equation decay length (20) evaluated using  $|A_0\alpha|$  when  $B_0 < -1$ .

160

Figure 1 shows  $X_{eG}/X_{eO}$  versus  $B_0$  for all three wave regimes. For  $\nu < 0$  where  $0 < B_0 < 1$ ,  $X_{eG}/X_{eO} \approx 1$  for  $0.2 < B_0 < 1$ . 1. There is a slight minimum of 0.9924 at  $B_0 = 0.55$ . As  $B_0 \rightarrow 0$  the ratio increases to  $X_{eG}/X_{eO} = 1.1842$  at  $B_0 = 10^{-14}$  and appears to approach a finite limit as  $B_0 \rightarrow 0$ . For  $B_0 > 1$  ( $\nu > 0$ ),  $X_{eG}/X_{eO}$  increases monotonically from one with  $B_0$ , but remains O(1) even for  $B_0 = 10$ . Similar behavior is found for  $B_0 < -1$  ( $A_0\alpha > 0$ ), although it should be remembered that in this regime the adiabatic theory gives an amplitude only until the limiting wave at  $B_e = -1$  is reached.

165

Figure 2 shows examples of normalized wave amplitude  $A/A_0$  as functions of  $x/X_{eO}$  for several values of  $B_0$  in each wave regime. For  $\nu < 0$  and  $B_0 \gtrsim 0.1$  (Figure 2a) the amplitude decay closely follows the solution (20) for the rKdV equation, but as  $B_0$  decreases the initial amplitude decay rate slows. As mentioned above, for  $\nu > 0$  and  $B_0 < -1$  (Figure 2c) the decay can only be followed until the limiting wave amplitude is reached.

#### 4 Comparisons with rG numerical solutions

170 In this section the adiabatic theory (24) is compared with full numerical solutions of the rG equation (3). Example cases employ spatially uniform stratifications and inhomogeneous effects are introduced by variations in the total water depth. The numerical solutions of (3) are found using a de-aliased pseudo-spectral scheme in *s* with a third-order Runge-Kutta integration in *x*. The relations for the coefficients,  $\alpha(x)$ ,  $\nu(x)$ , ..., are given in Appendix A.



Figure 2.  $A/A_0$  versus  $x/X_{eO}$  for homogeneous conditions. a)  $\nu < 0$  for  $B_0 = [10^{-1}, 10^{-2}, 10^{-4}]$  (solid lines from left to right). The dashed line is the rKdV equation solution (20). b)  $\nu > 0$  and  $B_0 = [1.1, 5, 10]$  (from left to right). c)  $\nu > 0$  and  $B_0 = [-2, -5, -10]$  (from left to right).

# 4.1 Rotating, homogeneous evolution

175 In the homogeneous case where the coefficients c,  $\alpha$ , ... are constants it is convenient to reduce (3) to an equation with only one parameter by introducing the scaling

$$u = \frac{\zeta}{U}, \quad \tau = \frac{s}{T}, \quad \xi = \frac{x}{L}$$

where

$$U = \frac{\alpha}{|\nu|}, \quad T = \left(\frac{\beta|\nu|}{c^2\alpha^2}\right)^{1/2}, \quad L = \frac{c}{\alpha^3} \left(\beta|\nu|^3\right)^{1/2}.$$

180 This change of variables gives

$$\frac{\partial}{\partial\tau} \left[ \frac{\partial u}{\partial\xi} + u \frac{\partial u}{\partial\tau} + \operatorname{sgn}(\nu) u^2 \frac{\partial u}{\partial\tau} + \frac{\partial^3 u}{\partial\tau^3} \right] = \epsilon u \quad \text{with} \quad \epsilon = \gamma \beta \frac{\nu^2}{\alpha^4}.$$
(30)



Figure 3. Comparison of the rG adiabatic radiation decay theory for homogeneous conditions (black lines) and rG numerical solutions for a)  $B_0 = 0.3$ , b)  $B_0 = 10^{-4}$ , c)  $B_0 = 5$ , and d)  $B_0 = -4$ . In (a) and (b) red (blue) indicates numerical solutions of the rG equation (30) for  $\epsilon = 2.5 \times 10^{-5} (2.5 \times 10^{-3})$ . In (c) and (d) red (blue) indicates  $\epsilon = 2.5 \times 10^{-3} (0.25)$ .

The solitary wave solution (6)-(8) and the adiabatic radiation decay results carry through after taking  $c = Q = \alpha = \beta = 1$ ,  $\nu = \pm 1$ , and  $\gamma = \epsilon$ . In these variables  $X_{eO} = \epsilon^{-1} (|A_0|/12)^{1/2}$  and  $A_0 = \operatorname{sgn}(\nu)(B_0 - 1)$ .

Figure 3a-d shows comparison of the scaled amplitude A/A<sub>0</sub> versus ξ/X<sub>eO</sub> from the adiabatic theory and full numerical solutions of the rG equation (30) with B<sub>0</sub> = [0.3, 10<sup>-4</sup>, 5, -4], respectively and 2.5 × 10<sup>-5</sup> ≤ ε ≤ 0.25 as indicated. (The small oscillations in the amplitude are due to the periodic boundary conditions used in the numerical solutions that allowed the radiated waves to re-enter the domain upstream of the solitary wave.) For B<sub>0</sub> = 10<sup>-4</sup> and 0.3 (ν < 0) the agreement between the adiabatic theory and the rG solutions is quite good for ε = 2.5 × 10<sup>-5</sup>. However, for ε = 2.5 × 10<sup>-3</sup> there is disagreement. Similarly, for B<sub>0</sub> = -4 and 5 the agreement also declines with increasing ε, although in these examples the agreement for ε = 2.5 × 10<sup>-3</sup> is good. Increasing ε generally results in slower amplitude decay. The exception is B<sub>0</sub> = 10<sup>-4</sup>, where the initial decay is more rapid. This rapid decay for near-maximal waves was also found by Obregon et al. (2018) which they attributed to a structural instability of large-amplitude, flat-top solitary waves.

The complicated evolution of the decaying solitary wave and the trailing radiation is illustrated in Figures 4a and b. Figure 4a is the  $B_0 = 10^{-4}$  and  $\epsilon = 2.5 \times 10^{-5}$  example in Figure 3b. As the initial solitary wave decays, the trailing radiation itself steepens to form a group of solitary-like waves that also decay by radiation damping. Over larger distances this radiation will likely organize into one or more nonlinear wave packets as found by Grimshaw and Helfrich (2008) for the rotating-KdV equation and Whitfield and Johnson (2015) in the rotating-Gardner equation.



Figure 4. Numerical solution of (30) with a)  $\epsilon = 2.5 \times 10^{-5}$  and  $B_0 = 10^{-4}$  and b)  $\epsilon = 2.5 \times 10^{-3}$  and  $B_0 = -4$ . The solid bar indicates the initial wave amplitude  $A_0 = \text{sgn}(\nu)(B_0 - 1)$ . Only a portion of the full  $\xi$  domain is shown. The locations of each time series are indicated by the vertical axis.

The example in Figure 4b has  $B_0 = -4$  and  $\epsilon = 2.5 \times 10^{-3}$  (see Figure 3d). For these parameters the evolution is further complicated since the radiation decay ceases at  $\xi \approx 50 \ (= 0.21 X_{eO})$  when the wave amplitude decays to the limiting ampli-200 tude  $A_{lim} = -2\alpha/\nu = -2$  in these scaled variables. The wave then rapidly forms what appears to be a small solitary wave of reversed polarity ( $B_0 > 1$ ) and a trailing wave packet that has characteristics similar to the breather solutions of the Gardner equation. However, this packet subsequently disintegrates due to rotational effects. The complicated nature of the wave evolution with rotation in homogeneous conditions suggests even more interesting features when both rotation and inhomogeneous effects are active.

#### 205 4.2 Combined inhomogeneous and rotation effects

Ostrovsky and Helfrich (2019) showed that for the rKdV equation the competition between extinction by radiation decay and at the point of polarity reversal,  $\alpha = 0$ , could be characterized by the ratio of the inhomogeneous and rotation terms of (1)

$$C \approx \frac{c}{\gamma} \frac{Q_x}{Q} \frac{\eta_x}{\eta} \sim \frac{c}{\gamma L \Delta} = \frac{X_{eO}}{L},\tag{31}$$

where L is the length scale over which inhomogeneous term Q varies, say the distance to the  $\alpha = 0$  location.  $\Delta$  is the solitary 210 wave scale, taken above to be the KdV wave scale  $\Delta_{KdV} = (12\beta/A_0\alpha)^{1/2}$ . For  $C \gg 1$  inhomogeneous effects dominate and conversely rotational decay dominates for  $C \ll 1$ . Alternatively  $\Delta$  might taken to be the Gardner solitary wave scale  $\Delta_G = \Delta_{KdV} [2/(1+B_0)]^{1/2}$ . However, for  $B_0 > 0$  and not too large, the term in parentheses is O(1) and (31) is a reasonable scaling estimate. One could simply define  $C = X_{eG}/L$ , but since  $X_{eG}/X_{eO} \approx 1$  for  $0 < B_0 < 10$  (see Figure 1) this also gives C from (31). The exception is for situations with  $B_0 < -1$ , since  $X_{eG}/X_{eO}$  can be much less than one.

215

To illustrate the combined effects of inhomogeneity and rotation a two-layer Boussinesq stratification with upper layer depth  $h_1$ , variable lower layer depth  $h_2(x)$ , reduced gravity  $g' = g(\rho_2 - \rho_1)/\rho_1$ , and Coriolis frequency f will be considered. Here  $\rho_1$  and  $\rho_2$  are the densities of the upper and lower layers, respectively. Relations for the coefficients  $\alpha(x)$ ,  $\nu(x)$ , etc. are given in (A.6). Note that  $\nu < 0$  for two-layer stratifications. Thus only the  $0 < B_0 < 1$  branch of solitary wave solutions is possible. The wave polarity is given by the sign of  $\alpha$  with  $\alpha < 0$  (> 0) for  $h_1/h_2 < 1$  (> 1).

220 The bottom slope will be taken constant and the lower layer depth given by

$$\frac{h_2(x)}{h_{20}} = 1 - \left(1 - \frac{h_1}{h_{20}}\right) \frac{x}{L},\tag{32}$$

where  $h_{20} = h_2(0)$  and L is the distance from the origin to the critical point where  $\alpha = 0$  (i.e.,  $h_1 = h_2$ ). The case  $h_1/h_{20} < 1$ corresponds to an initial solitary wave with  $A_0 < 0$  propagating from deep to shallower water. Propagation of a positive wave,  $A_0 > 0$ , from shallow to deeper water occurs for  $h_1/h_{20} > 1$ . While the adiabatic solutions can be obtained only up to the 225 critical point,  $x \leq L$ , numerical solutions of the rG equation are found beyond the critical depth. In the deep-to-shallow situation the bottom slope is continued until  $h_2 = h_1/2$ , beyond which  $h_2$  is constant over a shelf region.

Note that the linear bottom slope in (32) will not allow for the interesting, but rather special, topographic conditions Stepanyants (2019) found for the rKdV regime. In those cases rotation and inhomogeneity effects can be balanced and give a constant soliton amplitude. However, there is no balance for fluid velocity and, more important, soliton energy, which still decreases due to radiation.

The evolution of the wave amplitude A(x) is shown in Figure 5a for a deep to shallow case for a representative oceanographic situation with  $h_1 = 50$  m,  $h_{20} = 450$  m, g' = 0.005 m s<sup>-2</sup>,  $f = 10^{-4}$  s<sup>-1</sup>, and  $A_0 = -25$  m. For this initial wave  $B_0 = 0.764$ so that effects of the cubic nonlinearity are present, but do not dominate initially. Slope lengths L = 50, 100, 200, and 400 kmare considered and give C = 3.46, 1.73, 0.86, and 0.43, respectively, from (31). The solid lines show the rotating adiabatic 235 theory and the dashed line shows the f = 0 solution (equivalent for all L when plotted against x/L). As anticipated from the values of C, rotational decay effects increase significantly as L increases with complete, or nearly complete, extinction for the two longer slopes. However, even for L = 50 km, rotation causes a noticeable reduction in wave amplitude compared to the

non-rotating solution.

230

Figure 5b shows the wave energy,  $E_w(x)$  from (14), for the same parameters. The ratio  $E_w/E_{w0}$  is a measure of the fraction of initial wave energy that remains in the evolving solitary wave, with the difference lost to the trailing inertia-gravity wave 240 radiation. Even the shortest slope, L = 50 km, where the effects of rotation were relatively weak, more than half of the initial wave energy is lost to inertia-gravity wave radiation by  $x/L \approx 0.8$ .

The variation of wave mass,  $M_w$  from (15), is plotted in Figure 5c. In all cases rotation causes an initial reduction in wave mass, which is compensated in the trailing radiation and emerging shelf (with the same mass sign as the initial wave). The mass goes to zero for the two longer slopes prior to the critical point, while for the two shorter slopes the wave mass increases 245



Figure 5. Adiabatic theory for wave propagation from deep to shallow in a two-layer system with  $h_1 = 50$  m,  $h_{20} = 450$  m, g' = 0.005 m s<sup>-2</sup>,  $f = 10^{-4}$  s<sup>-1</sup>, and  $A_0 = -25$  m.  $A/A_0$ ,  $E_w/E_{w0}$ , and  $M_w/|M_{w0}|$  are shown as functions of x/L. The solid lines are for L = 50 (black), 100 (blue), 200 (red), and 400 km (magenta). The dashed lines are for f = 0. The dots are from solutions of (3).

rapidly as the critical point is approached, similar to the non-rotating result. However, increased slope length (i.e. rotational effects) delays this growth with consequences for the magnitude of the trailing shelf and subsequent wave structure transmitted onto the topographic shelf (see below).

250

255

Also shown in Figures 5a and b are results from the numerical solutions to the rG equation (3). The energy of the solitary wave in the rG model is found by integrating  $Q\eta^2$  in a small region encompassing the solitary wave that does not incorporate appreciable trailing radiation. The agreement is quite good for all the cases, except for  $x/L \gtrsim 0.8$  and L = 50 and 100, where the amplitude from the rG numerical calculation does not decay as rapidly as the adiabatic model. This is consistent with previous studies without rotation (Grimshaw et al., 1998b, 1999). As the point of polarity reversal is approached the wave elongates to form a rarefaction, and the trailing shelf with opposite polarity emerges. Figure 5 shows that the disagreement is associated with rapid changes in the wave amplitude and mass. In this region that assumption of slow variation of the inhomogeneity fails.

Two examples of the full rG solutions for L = 50 km and 200 km are shown in Figure 6 and the rG solutions at x/L = 1.2 for all five cases in Figure 5 are compared in Figure 7. Note that  $h_2 = h_1/2$  for  $x/L \ge 1.0625$  on the shelf. Again, even for L = 50 km rotation leads to a clear effect on the wave signal transmitted on to the shelf as the leading crest of the (weak)



Figure 6. rG numerical solutions for  $h_1 = 50$  m,  $h_{20} = 450$  m, g' = 0.005 m s<sup>-2</sup>,  $f = 10^{-4}$  s<sup>-1</sup> and  $A_0 = -25$  m. a) L = 50 km and b) L = 200 km. The wave amplitude  $\eta(x,t)$  is shown at x/L as function of normalized shifted time,  $-s(g'/H)^{1/2}$ , where  $H = h_1 + h_{20}$ . The initial wave amplitude is indicated by the scale on the lower right of each panel.

260 trailing inertia-gravity wave also steepens to form a second transmitted wave packet. The second packet is consistent with the breaking criteria obtained by Grimshaw et al. (2012) and similarly noted in Grimshaw et al. (2014). Further increases in rotation effects lead to an additional transmitted packet for L = 100 km. For the two longest slopes, L = 200 and 400 km the transmitted signal becomes increasingly disorganized. Figure 6b illustrates the evolution leading to this outcome. In this example the initial solitary wave is extinguished before the critical point is reached. However, the trailing inertia-gravity wave steepens to produce a solitary wave that is itself scattered through the critical point. The interaction of the scattered signal with 265 the trailing radiation gives rise to the disorganization. Recall that integration for calculating the energy of the leading wave,  $E_W = \int Q\eta^2 ds$ , shown in Figure 5 by the red solid dots, only includes the leading solitary wave and not the secondary solitary

wave that emerges from the trailing radiation.

270

In the examples above the cubic nonlinearity was not an essential feature of the evolution. Indeed the wave evolution is qualitatively similar to the rotating-KdV solutions in Figure 2 of Ostrovsky and Helfrich (2019). In Figure 8 another example with  $h_1 = 100 \text{ m}, h_{20} = 200 \text{ m}, g' = 0.01 \text{ m s}^{-2}, f = 10^{-4} \text{ s}^{-1}$ , and L = 40 km is shown. Initial wave amplitudes  $A_0 = -10$ m and -45m, corresponding to  $B_0 = 0.788$  and 0.0478, respectively. The larger wave is very close to the limiting amplitude  $A_{lim} = -47.06$  m. The competition parameter C = 4.56 and 9.68, respectively. The left column of Figure 8 shows the rG adiabatic solutions and for comparison the right column shows the adiabatic solutions obtained from the rKdV theory (18).



Figure 7.  $\eta/H$  at x/L = 1.2 for the parameters of Figure 5. The bottom curve is the non-rotating case with L = 50 km. The top four include rotation and have L = 50, 100, 200, and 400 km from bottom to top. The curves are offset by 0.05.



Figure 8. Adiabatic theory for wave propagation from deep to shallow in a two-layer system with  $h_1 = 100$  m,  $h_{20} = 200$  m, g' = 0.01 m s<sup>-2</sup>,  $f = 10^{-4}$  s<sup>-1</sup>, and L = 40 km. The left column shows  $A/A_0$ ,  $E_w/E_{w0}$  and  $M_w/|M_{w0}|$  for the rotating-Gardner theory. The right column shows the equivalent rotating-KdV solutions. The solid (dashed) lines are with (without) rotation for  $A_0 = -10$  m (black) and -45 m (blue). The open (solid) symbols are from corresponding numerical solutions of (3) with (without) rotation.



Figure 9.  $\eta/H$  at x/L = 1.5 for the parameters of Figure 8. a)  $A_0 = -10$  m and b)  $A_0 = -45$  m. The solid (dashed) lines are with (without) rotation.

Now the differences between the rG and rKdV solutions are substantial, especially for the large wave. The rKdV solution has this larger wave decaying more slowly than the smaller wave, while the rG model shows just the opposite. Similar to above, rotation, even for this relatively short slope (and hence weak rotational effect), causes significant energy loss in the primary wave as it climbs the slope. For A<sub>0</sub> = -45m, the mass M<sub>w</sub> remains finite at the critical point, indicating the generation of a relatively weak trailing shelf. The agreement between the rG adiabatic model and the full rG solutions for A(x) is not very good, although the qualitative prediction that the large wave should decay more rapidly is found in both cases. The origin of the disagreement is likely due to the lack of separation between the wave scales and scale of the inhomogeneity.

The transmitted signals at x/L = 1.5 from the full rG numerical solutions with and without rotation are shown in Figure 9. For both initial wave amplitudes rotation leads to significant changes in the transmitted signal. This is especially true for A<sub>0</sub> = -45 m, where the transmitted packet without rotation is replaced by a single, broad wave emerging onto the constant
depth shelf with a much weaker trailing signal. However, on the shelf A<sub>lim</sub>/H = 0.0745, so that this leading wave must further adjust and is also subject to continued rotational decay.

#### 5 Concluding Remarks

290

In this paper we have continued a series of studies of nonlinear internal waves of a moderate amplitude in the shallow, stratified areas of the ocean (see the references in the Introduction). Based on the classical Korteweg-de Vries equation, we added the main factors making the analysis closer to the physical reality: cubic nonlinearity, Earth's rotation, and sloping bottom. Interplay of these factors makes the problem rather complicated, both physically and mathematically. To better explain the qualitative effect of each of them, first we briefly reproduce the effect of rotation in the medium with quadratic nonlinearity (KdV with rotation), then that with both quadratic and cubic nonlinearities (Gardner with rotation) and, as the main content of this paper, the joint effect of rotation and inhomogeneity in the Gardner equation. The specific qualitative effect of the latter is

- 295 the limiting soliton amplitude and the corresponding increase of the wavelength so that the topography effect becomes especially important. Along with the approximate adiabatic approach, a direct numerical study of the rG equation was performed and confirmed the adiabatic theory and highlighted its limitations. This combined approach allowed us to demonstrate a rather complicated behavior of shoaling internal solitons. For example, whereas the soliton energy always decreases due to the radiation losses, the displacement amplitude and mass can increase in a shoaling wave at a finite distance due to the decrease
- 300 of total depth. In turn, the oscillating tail reveals a complicated behavior that includes, in particular, formation of nonlinear wave packets as in Grimshaw and Helfrich (2008), followed by an even more complex evolution. Another important point of bifurcation is that of wave passage through the condition  $h_1 = h_2$  when the polarity of a soliton is changed. According to the estimations, the interplay between the topography and rotation effects does exist under realistic oceanic conditions.

Future research should include further comparison of theory with observational results in different oceanic environments and extending the above results to the strongly nonlinear waves with rotation.

## Appendix A: Coefficients of the rotating-Gardner equation

The eigenvalue problem for the linear long wave phase speed  $c_n$  and vertical structure function for the isopycnal displacement  $\Phi_n(z)$  of vertical mode n is, in for the Boussinesq and rigid-lid limits (after dropping the subscript n),

$$\frac{d}{dz}\left[\left(c-\bar{u}\right)^{2}\frac{d\Phi}{dz}\right] + N^{2}(z)\Phi = 0, \quad \Phi(-h) = \Phi(0) = 0.$$
(A.1)

310 The buoyancy frequency is

$$N^2(z) = -\frac{g}{\rho_0} \frac{d\bar{\rho}}{dz},\tag{A.2}$$

where  $\bar{\rho}(z)$  is the background density profile,  $\bar{u}(z)$  is a background current in the direction of wave propagation, g is the gravitational acceleration, and  $\rho_0$  is a reference density. From Grimshaw et al. (2004) the coefficients of the rG equation (1)

for a particular mode are given by

0

315 
$$I = 2 \int_{-h}^{0} (c - \bar{u}) \Phi'^2 dz,$$
 (A.3a)

$$\alpha = I^{-1} \int_{-h}^{0} 3(c - \bar{u})^2 \Phi'^3 dz, \tag{A.3b}$$

$$\beta = I^{-1} \int_{-h}^{0} (c - \bar{u})^2 \Phi^2 dz, \tag{A.3c}$$

$$Q = c^2 I, \tag{A.3d}$$

$$\gamma = I^{-1} f^2 \int_{-h}^{0} \left( \Phi'^2 - \frac{\bar{u}'}{c - \bar{u}} \Phi \Phi' \right) dz.$$
(A.3e)

320 When  $\bar{u} = 0$ ,  $\gamma = f^2/2c$ , where f is the Coriolis frequency. The general relation for  $\gamma$  with  $\bar{u}' \neq 0$  was derived by Alias et al. (2014). The coefficient of the cubic nonlinear term is given by

$$\nu = I^{-1} \int_{-h}^{0} \left[ 3(c - \bar{u})^2 \left( 3T_z - 2\Phi'^2 \right) \Phi'^2 - \alpha^2 \Phi'^2 + \alpha(c - \bar{u}) \left[ 5\Phi'^2 - 4T_z \right] \Phi' \right] dz, \tag{A.4}$$

with the nonlinear correction to the vertical structure function, T(z), found from

$$\frac{d}{dz}\left[(c-\bar{u})^2\frac{dT}{dz}\right] + N^2T = -\alpha\frac{d}{dz}\left[(c-\bar{u})\frac{d\Phi}{dz}\right] + \frac{3}{2}\frac{d}{dz}\left[(c-\bar{u})^2\left(\frac{d\Phi}{dz}\right)^2\right]$$
(A.5)

325 with T(-h) = T(0) = 0. The solution T(z) is normalized through the addition of  $b\Phi(z)$  with b chosen so that  $T(z_{max}) = 0$ , where  $\Phi(z_{max}) = 1$ . This gives the isopycnal vertical displacement  $\xi(x, z, t) = \eta(x, t)\Phi(z) + \eta^2(x, t)T(z)$ .

In a two-layered stratification with depths  $h_1$  and  $h_2$  of the upper and lower layers, respectively, and  $\bar{u} = 0$ ,

$$c^{2} = g' \frac{h_{1}h_{2}}{h_{1}+h_{2}}, \ \alpha = \frac{3c}{2} \frac{h_{1}-h_{2}}{h_{1}h_{2}}, \ \nu = -\frac{3c}{8} \frac{\left(h_{1}^{2}+6h_{1}h_{2}+h_{2}^{2}\right)}{(h_{1}h_{2})^{2}}, \ \beta = \frac{c}{6}h_{1}h_{2}, \ \gamma = \frac{f^{2}}{2c}, \ Q = 2g'c.$$
(A.6)

Here  $g' = g\Delta\rho/\rho_0$  and  $\Delta\rho = \rho_2 - \rho_1$  is the difference in densities between the lower and upper layers.

#### 330 Author contributions. The authors contributed equally to this work.

Competing interests. There are no competing interests.

*Acknowledgements.* KRH was supported by Grant N00014-18-1-2542 from the Office of Naval Research and Grant OCE-1736698 from the National Science Foundation.

### References

- 335 Alford, M., Lien, R.-C., Simmons, H., Klymak, J., Ramp, S., Yang, Y. J., Tang, D., and Chang, M.-H., 2010: Speed and evolution of nonlinear internal waves transiting the South China Sea. J. Phys. Oceano., 121.
  - Alias, A., Grimshaw, R. H. J., and Khusnutdinova, K. R., 2014: Coupled Ostrovsky equations for internal waves in a shear flow. *Phys. Fluids*, **26**, 126603.
  - Apel, J. R., Ostrovsky, L. A., Stepanyants, Y. A., and Lynch, J. F., 2007: Internal solitons in the ocean and their effect on underwater sound.

- Farmer, D., Li, Q., and Park, J.-H., 2009: Internal wave observations in the South China Sea: the role of rotation and nonlinearity. *Atmos.-Ocean*, **47**, 267–280.
- Grimshaw, R., Guo, C., Helfrich, K., and Vlasenko, V., 2014: Combined effect of rotation and topography on shoaling oceanic internal solitary waves. *J. Phys. Oceano.*, **44**, 1116–1132.
- 345 Grimshaw, R., He, J.-M., and Ostrovsky, L., 1998a: Terminal damping of a solitary wave due to radiation in rotational systems. *Stud. Appl. Math*, 101, 197–210.
  - Grimshaw, R., Helfrich, K., and Johnson, E. R., 2012: The reduced Ostrovsky equation: Integrability and breaking. *Stud. Appl. Math.*, **129**, 414–436.

Grimshaw, R., Pelinovsky, E., and Talipova, T., 1998b: Solitary wave transformation due to a change in polarity. Stud. Appl. Math, 101,

350

365

357-388.

- Grimshaw, R., Pelinovsky, E., and Talipova, T., 1999: Solitary wave transformation in a medium with sign-variable quadratic nonlinearity and cubic nonlinearity. *Physica D*, **132**, 40–62.
- Grimshaw, R., Pelinovsky, E., Talipova, T., and Kurkin, A., 2004: Simulation of the transformation of internal solitary waves on oceanic shelves. *J. Phys. Oceano.*, **34**, 2774–2791.
- Grimshaw, R. H. J., and Helfrich, K. R., 2008: Long-time solutions of the Ostrovsky equation. *Studies Appl. Math.*, 121, 71–88.
  Helfrich, K. R., 2007: Decay and return of internal solitary waves with rotation. *Phys. Fluids.*, 19, 026601.
  Helfrich, K. R., and Grimshaw, R., 2008: Nonlinear disintegration of the internal tide. *J. Phys. Oceano.*, 38, 686–701.
  Holloway, P. E., Pelinovsky, E., and Talipova, T., 1999: A generalized Korteweg-de Vries model of internal tide transformation in the coastal ocean. *J. Geophys. Res.*, 104, 18333–18350.
- 360 Jackson, C., 2004: An Atlas of Internal Solitary-like Waves and Their Properties. 2nd ed., 560 pp., Global Ocean Assoc., Alexandria, Va. (Available at http://www.internalwaveatlas.com).
  - Karczewska, A., and Rozmej, P., 2020: Can simple KdV-type equations be derived for shallow water problem with bottom bathymetry? *Comm. Nonlin. Sci. Numer. Sim.*, **82**, 105073.

- Obregon, M., Raj, N., and Stepanyants, Y., 2018: Adiabatic decay of internal solitons due to Earth's rotation within the framework of the Gardner-Ostrovsky equation. *Chaos*, **28**, 033106.
- Ostrovsky, L., 1978: Nonlinear internal waves in a rotating ocean. Oceanology, 18, 119–1125.

Ostrovsky, L., and Grue, J., 2003: Evolution equations for strongly nonlinear internal waves. Phys. Fluids, 15, 2934–2948.

<sup>340</sup> J. Acoust. Soc. Am., 121, 695–722.

Nakoulima, O., Zahibo, N., Pelinovsky, E., Talipova, T., Sulnyaev, A., and Kurkin, A., 2004: Analytical and numerical studies of the variablecoefficient Gardner equation. *Appl. Math. Comp.*, **152**, 449–471.

370 Ostrovsky, L., and Helfrich, K. R., 2019: Some new aspects of the joint effect of rotation and topography on internal solitary waves. J. Phys. Oceano., 49, 1639–1649.

Ostrovsky, L., and Irisov, V. G., 2017: Strongly nonlinear internal solitons: Models and applications. J. Geophys. Res., 122, 3907–3916.

- Pelinovsky, E. N., and Grimshaw, R., 1997: Structural transformation of eigenvalues for a perturbed algebraic soliton potential. *Physics Letters A*, **229**, 165–172.
- 375 Pelinovsky, E. N., Polukhina, O. E., and Lamb, K. 2000: Nonlinear internal waves in the ocean stratified in density and current. *Marine Phys.*, 40, 757–766.
  - Stanton, T. P., and Ostrovsky, L. A., 1998: Observations of highly nonlinear internal solitons over the continental shelf. *Geophys. Res. Let.*, 25, 2695–2698.

Stepanyants, Yu., 2019: The effects of interplay between the rotation and shoaling for a solitary wave on variable topography. Stud. Appl.

- 380 *Math.*, 1–22. https://doi.org/10.1111/sapm.12255
  - Talipova, T. G., Kurkina, O. E., Rouvinskaya, E. A., and Pelinovsky, E. N., 2015: Propagation of solitary internal waves in two-layer ocean of variable depth. *Izvestiya, Atmos. Oceanic Phys.*, **51**, 89–97.
  - Whitfield, A. J., and Johnson, E. R., 2015: Wave-packet formation at the zero-dispersion point in the Gardner-Ostrovsky equation. *Phys. Rev.*, **91**, 051201.
- 385 Zhao, Z., and Alford, M., 2006: Source and propagation of nonlinear internal waves in the northeastern South China Sea. J. Geophys. Res., 111, C11012.